

SECOND EDITION 2023

ANTIMICROBIAL GUIDELINE

HOSPITAL RAJA PERMAISURI BAINUN

ANTIMICROBIAL GUIDELINE HOSPITAL RAJA PERMAISURI BAINUN SECOND EDITION 2023

Published by:
Antimicrobial Stewardship (AMS) Team, Hospital Raja Permaisuri Bainun

This guide can be downloaded from the
Hospital Raja Permaisuri Bainun website: hrpb.moh.gov.my

FOREWORDS

Antimicrobial resistance in disease pathogens has become a matter of great public health concern globally including in our country. The factors responsible for this are widespread use of all the antimicrobials across the country meant for human as well as animal consumption. Inappropriate selection of antimicrobials may lead to antimicrobial resistance, which is fast becoming a growing public health threat worldwide. Detecting, preventing and controlling antibiotic resistance requires sustained effort and engagement from all healthcare professionals.

I am happy to note that the Antimicrobial Stewardship Team of HRPB in collaboration with various departments in the hospital has produced the second antimicrobial guideline HRPB adapted to the local practice, thereby reducing the development of antimicrobial resistance. The guideline is important to provide guidance on the selection of antimicrobials to healthcare professionals.

I congratulate all the working members from various departments for their contribution in preparing this guideline. I hope that the healthcare professionals in HRPB would use these guidelines to the maximum for the benefit of the patients.

Thank you.



DR. MEGAT ISKANDAR BIN MEGAT ABDUL HAMID (P.M.P)
Director of Hospital Raja Permaisuri Bainun (HRPB)

FOREWORDS

As we all know, the use of antimicrobials in one person has the potential to affect the drug's efficacy in others, making the overuse of antimicrobials a public health problem not just hospital wide, nationwide but also a global wide issue. Inadequate infection prevention and control practice can further worsen the spread of antimicrobial resistance.

The practice of appropriate and judicious utilisation of antimicrobials is of utmost importance in order to improve patient outcomes and to reverse or slow down antimicrobial resistance. Hence, antimicrobial guidelines have always been used to provide guidance to healthcare personnels in the management of various infections. Our first *HRPB Antimicrobial Guideline 2017* was published on the HRPB website. Now I am indeed honoured to introduce this new updated edition of *HRPB Antimicrobial Guideline 2023*. This edition sees some new and exciting changes done to the format and the content, making it more user friendly especially to the junior doctors.

This guideline was reviewed by relevant specialists from respective departments in HRPB. My heartfelt thank you goes to those who are involved and work hard to update this document despite their busy schedules.

This endeavour would not be possible without the tremendous efforts by our editorial team who worked diligently, from organising everything up to the final editing. I salute you all! I would like to thank the continuous support and trust from hospital top management, head of clinical departments and pharmacy.

Last but not least, I sincerely hope that our clinicians will find this updated guideline useful in their daily practice. Let's use antimicrobial wisely and prevent antimicrobial resistance together!

Thank you.



DATO DR. KER HONG BEE

Consultant Infectious Disease Physician
Department of Medicine
Hospital Raja Permaisuri Bainun (HRPB)

FOREWORDS

Overuse and misuse of antimicrobials have been associated with more rapid emergence of antimicrobial resistance. Therefore, promoting the appropriate use of antimicrobials through an updated guideline is important to reduce antimicrobial resistance and infections caused by multidrug resistant organisms.

The second edition of the *HRPB Antimicrobial Guideline 2023* has been developed to suit the local practice. This guideline is adapted from the *National Antimicrobial Guideline 2019*. I hope it serves as a guiding tool to provide relevant information on the rational use of antibiotics. This guideline is intended to be used by all the healthcare professionals in HRPB.

I want to congratulate all who are involved for their commendable efforts in reviewing and updating this guideline. I hope this updated guideline will be implemented effectively to ensure rational use of antimicrobials, thereby reducing antimicrobial resistance.

Thank you.

A handwritten signature in black ink, consisting of a large, stylized 'N' followed by a checkmark-like flourish.

NORMI BT KAMARUZAMAN

Head of Pharmacy Department
Hospital Raja Permaisuri Bainun (HRPB)

WORKING COMMITTEE

Advisors

Dr. Megat Iskandar bin Megat Abdul Hamid	Hospital Director
Ms. Normi bt Kamaruzaman	Chief Pharmacist

Infectious Disease Physicians

Dato' Dr. Ker Hong Bee
Dr. Cheng Joo Thye
Dr. Steven Lim Chee Loon
Dr. Marsilla Mariaty binti Marzukie

Infectious Disease Paediatrician

Dr. Haema Shunmugarajoo

Clinical Microbiologists

Dr. Zahrul Laili binti Abd Hadi
Dr. Fong Kang Chyi @ Foong Kang Chyi

Pharmacists

Dr. Doris a/p George Visuvasam	Mr. Muhammad Azuan bin Azman
Ms. Ros Sakinah Binti Kamaludin	Mr. Ahmad Ayzat bin Ahmad Jaafar
Ms. Lee Pooi Mun	Ms. Lai Qian Xin
Mr. Ho Chee Wah	Ms. Tan Rou Wei
Ms. Wong Hong Yean	Mr. Leow Hon Lunn
Ms. Voon Bee Peng	Ms. Intan Nurul'Ain bt Ismail
Ms. Siti Nur Sharida binti Abd Kadir	Ms. Kwek Lih Ling
Mr. Cheah Meng Fei	Ms. Amutha Vaani a/p Vinthaikumaran
Ms. Manimegahlai a/p Selvaratnam	Ms. Ghaneshwari a/p Ramamuthie
Ms. Nurulhusna Syuhaidah bt Yusof	Ms. K. Oganeshwary a/p Kobi
Ms. Thanushiapriya a/p Selva Raj	Mr. Sathesh Kumar a/l Ramayah
Ms. Lee Pei Jia	Ms. Roseanna Prescilla a/p Ramakrishnan
Ms. Kong Pui Chin	Ms. Low Yuet Man
Ms. Liew Wei Mun	Ms. Malarvili a/p Manika Wasagam
Ms. Choong Lai Yuan	
Ms. Kuginesh a/p Ganapathy	
Ms. Chow Li Wei	
Ms. Chan Pooi Yee	
Mr. Loo Yew Win	
Mr. Nicholas Lee Chor Teng	
Ms. Wan Alia Amalina bt Adenan	
Ms. Hemalathah a/p Paramalingam	
Ms. Lim Wan Lin	
Ms. Madhuri a/p G. Sabapathy	
Ms. Nur Atikah binti Basharuddin	

CONTRIBUTORS

Surgery

Ms. (Dr.) Elaine Ng Hui Been
Mr. (Dr.) L Sivanewaran Lechmiannandan
Ms. (Dr.) Nurdaliza M Badaruddin

Neurosurgery

Ms. (Dr.) Neoh Yee Yik

Plastic and Reconstructive Surgery

Mr. (Dr.) Jeremiah See Un Chang

Obstetrics and Gynaecology

Dr. Kaamini a/p Planisamy
Dr. S. Hamsavahini a/p Sorano @ Soranam

Orthopaedics

Mr. (Dr.) Choo Tze Yong
Mr. (Dr.) Yap Khoo Sim

Otorhinolaryngology

Dr. Maithrea Suresh Narayanan
Dr. Benjamin Goh Hong Beng

Ophthalmology

Dr. Umami Faradiana binti Abdul Rahim

Dermatology

Dr. Madiha binti Muhamad Sarkan

Nephrology

Dr. Loh Chek Loong
Dr. Thong Kah Mean

Neurology

Dr. Cheah Chun Fai

Respiratory

Dr. Kumaresh Raj a/l Lachmanan

Oncology

Dr. Angel Kwan Khor Nee

Haematology

Dr. Kamini a/p Kirubamoorthy

Cardiovascular Infections

Dr. Kwan Yew Fung

Gastrointestinal Infections

Dr. Tan Phei Oon

Oral Health

Dr. Yuen Kar Mun

Paediatrics

Dr. Chew Eng Lai

Paediatric Dentistry

Dr. Sumathy Perumal

Microbiology

Ms. Azlina binti Yahya
Ms. Husna binti Khairam

TABLE OF CONTENTS

Contents	Page
SECTION A: ADULTS	
Cardiovascular Infections	10
Central Nervous Infections	19
Gastrointestinal Infections	28
Infections in Immunocompromised Patients:	
A. Haematology/Oncology	37
B. Human Immunodeficiency Virus (HIV)	44
Obstetrics and Gynaecological Infections	57
Ocular Infections	63
Oral/Dental Infections	73
Otorhinolaryngology Infections	82
Respiratory Infections	87
Sexually Transmitted Infections	92
Skin and Soft Tissue Infections	102
Surgical Infections	118
Tropical Infections	135
Urinary Tract Infections	144
Infections In Patients On Long-Term Renal Replacement Therapy	152
Chemoprophylaxis	
A. Surgical	157
B. Non-surgical	174
 SECTION B: PAEDIATRICS	
Cardiovascular Infections	177
Central Nervous Infections	182
Gastrointestinal Infections	185
Infections In Immunocompromised Patients	190
Neonatal Infections	193
Oral/Dental Infections	199
Otorhinolaryngology Infections	201
Respiratory Infections	203
Skin and Soft Tissue Infections	205
Surgical Infections	208
Tropical Infections	211
Tuberculosis in Children	217

Urinary Tract Infections	219
Vascular Infections	221
Chemoprophylaxis	
C. Surgical	224
D. Non-surgical	229
APPENDICES	
Appendix 1: Antibiotic Resistance Data	239
Appendix 2: Antibiotic Utilisation Data	250
Appendix 3: Antimicrobial Dosages in Patients with Impaired Renal Function (Adults)	253
Appendix 4: Clinical Pharmacokinetics Guidelines (Aminoglycosides and Vancomycin)	266

(SECTION A: ADULTS)

CARDIOVASCULAR INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
INFECTIVE ENDOCARDITIS			
Empirical Treatment			
Community-acquired native valves	Benzylpenicillin 18MU/day IV q4-6h OR Ampicillin 12g/day IV q4-6h PLUS Gentamicin 3mg/kg IV q24h PLUS/ MINUS *Cloxacillin 12g/day IV q4-6h	Penicillin Allergy: Vancomycin 25-30mg/kg IV loading dose, then 15-20mg/kg/dose IV q8-12h (actual body weight, not to exceed 2g/dose) PLUS Gentamicin 3mg/kg IV q24h	Treatment can be modified once the blood result is known. *For patients with suspected <i>S. aureus</i> infections (such as IVDU or patients with prosthesis) and acute presentation.
Early PVE (<12 months post-surgery) or nosocomial and non-nosocomial healthcare associated endocarditis	Vancomycin 25-30mg/kg IV loading dose, then 15-20mg/kg/dose IV q8-12h (actual body weight, not to exceed 2g/dose) PLUS Gentamicin 3mg/kg IV q24h PLUS/ MINUS *Rifampicin 300-450mg PO q12h PLUS/MINUS **Cefepime 2g IV q8h		*Rifampicin is only recommended for PVE and it should be started 3-5 days later than vancomycin and gentamicin. **Cefepime is indicated if local epidemiology suggests for non-HACEK Gram-negative rod infections (such as <i>Pseudomonas</i>).
Late PVE (> 12months)	Ampicillin 12g/day IV q4-6hr PLUS Gentamicin 3mg/kg IV q24h	Penicillin Allergy:	*For patients with suspected <i>S. aureus</i> infections (such as IVDU or

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	PLUS/ MINUS *Cloxacillin 12g/day IV q4-6h	Vancomycin 25-30mg/kg IV loading dose, then 15-20mg/kg/dose IV q8-12h (actual body weight, not to exceed 2g/dose) PLUS Gentamicin 3mg/kg IV q24h	patients with prosthesis) and acute presentation.
Definitive Treatment <i>(Viridans streptococci & Streptococcus bovis: It is recommended MIC estimation is done for these isolates to facilitate management)</i>			
Native and Prosthetic Valves MIC: < 0.125mg/L Penicillin-susceptible viridans streptococci & <i>Streptococcus bovis</i>	Benzylpenicillin 3MU IV q4-6h (or 12-18MU/day as continuous infusion)	Ampicillin 2g IV q4h OR Ceftriaxone 2g IV/IM q24h Penicillin Allergy: Vancomycin 25-30mg/kg IV loading dose, then 15-20mg/kg/dose IV q8-12h (actual body weight, not to exceed 2g/dose)	Duration of therapy: 4 weeks (native valves) 6 weeks (prosthetic valves) Penicillin-susceptible viridans streptococci, monotherapy with benzylpenicillin, ampicillin or ceftriaxone is adequate.
Native Valves and Prosthetic Valves MIC > 0.125mg/L- 2mg/L Penicillin-relatively resistant viridans streptococci & <i>Streptococcus bovis</i>	*Benzylpenicillin 4MU IV q4h or 24MU IV as continuous infusion PLUS **Gentamicin 3mg/kg IM/IV q24h	*Ceftriaxone 2g IV/IM q24h PLUS **Gentamicin 3mg/kg IV/IM q24h Penicillin Allergy (immediate-type): *Vancomycin 25-30mg/kg IV loading dose, then 15-20mg/kg/dose IV q8-12h (actual body weight, not to exceed 2g/dose) PLUS **Gentamicin 3mg/kg IV/IM q24h	*Duration of therapy: 4 weeks (native valves) 6 weeks (prosthetic valves) **Penicillin-relatively resistant viridans streptococci, gentamicin has to be added to regime; 2 weeks (native valves) 6 weeks (prosthetic valves) **Gentamicin pre-dose (trough): <1mg/L

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Native Valves & Prosthetic Valves MIC > 2mg/L Penicillin-relatively resistant viridans streptococci & <i>Streptococcus bovis</i>	Treat as resistant enterococcal endocarditis		
Nutritionally variant <i>Streptococci</i> (NVS) <i>Abiotrophia defectiva</i> and <i>Granulicatella species</i> , both formerly known as NVS	Ampicillin 2g IV q4h for 6 weeks PLUS Gentamicin 1mg/kg IV q8h for 2 weeks OR Benzylpenicillin 4MU IV q4h (or 24MU IV as continuous infusion) for 6 weeks PLUS Gentamicin 1mg/kg IV q8h for 2 weeks	Ceftriaxone 2g IV/IM q24h for 6 weeks PLUS Gentamicin 1mg/kg IV q8h for 2 weeks Penicillin Allergy (immediate-type): Vancomycin 25-30mg/kg IV loading dose, then 15-20mg/kg/dose IV q8-12h (actual body weight, not to exceed 2g/dose) for 6 weeks	NVS are fastidious and slow growing making it technically difficult to determine antimicrobial susceptibility.
Definitive Treatment (<i>Enterococcus</i> species)			
Native and Prosthetic Valves (sensitive to gentamicin) MIC ≤8mg/L Fully penicillin-susceptible strains	Ampicillin 2g IV q4h PLUS *Gentamicin 1mg/kg IM/IV q8h		Duration of symptoms < 3 months and native valves: Ampicillin duration - 4 weeks Gentamicin duration - 2 weeks Duration of symptoms > 3 months or prosthetic valves: Ampicillin duration - 6 weeks Gentamicin duration - 6 weeks *To maximize synergistic effect, administer gentamicin at the same

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			time or temporally close to ampicillin/penicillin.
Native and Prosthetic Valves (sensitive to gentamicin) (renal impairment and elderly)	Ampicillin 2g IV q4h for 6 weeks PLUS Ceftriaxone 2g IV q12h for 6 weeks		
Native and Prosthetic Valves Sensitive to penicillin and vancomycin but high level resistance to gentamicin (MIC > 500mg/L)	Ampicillin 2g IV q4h for 6 weeks PLUS *Ceftriaxone 2g IV q12h for 6 weeks		*Ceftriaxone should not be used alone for <i>enterococcus</i> infection, as they are intrinsically resistant. This combination is not active against <i>E. faecium</i> .
Native and Prosthetic Valves Resistant to penicillin and susceptible to aminoglycosides and vancomycin	Vancomycin 25-30mg/kg IV loading dose, then 15-20mg/kg/dose IV q8-12h (actual body weight, not to exceed 2g/dose) for 6 weeks PLUS *Gentamicin 1mg/kg IV/IM q8h for 6 weeks		*Gentamicin pre-dose (trough): <1mg/L
Definitive Treatment (<i>Staphylococcus aureus</i>)			
Native Valves Methicillin-susceptible staphylococci (MSSA)	Left sided endocarditis or complicated right sided endocarditis: Cloxacillin 2g IV q4h for 4- 6 weeks	Both left-sided and right sided: Immediate-type penicillin allergy: Vancomycin 25-30mg/kg IV loading dose, then 15-20mg/kg/dose IV q8-12h (actual body weight, not to exceed 2g/dose) for 4- 6 weeks Non-immediate type hypersensitivity:	Vancomycin is inferior to cloxacillin for treatment of MSSA.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
		Cefazolin 2g IV q8hr for 4- 6weeks	
Native Valves Methicillin-susceptible staphylococci (MSSA)	Right sided endocarditis (tricuspid valve) in uncomplicated endocarditis: Cloxacillin 2g IV q4h for 2- 4 weeks		2-week regime is sufficient provided the patient fulfils all the following criteria: ● MSSA ● Good response to treatment ● Absence of metastatic sites of infection or empyema ● Absence of cardiac and extracardiac complications ● Absence of associated prosthetic valve or left-sided valve infection ● <20mm vegetation ● Absence of severe immuno-suppression (<200 CD4 cells/mL) with or without AIDS
Native Valves Methicillin-resistant staphylococci (MRSA)	Vancomycin 25-30mg/kg IV loading dose, then 15-20mg/kg/dose IV q8-12h (actual body weight, not to exceed 2g/dose) for 4-6 weeks	*Daptomycin 10mg/kg IV q24h for 4-6 weeks	*Daptomycin is superior to vancomycin for MRSA bacteremia with vancomycin MIC > 1mg/L *Daptomycin requires special medicine approval
Prosthetic Valves Methicillin-susceptible staphylococci (MSSA)	Cloxacillin 2g IV q4h for ≥ 6 weeks PLUS Rifampicin 300-450mg PO q12h for ≥ 6 weeks PLUS Gentamicin 1mg/kg IM/IV q8h for 2 weeks	Regimen for β-lactam allergic patients: Immediate type hypersensitivity to penicillin: Vancomycin 25-30mg/kg IV loading dose. then 15-20mg/kg/dose IV q8-12h (actual	Rifampicin has better penetration, however, it should be started after 3-5 days of effective initial cloxacillin therapy and/or once the bacteraemia has been cleared to avoid resistance.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
		body weight, not to exceed 2g/dose) for 4- 6 weeks OR For non-immediate type hypersensitivity: Cefazolin 2g IV q8h for 4- 6 weeks	Gentamicin can be given in a single daily dose in order to reduce renal toxicity
Prosthetic Valves Methicillin-resistant staphylococci (MRSA)	Vancomycin 25-30mg/kg IV loading dose, then 15-20mg/kg/dose IV q8-12h (actual body weight, not to exceed 2g/dose) for ≥ 6 weeks PLUS Gentamicin 1mg/kg IM/IV q8h for 2 weeks PLUS *Rifampicin 300-450mg PO q12h for ≥ 6 weeks		*Rifampicin has better penetration, however, it should be started after 3-5 days of effective initial vancomycin therapy and/or once the bacteraemia has been cleared to avoid resistance.
Definitive Treatment HACEK Microorganisms (<i>Haemophilus parainfluenzae</i> , <i>Aggregatibacter aphrophilus</i> , <i>Aggregatibacter actinomycetemcomitans</i> , <i>Cardiobacterium hominis</i> , <i>Eikenella corrodens</i> , and <i>Kingella kingae</i>)			
Native and Prosthetic Valves	Ceftriaxone 2g IV/IM q24h for 4 weeks (native valve) or 6 weeks (prosthetic valve)	Ampicillin/sulbactam 3g IV q6h for 4 weeks (native valve) or 6 weeks (prosthetic valve) OR *Ciprofloxacin 400mg IV q12h or 500mg PO q12h for 4 weeks (native valve) or 6 weeks (prosthetic valve)	*Fluoroquinolones can be switched to appropriate oral doses if patients can tolerate oral medications.
Definitive Treatment			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Non-HACEK Gram-negative microorganisms (eg. <i>Pseudomonas aeruginosa</i> , <i>Escherichia coli</i> and <i>Salmonella</i>)			
<i>Pseudomonas</i> Endocarditis	Piperacillin-tazobactam 4.5g IV q6h PLUS Gentamicin 5-7mg/kg IV q24h for 6 weeks OR Piperacillin-tazobactam 4.5g IV q6h PLUS Amikacin 15mg/kg IV q24h for 6 weeks	Cefepime 2g IV q8h OR Ceftazidime 2g IV q8h PLUS Gentamicin 5-7mg/kg IV q24h OR Amikacin 15mg/kg IV q24h OR Cefepime 2g IV q8h OR Ceftazidime 2g IV q8h PLUS *Ciprofloxacin 400mg IV q8h	Choice of antimicrobials depends on antimicrobial susceptibility patterns. Often needs to be combined with cardiac surgery. *Fluoroquinolones can be switched to appropriate oral doses if patients can tolerate oral medications.
Definitive Treatment: <i>Candida</i>			
Native and Prosthetic Valves	Amphotericin B deoxycholate 0.6-1mg/kg IV q24h OR Lipid formulation Amphotericin B 3-5mg/kg IV q24h PLUS/ MINUS *Flucytosine 25mg/kg PO q6h for at least 6 weeks after surgery	Micafungin 150mg IV q24h OR Caspofungin 150mg q24h OR Anidulafungin 200mg IV q24h for at least 6 weeks after surgery	Valve replacement is mandatory. Continue therapy for 6 weeks after replacement or longer in patients with perivalvular abscess. If prosthetic valve cannot be replaced, lifelong suppressive therapy with fluconazole 400mg (6mg/kg) daily is recommended.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			Duration of therapy depends on the patient's response and surgical intervention and should be referred to an ID physician. *For synergistic effect, causes dose related marrow toxicity, avoid using in patients with renal failure.
Definitive Treatment: Other microorganisms			
<i>Brucella spp.</i>	Doxycycline 100mg PO q12h PLUS Rifampicin 300-600mg PO daily PLUS *Streptomycin 15mg/kg IM q24h OR *Gentamicin 5mg/kg IV 24h		Duration of therapy: 3-6 months * For first 2-4 weeks only
<i>C. burnetii</i> (agent of Q fever)	Doxycycline 100mg PO q12h PLUS Hydroxychloroquine 600mg PO q24h or 200mg PO q8h		Duration: 18-24 months based on clinical and serological response
<i>Bartonella spp.</i>	Doxycycline 100mg PO q12h for 6 weeks PLUS Gentamicin 3mg/kg IV q24h for 2 weeks		
Therapy for Culture-Negative Endocarditis - Consultation with an infectious disease specialist needed			

Footnotes for antibiotic treatment of endocarditis:

1. Vancomycin : aim for serum trough level of 15-20 mg/for both adults and pediatrics. Dose should be adjusted in patients with renal impairment.
2. For patients on gentamicin:
 - Monitor gentamicin level and renal function weekly. There should be a low threshold for stopping gentamicin in patients with deteriorating renal function or other signs of toxicity.
 - Aim for peak serum level of 3-5mcg/mL & pre-dose (trough) serum level of 1mcg/mL when 2-3 divided doses are given.
 - Consider biweekly clinical screening for ototoxicity: Check baseline visual acuity using a Snella pocket card, to screen for ototoxicity, have patient shake head and reread the card, consider formal audiology test if patient loses 2 lines of visual acuity.
 - For obese patients use ideal body weight
 - For high level gentamicin resistance, ampicillin/sulbactam or vancomycin continued for ≥ 6 weeks, .Refer ID physician.
3. Rifampicin should be combined with another effective antistaphylococcal drug to minimize risk of resistance. Rifampicin increases hepatic clearance of warfarin & other drugs

References:

1. Clinical Practice Guidelines for the Prevention, Diagnosis and Management of Infective Endocarditis 2017, National Heart Association of Malaysia and the Academy of Medicine Malaysia
2. National Antibiotic Guideline 2019 - Cardiovascular Infections for Adults
3. Gilbert Habib et al. ESC Scientific Document Group, 2015 ESC Guidelines for the management of infective endocarditis: The Task Force for the Management of Infective Endocarditis of the European Society of Cardiology (ESC)
Endorsed by: European Association for Cardio-Thoracic Surgery (EACTS), the European Association of Nuclear Medicine (EANM), *European Heart Journal*, Volume 36, Issue 44, 21 November 2015

CENTRAL NERVOUS INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Meningitis (acute): Empirical Therapy			
Common organisms: <i>Streptococcus pneumoniae</i> <i>Haemophilus influenzae</i> <i>Neisseria meningitidis</i> Other organisms: Gram-negative rods	Ceftriaxone 2g IV q12h OR Cefotaxime 2g IV q6h	Alternative ONLY for immunocompromised host: Meropenem 2g IV q8h	Antibiotic should not be delayed if lumbar puncture (LP) is delayed by radiological investigation. If no organism is isolated from CSF C&S but LP is suggestive of bacterial meningitis and patient is responding, continue antibiotics for 14 days Dexamethasone 10mg IV q6h is recommended 15 to 20 minutes before or at the time of first dose of antibiotics. Continue for 4 days if the Gram stain and/or cultures consistent with <i>S. pneumoniae</i>. Discontinue if not <i>Streptococcus pneumonia</i> or if bacterial meningitis is subsequently thought not to be present. Incidence of listeriosis increases in people > 60 years of age, immunosuppressed & pregnancy. Consider empirical cover for this organism especially if the course of disease is indolent or there is

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			epidemiological risk (refer section on treatment of listeriosis) Duration: 10-14 days
Meningitis (acute): Causative organism isolated			
<i>Haemophilus influenzae</i> (Gram-negative bacilli)	Ceftriaxone 2g IV q12h OR Cefotaxime 2g IV q6h	Cefepime 2g IV q8h. If organism is susceptible and patient is allergic to cephalosporins: Ciprofloxacin 400mg IV q8h	Duration: 7-10 days
<i>Streptococcus pneumoniae</i> (Gram-positive cocci)	Penicillin-sensitive strains (MIC to Penicillin < 0.12 mcg/mL) Benzylpenicillin 4MU IV q4h Penicillin-resistant strains (MIC to Penicillin ≥0.12 mcg/mL) Ceftriaxone 2g IV q12h OR Cefotaxime 2g IV q6h Cephalosporin-resistant strains (MIC to Cephalosporin ≥2 mcg/mL): Vancomycin 25-30mg/kg loading dose then 15-20mg/kg IV q8-12h; not to exceed 2g per dose PLUS Ceftriaxone 2g IV q12h	Penicillin-resistant strain (MIC to Penicillin ≥0.12 mcg/ml) Cefepime 2g IV q8h OR Meropenem 2g IV q8h	All attempts should be made to ascertain the MIC of isolated pneumococcus. Ceftriaxone or cefotaxime should be de-escalated to benzylpenicillin once the MIC result has been confirmed. Duration: 10-14 days

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	OR Cefotaxime 2g IV q6h		
<i>Neisseria meningitidis</i> (Gram-negative diplococci)	If MIC to Penicillin < 0.1 mcg/mL: Benzylpenicillin 4MU IV q4h If MIC to penicillin is > 0.1 mcg/mL: Ceftriaxone 2g IV q12h OR Cefotaxime 2g IV q6h		Duration: 5-7 days
Prophylaxis for household and close contacts of meningococcal meningitis cases	Age > 15 years: *Ciprofloxacin 500mg PO as single dose OR *Rifampicin 600mg PO q12h for 2 days (4 doses)	Ceftriaxone 250mg IM as single dose (especially in pregnancy and lactating mothers) OR Azithromycin 500mg PO as single dose	Close contacts are defined as those individuals who have had contact for > 8 hours and within 1 metre of the index case. Individuals who were in contact with oropharyngeal secretions of the index case in the last 7 days before onset of symptoms up to 24 hours after appropriate antibiotics should also receive chemoprophylaxis. For index cases who received only benzylpenicillin as therapy, chemoprophylaxis should also be given upon discharge to eliminate nasopharyngeal carriage. *Not recommended in pregnant women

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Listeriosis	Ampicillin 2g IV q4h OR Benzylpenicillin 4MU IV q4h PLUS/MINUS Gentamicin 5mg/kg/day IV in 3 divided doses	Trimethoprim/ sulfamethoxazole 10-20mg/kg/day [based on the TMP component] IV q6-12h OR Meropenem 2g IV q8h	Duration of treatment is 3 weeks depending on clinical response. May be longer in immunocompromised host. Gentamicin is given until symptoms improve (minimum of 1 week)
Brain abscess/subdural empyema: Empirical Therapy			
Common organisms: <i>Streptococcus</i> <i>Staphylococcus</i> Gram-negative bacilli Anaerobes Source: 1. Dental or Oral 2. Paranasal Sinus or Orogenic 3. Hematogenous spread or penetrating trauma (community acquired) Source: Hematogenous spread (hospital acquired) or post neurosurgical operation	Ceftriaxone 2g IV q12h PLUS Metronidazole 500mg IV q8h Ceftriaxone 2g IV q12h If patient has history of MRSA infections, consider adding: Vancomycin 25-30mg/kg loading dose then 15-20mg/kg IV q8-12h; not to exceed 2g per dose.		Surgical intervention: excision or aspiration of the abscess is treatment of choice if the size of the abscess is more than 2.5cm Blood cultures, WBC and CRP should be taken when a diagnosis of brain abscess is suspected. Other screenings needed are ECHO (for patient with heart problem-to see possibilities of vegetation) and scope (for paranasal sinus)

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Brain abscess/ subdural empyema: Causative organism isolated			
<i>Streptococcus milleri</i>	Benzylpenicillin 4MU IV q6h	Cefotaxime 2g IV q6h OR Vancomycin 25-30mg/kg loading dose then 15-20mg/kg IV q8-12h; not to exceed 2g per dose	Duration to be determined by clinical response (usually 4-8 weeks with IV therapy for 2 weeks minimum depending on whether surgical drainage done, clinical and radiological response).
<i>Staphylococcus aureus</i> (Methicillin-sensitive)	Cloxacillin 2g IV q4h OR Cefazolin 2g IV q8h	*Vancomycin 25-30mg/kg loading dose then 15-20mg/kg IV q8-12h; not to exceed 2g per dose OR If no bacteremia: Trimethoprim/sulfamethoxazole 10mg/kg/day [based on TMP component] IV q6-12h	
Enterobacterales	Cefotaxime 2g IV q6h	Meropenem 2g IV q8h	Duration of Treatment: -3-4 weeks if abscess has been excised -4-6 weeks if abscess has been aspirated
<i>Pseudomonas aeruginosa</i>	Ceftazidime 2g IV q8h	Meropenem 2g IV q8h	(Minimum of 4 weeks if abscess treated with antibiotic alone and CT showing resolution of ring enhancing lesions & good clinical progress) Patient can be switched onto oral therapy when: 1. CRP falls to normal levels. 2. No signs of fever

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			3. Appropriate oral antibiotics are available 4. Patient can tolerate oral antibiotics
Spinal Epidural abscess Common organisms: <i>Streptococci</i> <i>Staphylococcus</i> Gram-negative bacilli	Cloxacillin 2g IV q4h OR *Vancomycin 25-30mg/kg loading dose then 15mg/kg IV q8-12h; not to exceed 2g per dose PLUS Gentamicin 4-7mg/kg/day IV in 3 divided doses OR **Ceftriaxone 2g IV q12h OR **Cefotaxime 2g IV q4-6h		Source control is strongly recommended. Duration to be determined by clinical response (usually 2-6 weeks with IV therapy for 2 weeks minimum, followed by oral depending on whether surgical drainage done, clinical and radiological response). *Vancomycin is indicated when suspecting MRSA or allergy to Cloxacillin **Third-generation cephalosporin indicated when gentamicin is contraindicated.
Viral encephalitis Common organisms: Herpes simplex Varicella zoster	*Acyclovir 10mg/kg IV q8h		*Consider using ideal body weight in obese patients. Duration: 14-21 days
Meningitis (Chronic)			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Tuberculous meningitis <i>(Mycobacterium tuberculosis)</i>	Intensive 2 months S/EHRZ and 10 months HR Isoniazid (H) 5 (4-6) mg/kg/day PO (max: 300 mg/day) PLUS Rifampicin (R) 10 (8-12) mg/kg/day PO (max: 600mg/day) PLUS Pyrazinamide (Z) 25 (20-30)mg/kg/day PO (max: 2000 mg/day) PLUS Streptomycin (S) 15 (12-18) mg/kg/day IM (max: 1000mg/day) OR Ethambutol (E) 15 (15-20)mg/kg/day PO (max: 1600mg/day) Pyridoxine 10-50mg PO q24h needs to be prescribed together with Isoniazid	In HIV patients: Recommendations for the treatment of TB in HIV-infected adults are identical to those for HIV-uninfected adults when the disease is caused by organisms that are known or presumed to be susceptible to the first-line drugs. Daily dosing is recommended rather than intermittent dosing. Rifampicin is not recommended in combination with all protease inhibitors (PIs) and rifabutin should be used with PI-based ART for HIV-TB co-infected adults.	Infection in HIV patients: Add IV Dexamethasone 12-16 mg daily in divided doses for 6 weeks in tapering doses, (intravenously initially then switch to oral when safe to do so). Alternatively, prednisolone 30-40mg/day PO in tapering doses for 6 weeks. Treatment is continued for 12 months. Refer to Clinical Practice Guidelines on Management of Tuberculosis (4th edition).
Cryptococcal meningitis <i>Cryptococcus neoformans</i> (non-HIV, non-transplant patients)	Induction Therapy: Amphotericin B 0.7-1.0mg/kg/day IV q24h PLUS 5-Flucytosine 100mg/kg/day PO q6h OR Fluconazole 1200mg PO q24h.	Induction Therapy: Fluconazole 1200mg PO q24h PLUS 5-flucytosine 100mg/kg/day PO q6h	Lipid formulations of amphotericin may be used in cases of severe nephrotoxicity: *The lipid formulations: Amphotericin B lipid complex: 5mg/kg IV q24h Amphotericin B liposomal: 3-4mg/kg IV q24h may be used.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			*Both lipid formulations require special medicine approval Duration of induction therapy : 4-6 weeks
	Consolidation Therapy: Fluconazole 400-800mg PO q24h		Duration of consolidation therapy : 8 weeks
	Maintenance Therapy: Fluconazole 200mg PO q24h		Duration of maintenance therapy : up to 12 months
Healthcare-associated ventriculitis and meningitis	Empirical treatment should be decided by the primary team based on local antibiogram and CSF gram stain result.		De-escalate antibiotics to targeted therapy when the culture results are available. *Vancomycin trough level should be 15- 20mcg/L
	If C&S is not available: Ceftazidime 2g IV q8h PLUS/MINUS *Vancomycin 25-30mg/kg loading dose then 15-20mg/kg IV q8-12h; not to exceed 2g per dose	Meropenem 2g IV q8h PLUS/MINUS *Vancomycin 25-30mg/kg loading dose then 15-20mg/kg IV q8-12h; not to exceed 2g per dose	
Cranial Trauma 1. Open fracture & 2. Penetrating injuries	Amoxicillin/clavulanate 1.2g IV q8h	Cefuroxime 1.5g IV q8H PLUS Metronidazole 500mg IV q8H	Duration: 5-7 days
Penetrating craniocerebral injuries	Ceftriaxone 2g IV q12h PLUS Metronidazole 400mg PO q8h		Duration: for 2 weeks initially and then review with microbiology
Neurosyphilis	Refer to section (Sexually Transmitted Infections)		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
HIV related CNS infection	Refer to section (Infections in Immunocompromised Patients - Human Immunodeficiency Virus)		

References:

1. Brouwer MC et al. Corticosteroids for acute bacterial meningitis. Cochrane Database of Systematic Reviews 2015, Issue 9. Art. No.: CD004405
2. Pasquale Pagliano et al. Listeria monocytogenes meningitis in the elderly: epidemiological, clinical and therapeutic findings. Le Infezioni in Medicina, n. 2, 105-111, 2016
3. van de Beek, D. et al. ESCMID guideline: diagnosis and treatment of acute bacterial meningitis. Clinical Microbiology and Infection, Volume 22 , S37 - S62
4. McGill, F. et al. The UK joint specialist societies guideline on the diagnosis and management of acute meningitis and meningococcal sepsis in immunocompetent adults. Journal of Infection, Volume 7, Issue 4, 405 – 438.
5. Solomon, T. et al. Management of suspected viral encephalitis in adults – Association of British Neurologist and British Infection Association National Guidelines. Journal of Infection Volume 64, Issue 4 , 347 – 373.
6. Allan R. Tunkel et al. 2017 Infectious Diseases Society of America’s Clinical Practice Guidelines for Healthcare-Associated Ventriculitis and Meningitis, Clinical Infectious Diseases Volume 64, Issue 6, 15 March 2017,
7. Peter R. Williamson et al. Cryptococcal meningitis: epidemiology, immunology, diagnosis and therapy. Nature Reviews Neurology Volume 13, pages 13–24 (2017)
8. The Sanford Guide to Antimicrobial Therapy 2018.
9. Antibiotic Expert Groups. Therapeutic guidelines: antibiotics. Version 15. Melbourne: Therapeutic Guidelines Limited; 2014.
10. R Bayston, J de Louvois, E M Brown, R A Johnston, P Lees, I K People, “Infection in Neurosurgery ” Working Party of British Society for Antimicrobial Chemotherapy. Lancet 2000; 355: 1813–17
11. Chaudhuri A. et al. EFNS guideline on the management of community-acquired bacterial meningitis: report of an EFNS Task Force on acute bacterial meningitis in older children and adults. European Journal of Neurology 2008, 15: 649–659
12. Allan R. Tunkel et al. Practice Guidelines for the Management of Bacterial Meningitis Clinical Infectious Diseases 2004; 39:1267–84
13. Clinical Practice Guidelines on Management of Tuberculosis (3rd edition) MOH/P/PAK/258.12(GU).
14. Clinical Practice Guidelines for the management of cryptococcal disease:2010 update by IDSA.
15. Guideline for Management of Brain Abscess and Subdural Empyema, Leeds Teaching Hospitals NHS Trust 2022 (website: <http://www.lhp.leedsth.nhs.uk/detail.aspx?id=1431>)- last review 3/1/22 .

GASTROINTESTINAL INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Oropharyngeal Candidiasis			
Oral Thrush	Refer to section (Infections in Immunocompromised Patients - HIV)		
Esophagitis			
Candida	Refer to section (Infections in Immunocompromised Patients - HIV)		
Herpes simplex virus	Refer to section (Sexually Transmitted Infections)		
Cytomegalovirus	Refer to section (Infections in Immunocompromised Patients - HIV)		
Helicobacter pylori Infection			
	First line treatment* Triple Therapy: Proton Pump Inhibitors PO q12h PLUS Amoxicillin 1g PO 12h PLUS Clarithromycin 500mg PO q12h OR Metronidazole 400mg PO q12h Second Line Treatment¹ <u>Fluoroquinolone triple therapy:</u> Proton Pump Inhibitors PO q12h PLUS	Antibiotic allergy: Proton Pump Inhibitors PO q12h PLUS Clarithromycin 500mg PO q12h PLUS Metronidazole 400mg PO q12h	Dosages of Proton Pump Inhibitors: -Pantoprazole 40mg PO q12h -Esomeprazole 40mg PO q12h *First line treatment recommended in areas with < 15-20% of Clarithromycin resistance Duration of therapy = 14 days -Meta analysis of RCTs found 14 days duration of therapy showed greater eradication rate

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	Levofloxacin 500mg PO q24h PLUS Amoxicillin 1g PO q12h OR Metronidazole 400mg PO q12h		
References : 1. Malfertheiner P et al. Management of Helicobacter pylori infection--the Maastricht IV/ Florence Consensus Report .Gut 2012;61: 646 – 64. 2. V. Mahachaiet al., “H. Pylori management in ASEAN: the Bangkok consensus report,” Journal of Gastroenterology and Hepatology, vol. 33, no. 1, pp. 37– 56, 2017 3. Chey WD et al. ACG clinical guideline: Treatment of Helicobacter pylori infection. Am J Gastroenterol 2017 Jan10.			
Infectious Diarrhoea Most community-acquired diarrhoea is viral in origin (norovirus, rotavirus, and adenovirus), antibiotic therapy does not shorten the duration of symptoms, therefore should be discouraged. When to consider empirical treatment for acute diarrhoea? • Immunocompetent host with high grade or persistent fever, or with dysentery or in sepsis • Immunocompromised host • Suspected enteric fever			
Empirical treatment	Ciprofloxacin 500mg PO q12h OR Azithromycin 500mg PO q24h		Duration: 3 days
Non-Shiga toxin producing <i>E.coli</i> (STEC), <i>Aeromonas/ Plesiomonas</i> <i>Yersinia species</i>	Trimethoprim/sulfamethoxazole 160/800mg PO q12h	Ciprofloxacin 500mg PO q12h	Duration: 3 days
<i>Campylobacter jejuni</i>	Azithromycin 500mg PO q24h		Duration: 3 days

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			In immunocompromised: consider longer duration of therapy
<i>Salmonella</i> , non-typhi	Trimethoprim/sulfamethoxazole 160/800mg PO q12h	Ciprofloxacin 500mg PO q12h OR Azithromycin 500mg PO q24h OR Ceftriaxone 2gm IV q24h	Duration : Immunocompetent: 5-7 days Immunocompromised: 14 days Antibiotic is usually not indicated, except in patient • < 6 months old or >50 years old • with severe illness or in septic shock • with prostheses, valvular heart disease or severe atherosclerosis • with malignancy • who is immunocompromised
<i>Salmonella</i> , non-typhi - in HIV patients	Refer to section (Infections in Immunocompromised Patients - HIV)		
<i>Salmonella enterica</i> serovar Typhi and Paratyphi (Typhoid Fever)	Refer to section (Tropical Infections)		
<i>Vibrio cholera</i>	Primary therapy is rehydration Azithromycin 1g PO single dose OR Doxycycline 300mg PO single dose	Erythromycin Ethylsuccinate 800mg PO q12h for 3 days	Antibiotic is to reduce the shedding time

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
<i>Shigella</i> sp. (Fever and bloody stool)	Ciprofloxacin 750mg PO q12h for 3 days	Azithromycin 500mg PO q24h for 3 days OR Trimethoprim/sulfamethoxazole 160/800mg PO q12h for 3 days For severe disease: Ceftriaxone 2gm IV q24h for 2-5 days	In immunocompromised patients, duration of antibiotic is 7-10 days.
<i>Giardiasis</i>	Metronidazole 400mg PO q8h		Duration : 7-10 days
<i>Entamoeba histolytica</i>	Metronidazole 800mg PO q8h for 5–10 days PLUS *Paromomycin 500mg PO q8h for 7 days		*Requires special medicine approval
<i>Clostridium difficile</i> Infection (CDI) - Discontinue therapy with inciting antibiotic agent as soon as possible as may influence risk of CDI recurrence			
Initial (Non severe)	Vancomycin 125mg PO q6h	Metronidazole 400mg PO q8h	Duration : 10 days
Initial (Severe)	Vancomycin 125mg PO q6h	Metronidazole 500mg IV q8h PLUS Vancomycin 125mg PO q6h	Duration : 10 days Symptoms to indicate severe colitis: -WCC >15 x 10 ⁹ -Creatinine 50% increase from baseline - Temperature > 38.5°C - Evidence of severe colitis (abdominal signs; radiography)

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Initial (Fulminant/ Severe with Complications)	*Vancomycin 500mg PO q6h PLUS Metronidazole 500mg IV q8h		Duration: 10 days *If ileus, consider rectal instillation of Vancomycin (enema): Vancomycin 500mg (in 100 mL normal saline) q6h via enemas or by nasogastric tube
Recurrent	If vancomycin was used for the first episode: <u>Vancomycin pulsed-tapered regimen</u> 125mg PO q6h for 10-14 days then 125mg PO q12h for 7 days then 125mg PO q24 for 7 days then 125mg PO q48-72h for 2-8 weeks	If metronidazole is used for the first episode: Vancomycin 125mg PO q6h for 10- 14 days	Duration: 10 - 14 days
References: 1. Riddle et al. ACG clinical guideline: diagnosis, treatment, and prevention of acute diarrheal infections in adults. Am J Gastroenterol. 2016;111:602–622 2. Shane et al. 2017 IDSA CPG for the Diagnosis and Management of Infectious Diarrhea.CID 2017:65. 3. McDonald et al. Clinical Practice Guidelines for Clostridium difficile Infection in Adults and Children: 2017 Update by IDSA and Society for Healthcare Epidemiology of America (SHEA).Clin Infect Dis.Feb 15,2018. doi:10.1093/cid/cix1085 4. UK: https://www.guidelines.co.uk/infection/phe-infectious-diarrhoea-guideline/252651.article 5. AUS: https://www.ncbi.nlm.nih.gov/pubmed/27062204			
Traveller's Diarrhea			
Mild Diarrhoea -Diarrhoea that is tolerable, not distressing	Antibiotic is not indicated		
Moderate Diarrhoea	Azithromycin 1g PO single dose or 500mg PO q24h	Ciprofloxacin 750mg PO stat or 500mg PO q12h	Antibiotic therapy may be considered Duration : 3 days

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
-Diarrhoea that is distressing or interferes with planned activities Severe Diarrhoea -Diarrhoea that incapacitating or completely prevents planned activities; including all dysentery	Azithromycin 1g PO single dose or 500mg PO q24h	Ciprofloxacin 750mg PO stat or 500mg PO q12h	Antibiotic therapy may be considered Duration : 3 days
References : 1. Riddle et al. Guidelines for the prevention and treatment of travellers' diarrhoea: a graded expert panel report. J Travel Med 2017; 24:S7 2. https://www.racgp.org.au/afp/2015/januaryfebruary/advising-travellers-about-management-of-travellers%E2%80%99-diarrhoea/ (aus)			
Liver Abscess			
Pyogenic liver abscess <i>Klebsiella spp.</i> , <i>Escherichia coli</i>	Empirical therapy Ceftriaxone 2g IV q24h OR Cefotaxime 2g IV q8h PLUS/MINUS *Metronidazole 500mg IV q8h	Amoxicillin/clavulanate 1.2g IV q6-8h	Duration : 4-6 weeks To consider drainage if abscess size is ≥ 5 cm or impending rupture. *Metronidazole: Risk factors of anaerobic liver abscess: • acute and chronic inflammatory bowel disease with or without perforation • malignancy of gastrointestinal tract • surgery of the gastrointestinal tract or pelvic organs Metronidazole has excellent

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			bioavailability: consider IV to PO switch
Amoebic liver abscess <i>Entamoeba histolytica</i>	Amoebicidal agent: *Metronidazole 750mg IV q8h for 10 days followed by Luminal agent: -To eradicate intestinal colonisation after Amoebicidal treatment **Paromomycin 25-35mg/kg/day PO q8h for 7 days		*May switch to PO Metronidazole 800mg q8h when there is satisfactory clinical improvement. Drainage of amoebic liver abscess is not usually required but is necessary if : 1. The patient does not respond to antibiotic therapy 2. The abscess is >5 cm in diameter 3. The abscess is in the left lobe of the liver 4. The diagnosis remains in doubt **Requires special medicine approval.
References : 1. Anesi,J and Gluckman S.Amebic Liver Abscess Clinical Liver Disease, Vol 6, No 2, August 2015 2. Stanley et al. Amoebiasis, Lancet 2003; 361: 1025–34 3. Hope et. Optimal treatment of hepatic abscess. Am Surg. 2008;74:178-182			
Cholecystitis and Cholangitis			
Community-acquired: - <i>Enterobacteriales</i> (>50%) [†] -Bacteroides only comprise 4-20% of biliary infection.	Amoxicillin/clavulanate 1.2g IV q8h	3 rd gen. cephalosporins: Cefoperazone 1-2g IV q12h OR Cefotaxime 2g IV q8h	Duration: 4-7 days Appropriate source control to drain infected foci and restoration of

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
		PLUS/MINUS Metronidazole 500mg IV q8h (if biliary enteric anastomosis is present)	anatomic and physiological function is recommended for all patients, as antibiotics will not enter bile in the presence of obstruction.
Hospital-acquired	*Piperacillin/tazobactam 4.5g IV q6-8h		* Piperacillin/tazobactam: If given as q8h, to be given as extended infusion (over 3-4 hours).
Reference s: 1. Gomi,H. et al. Tokyo Guidelines 2018: antimicrobial therapy for acute cholangitis and cholecystitis. J Hepatobiliary Pancreat Sci, 25: 3–16. doi:10.1002/jhbp.518 2. Joseph et al, Diagnosis and Management of Complicated Intra-abdominal Infection in Adults and Children: Guidelines by the Surgical Infection Society and the Infectious Diseases Society of America , CID 2010; 50:133–64			
Spontaneous bacterial peritonitis (SBP)			
Primary SBP Enterobacterales (<i>eg. E. coli</i> <i>K. pneumoniae</i> , <i>Streptococcus sp.</i>)	Cefotaxime 2g IV q8h OR Ceftriaxone 1g IV q12h	Amoxicillin/clavulanate 1.2g IV q8h	Duration : 5 days
In cirrhotic with upper gastrointestinal haemorrhage	Ceftriaxone 1g IV q12h OR Cefotaxime 2g IV q8h	*IV Ciprofloxacin 400mg IV q12h	Duration : 7 days *Ciprofloxacin does not penetrate into ascitic fluid to the same extent as cefotaxime
References : 1. Bhuvu et al. Spontaneous bacterial peritonitis: An update on evaluation, management, and prevention Am J Med 1994;97:169–175. 2. Rimola et al. Diagnosis, treatment and prophylaxis of spontaneous bacterial peritonitis: a consensus document. Journal of Hepatology 2000; 32: 142-153 3. Rimola A, Salmerón JM, Clemente G, Rodrigo L, Obrador A, Miranda ML,et al. Two different dosages of cefotaxime in the treatment of spontaneous bacterial peritonitis in cirrhosis: results of a prospective, randomised,multicenter study. Hepatology 1995;21:674–679.			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
4. Navasa M, Follo A, Llovet JM, Clemente G, Vargas V, Rimola A, et al. Randomised, comparative study of oral ofloxacin versus intravenous cefotaxime in spontaneous bacterial peritonitis. <i>Gastroenterology</i> 1996;111:1011–1017. 5. Sort P, Navasa M, Arroyo V, Aldeguer X, Planas R, Ruiz delArbol L, et al. Effect of intravenous albumin on renal impairment and mortality in patients with cirrhosis and spontaneous bacterial peritonitis. <i>N Engl J Med</i> 1999;341:403–409. 6. Ricart E, Soriano G, Novella MT, Ortiz J, Sàbat M, Kolle L, et al. Amoxicillin– clavulanic acid versus cefotaxime in the therapy of bacterial infections in cirrhotic patients. <i>J Hepatol</i> 2000;32:596–602. 7. UK: https://gut.bmj.com/content/55/suppl_6/vi1 8. Terg R, et al. Oral ciprofloxacin after a short course of intravenous ciprofloxacin in the treatment of spontaneous bacterial peritonitis: results of a multicenter, randomised study. <i>J Hepatol</i>. 2000;33(4):564-569			

INFECTIONS IN IMMUNOCOMPROMISED PATIENTS

A. HAEMATOLOGY/ONCOLOGY

1. Any infection in the immune-compromised host is life-threatening and needs immediate attention.
2. Febrile neutropenia is defined as a temperature $>38.3^{\circ}\text{C}$ on a single occasion or $>38^{\circ}\text{C}$ over one hour and ANC (Absolute Neutrophil Count) <500 cells/uL or <1000 cells/uL in those with anticipated declining counts.
3. Cultures may be positive in $< 40\%$ of cases. Patients have impaired inflammatory responses and hence may have no localising signs. The usual sign is fever $>38^{\circ}\text{C}$ or hypothermia.
4. The common portals of infection include the oral cavity, gastrointestinal tract, perianal region, lungs and IV lines. Potential pathogens are dependent on the underlying defect, e.g.

Neutropenia	Gram-positive organisms, Gram-negative organisms, Fungi
Hypogammaglobulinemia; Post splenectomy or hyposplenic patients	Encapsulated organisms
Defective cellular immunity	Pneumocystis, Toxoplasma, Fungi, Viruses, Mycobacteria

5. The choice of antibiotic is based on local organisms and sensitivity patterns, sound clinical judgement, the clinical state of the patient, prior infections with drug-resistant bacteria, recent outbreaks (e.g. MRSA or multidrug resistant *Enterobacteriales*) as well as the availability and cost of the antibiotics.
6. Surveillance for **CRE**, *Stenotrophomonas maltophilia* and multi-resistant organisms should be carried out by the infection control team. The incidence of these organisms must be borne in mind when selecting agents for use in the first line setting.
7. Validated risk assessment scoring systems included the Multinational Association for Supportive Care in Cancer (MASCC) score and the Clinical Index of Stable Febrile Neutropenia (CISNE) score should be used for the classification of low or high risk group of patients

8. LOW RISK PATIENTS

The use of **oral antibiotics** with **Amoxicillin/clavulanate 625mg q8h (Clindamycin 600mg q6h if penicillin allergic) PLUS/MINUS Ciprofloxacin 500mg q12h** (if previously colonised by *Pseudomonas aeruginosa* and patient not on prior fluoroquinolone prophylaxis) may be considered after careful assessment of risk factors and a consult from the hematologist.(i.e no evidence of dehydration or hypotension, no pneumonia/COAD). Besides, it is important that patients must be able to access prompt medical attention if their condition deteriorates.

9. HIGH RISK PATIENTS

Risk assessment for complications of severe infection should be done during triage. Patient should be admitted to hospital for **IV antibiotics** if deemed **high risk** with the presence of:

- a) prolonged (> 7 days) and profound ANC < 0.1x10⁹/L,
- b) hypotension
- c) pneumonia
- d) new onset abdominal pain or neurological signs
- e) MASCC <21 or CISNE ≥ 3 at the time of assessment

10. The administration of the first dose of empirical antipseudomonal antibiotic should be done as soon as possible following triage (within the 1st hour) after taking blood cultures. The following regimens are suggested: Empirical antibiotics should be continued until ANC > 0.5 x 10⁹/L and the patient is clinically stable and afebrile.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Febrile Neutropenia			
<i>Pseudomonas aeruginosa</i>	First line Piperacillin/tazobactam 4.5g IV q6h OR Cefepime 2g IV q8h PLUS/MINUS *Gentamicin 5mg/kg IV q24h OR Amikacin 15mg/kg q24h Second line	**Ceftazidime 2g IV q8h	Duration: until neutrophils count >500 cells/uL or longer if clinically indicated (> 1 x 10 ⁹ /L) *Monotherapy (Piperacillin/tazobactam, cefepime, imipenem or meropenem) is likely as efficacious and less toxic. Aminoglycosides may be added in patients with severe sepsis for broader Gram-negative coverage. It

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	After 3 days, no improvement or condition worsening (severe sepsis or impending shock): Meropenem 1g IV q8h (2g IV q8h if suspected central nervous system infection) OR Imipenem 500mg IV q6h (1g q8h if severe sepsis)		can be discontinued if microbiological cultures showed isolated organisms sensitive to piperacillin/tazobactam or cefepime and the patient is clinically stable. **Weak Gram-positive coverage and increased breakthrough infections limit utility Carbapenem NOT commonly recommended as 1st line unless - severe sepsis - patients with previous infections of ESBL organisms or gram-negative organisms resistant to narrow spectrum B-lactams
Anaerobes	PLUS Metronidazole 500mg IV q8h		Metronidazole may be added to cefepime in the presence of 1) severe mucositis 2) intra-abdominal infections 3) peri-anal abscesses or 4) pseudomembranous colitis Piperacillin/tazobactam and carbapenems have good anaerobic coverage and therefore do not need addition of metronidazole.
MRSA	PLUS	VRE/VISA/VRSA:	Empirical used of vancomycin if:

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	Vancomycin 15mg/kg IV q8-12h	**Linezolid 600mg PO/IV q12h	a. Known colonisers by MRSA not responding to adequate empirical antibiotics b. Suspected catheter-related infections c. Suspected/ proven skin and soft tissue gram positive infections d. Positive blood culture for gram positive bacteria/ proven methicillin-resistant gram-positive infections e. Severe mucositis f. Severe sepsis, septic shock or respiratory distress To stop after 48 hours if there is no evidence of Gram-positive infection. **To change to Linezolid if MRSA/MRSE not responding to or intolerant to Vancomycin and in those with suspected or confirmed VRE, VISA or VRSA.
11. Fungal infection Invasive fungal infection (IFI) is one of the most important infectious complications seen in patients with hematological malignancies and haematopoietic stem cell transplant (HSCT) recipients. <i>Aspergillus</i> species remain the most common pathogen followed by <i>Candida</i> species. The main risk factors for IFI are patients with underlying hematological malignancies, profound neutropenia, following a myelosuppressive chemotherapy, allogeneic HSCT and graft versus host disease. Suspect fungal infection when there is:			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
a) mucositis, thrush b) difficulty or pain when swallowing c) suspicious skin lesions or pulmonary infiltrates d) fundal exudates e) persisted fever not responding to antibiotics (4 - 7 days) f) prolonged steroid/ antibiotic usage (>2 week)			
Invasive fungal infection	Micafungin 100-150mg IV q24h OR Caspofungin 70mg IV loading, then 50 mg IV q24h OR Anidulafungin 200mg IV loading, then 100mg IV q24h	Voriconazole 6mg/kg q12h for 2 doses, then 4mg/kg q12h OR Amphotericin B deoxycholate 0.7 - 1mg/kg IV q24h	Azoles should not be used for empirical therapy in patients who have received an azole for prophylaxis.
Invasive Aspergillosis	Voriconazole 6 mg/kg q12h for 2 doses, then 4mg/kg q12h	Amphotericin B deoxycholate 0.7 -1mg/kg IV q24h OR IV Micafungin 150mg od OR Anidulafungin 200mg IV loading, then 100mg IV q24h OR Caspofungin 70mg IV loading, then 50 mg IV q24h	
Candidaemia	Micafungin 100-150mg IV q24h OR Anidulafungin 200mg IV loading, then 100mg IV q24h OR	*Fluconazole 800 mg [12 mg/kg] stat, then 400 mg [6 mg/kg] q24h) **Voriconazole: 400 mg (6 mg/kg) q12h for 2 doses, then 200 mg (3 mg/ kg) q12h	*For patients who are less critically ill and who have no recent azole exposure.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	Caspofungin 70mg IV loading, then 50 mg IV q24h		**Can be used in situations in which additional mould coverage is desired. Recommended duration of therapy for candidemia without persistent fungemia or metastatic complications is 2 weeks after documented clearance of Candida from the bloodstream and resolution of symptoms attributable to candidemia and resolution of neutropenia.
12. The use of prophylactic growth factors (e.g.G-CSF) i) Significantly reduced the relative risk for severe neutropenia, febrile neutropenia and infection and should be considered in high-risk patients with ANC < 100/UI, multiple organ dysfunction syndrome, pneumonia, invasive fungal infections, septic shock or patients with reduced bone marrow reserve due to extensive radiotherapy. ii) However, there is no evidence that either G-CSF reduced the number of patients requiring intravenous antibiotics or lowered infection related mortality. iii) The role of granulocytes remains controversial and should be discussed with haematologists. iv) Granulocyte transfusions may be used in patients with serious bacterial or fungal infections not responding to appropriate treatment and who will likely recover in the neutrophil count in the short term. The risk of disease transmission e.g. CMV must be borne in mind.			
References: 1. NCCN Clinical Practice Guidelines in Oncology. Prevention and Treatment of Cancer-Related Infections, version 2. 2017. National Cancer Comprehensive Network website. www.nccn.org/professionals/physician_gls/PDF/infections.pdf. Published February 21, 2017. Accessed September 27, 2017. 2. Hughes W T, Armstrong D, Bodey G P et al. 2002 Guidelines for the use of antimicrobial agents in neutropenic patients with cancer.Clinical Infectious disease 2002; 34:730-751 3. Herbrecht R, Denning D W, Patterson T F et al. Voriconazole versus amphotericin B for primary therapy of invasive aspergillosis. NEJM 2002 ; 347: 408-415 4. WalshTJ, Teppler H, Donowitz G R et al .Caspofungin versus liposomal amphotericin B for empirical antifungal therapy in patients with persistent fever and neutropenia. NEJM 2004;351(14):1391-1402 5. Dellinger RP, Levy MM, Carlet JM et al.Surviving sepsis campaign: international guidelines for management of severe sepsis and septic shock. Intensive Care Med 2008, 34 : 17-60			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
6.	6. Bohlius J, Herbst C, Reiser M, Schwarzer G, Engert A. Granulopoiesis-stimulating factors to prevent adverse effects in the treatment of malignant lymphoma. Cochrane Database of Systematic Reviews 2008, Issue 4.		
7.	Alison GF, Eirc JB, Kent A S et al. IDSA guideline: Clinical Practice guideline for the Use of Antimicrobial Agents in Neutropenic Patients with Cancer: 2010 update by the Infectious Diseases Society of America. CID 2011; 52: 56 -93		
8.	Mica P, Sara B, Abigail F, Liat v and Leonard L. Empirical antibiotics against Gram -positive infections for febrile neutropenia: Systemic review and meta-analysis of randomised controlled trials. J Antimicrob Chemother 2005; 55: 436-444.		
9.	Diana A, ChristinaO, Catherine C et al. European guidelines for empirical antibacterial therapy for febrile neutropenic patients in the era of growing resistance: summary of the 2011 4th European conference on the infections in Leukaemia. Haematologica 2013;98(12) :1826-1835		
10.	Paul M, Yahav D, Fraser A, et al. Empirical antibiotic monotherapy for febrile neutropenia: Systematic review and meta-analysis of randomised controlled trials. J Antimicrob Chemother 2006; 57:176-89.		
11.	Engelhard D. Akova M, Boeckh MJ et al. Bacterial infection prevention after hematopoietic cell transplantation. Bone marrow Transplant 2009; 44:467-470.		
12.	Cornely OA, Bohme A, Buchheidt D et al. Primary prophylaxis of invasive fungal infections in patients with hematologic malignancies. Recommendations of the Infectious Diseases Working Party of the German Society for Haematology and oncology. Haematologica 2009; 94: 113-22.		
13.	Zaia J, Baden L, Boeckh MJ et al. viral disease protection after hematopoietic cell transplantation. Bone marrow Transplant 2009; 44:471-4		
14.	NCCN Clinical Practice Guidelines in Oncology. Myeloid Growth Factors, version 1.2022. National Cancer Comprehensive Network website. www.nccn.org/professionals/physician_gls/PDF/myeloid_growth.pdf. Published April 28, 2017. Accessed May 22, 2022.		
15.	Taplitz RA, Kennedy EB, Bow EJ, Crews J, Gleason C, Hawley DK, Langston AA, Nastoupil LJ, Rajotte M, Rolston K, Strasfeld L. Outpatient management of fever and neutropenia in adults treated for malignancy: American Society of Clinical Oncology and Infectious Diseases Society of America Clinical Practice Guideline Update. J Clin Oncol. 2018 May 10;36(14):1443-53.		
16.	Zimmer AJ, Freifeld AG. Optimal management of neutropenic fever in patients with cancer. Journal of oncology practice. 2019 Jan ;15(1):19-24.		
17.	Klastersky J, De Naurois J, Rolston K, Rapoport B, Maschmeyer G, Aapro M, Herrstedt J. Management of febrile neutropenia: ESMO clinical practice guidelines. Annals of Oncology. 2016 Sep 1;27:v111-8.		
18.	Claeys KC, Zasowski EJ et al. Improves Outcomes Regardless of Vancomycin MIC in a Propensity-Matched Analysis of Methicillin-Resistant staphylococcus aureus Bloodstream Infections. Antimicrob Agents Chemother. 2016;60(10):5841.		

B. HUMAN IMMUNODEFICIENCY VIRUS (HIV)

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
<i>Pneumocystis jiroveci (carinii)</i> interstitial pneumonia (PJP/PCP)			
Treatment	Trimethoprim/sulfamethoxazole 15-20mg/kg/day [TMP component]/ IV/PO (excellent bioavailability) in 3-4 divided doses	Primaquine 30mg (base) PO q24h PLUS Clindamycin 600mg IV q6h or 900mg q8h OR *Pentamidine 4mg/kg/day IV (in 1 pint D5% or NS run over 1-2 hours)	Duration: 21 days Patients with severe disease (PO2 < 70mmHg at room air) should receive steroids as soon as possible (Within 72 hours of starting PCP treatment): Prednisolone* dose: 40mg PO q12h for 5 days then 40mg PO q24h for 5 days then 20mg PO q24h for 11 days IV hydrocortisone or methylprednisolone at equivalent dose to prednisolone can be substituted for oral prednisolone. Patients should be screened for G6PD deficiency before dapsone or primaquine is administered. TMP/SMX PO has excellent bioavailability and should be preferred for patients who can tolerate orally without concerns over drug absorption.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			Rates of adverse reaction to TMP-SMX are high. However, supportive care for non-life-threatening adverse effects should be attempted before the drug is discontinued. Lower doses of TMP-SMX may also be effective, potentially with lower toxicity. *Requires special medicine approval
Prophylaxis (primary and secondary) <i>Indications:</i> • CD4 count <200 cells/mm³ or percentage less than 14%. • Evidence of AIDS-defining illness, eg. oropharyngeal candidiasis, pulmonary TB etc. • CD4 count 200-250 and if CD4 monitoring (e.g., every 3 months) is not possible or ART cannot be initiated	Trimethoprim/sulfamethoxazole 80/400mg 1 - 2 tablets PO q24h	Dapsone 100mg PO q24h OR *Aerosolized Pentamidine 300mg monthly (recommended to be delivered via Respigard II nebulizer)	Patients with CD4<100 cells/μL should receive at least 2 tablets of TMP/SMX daily to extend prophylaxis against toxoplasmosis. Prophylactic regimens other than TMP/SMX should only be considered if TMP/SMX is absolutely contraindicated. Aerosolized pentamidine or dapsone alone (without pyrimethamine) cannot prevent toxoplasmosis, hence they may not be suitable for patients who are at high risk of toxoplasmosis infection (seropositive toxoplasma IgG +ve and CD4<100 cells/μL) *Requires special medicine approval

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			Discontinuation of prophylaxis: Consider in patients with CD4>200 cells/μL and undetectable HIV viral load in response to successful ART. Restarting prophylaxis: CD4 count falls to <200 cells/μL OR PCP recurred at a CD4 count >200 cells/μL (lifelong prophylaxis should be considered)
Mucocutaneous Candidiasis			
Oropharyngeal (oral thrush)	Fluconazole 100mg PO q24h	Nystatin suspension 500,000 units PO q6h OR Itraconazole oral solution 200mg PO q24h	Duration: 7-14 days Chronic suppressive therapy is usually not recommended unless patients have frequent or severe recurrences. If used, it is reasonable to discontinue therapy if CD4 >200 cells/ μ L.
Oesophageal	Fluconazole 200mg PO/IV q24h for 14 days	Itraconazole oral solution 200mg PO q24h OR Voriconazole 200mg PO/IV q12h OR Echinocandins (anidulafungin, caspofungin or micafungin) OR	Candidiasis is the most common cause of esophagitis with HIV infection, but CMV, HSV and aphthous ulcerations can present with similar complaints.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
		Amphotericin B deoxycholate 0.6mg/kg IV q24h	Endoscopy required with unusual presentations or lack of response to azole within several days.
Vulvovaginal	Refer to section (Obstetrics & Gynaecology Infections)		
Cryptococcal meningitis or meningoencephalitis (<i>Cryptococcus neoformans</i>)			
Induction Therapy (for at least 2 weeks)	<i>If liposomal amphotericin B is not available:</i> Amphotericin B deoxycholate 0.7-1mg/kg IV q24h PLUS Flucytosine 25mg/kg PO q6h for 7 days, followed by Fluconazole 1200mg PO q24h for 7 days OR <i>If liposomal amphotericin B is available:</i> Liposomal Amphotericin B 10mg/kg IV (single dose) PLUS 14 days of: Flucytosine 25mg/kg PO q6h PLUS Fluconazole 1200mg PO q24h	<i>If flucytosine is not available:</i> Amphotericin B deoxycholate 0.7-1mg/kg IV q24h PLUS Fluconazole 1200mg PO q24h OR Liposomal Amphotericin B 3-4mg/kg IV q24h PLUS Fluconazole 1200mg PO q24h <i>If no amphotericin B formulations are available:</i> Fluconazole 1200mg PO q24h PLUS Flucytosine 25mg/kg POq6h	Test dose of Amphotericin B is not required. Ensure adequate hydration (e.g. 2 pints of IVD NS before and after administration of Amphotericin B) to prevent nephrotoxicity. Pre-medication may be given before the administration of Amphotericin B (Promethazine 25mg IV and Paracetamol 1g PO/ Hydrocortisone 50mg IV). If clinical signs of cerebral oedema do not improve after about 2 weeks of daily LPs, consider placement of a lumbar drain or VP shunt.
Consolidation therapy	Fluconazole 800mg PO q24h	Itraconazole 200mg PO q12h	Duration: For 8 weeks

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Maintenance Therapy (Secondary prophylaxis)	Fluconazole 200mg PO q24h	Itraconazole 200mg PO q24h	Discontinuation: Consider in patients with CD4>200 cells/ μ L and undetectable HIV viral load in response to successful ART.
<i>Toxoplasma gondii</i> Encephalitis			
Acute Infection (up to 97% patients are Toxo IgG +ve) Induction therapy (for at least 6 weeks)	Trimethoprim/sulfamethoxazole 10mg/kg/day (TMP component) PO/IV in 2 divided doses	*Pyrimethamine 200mg PO loading dose followed by - 50mg q24h (BW $\leq$ 60kg) - 75mgq24h (if BW>60kg) PLUS Folinic acid 10-25mg PO q24h PLUS Clindamycin 600mg IV/PO q6h	Adjunctive corticosteroids (e.g., dexamethasone) should be administered when clinically indicated to treat mass effect associated with focal lesions and perilesional edema. Because of the potential immunosuppressive effects, corticosteroids should be tapered down and discontinued as soon as clinically feasible. Sulfadoxine/pyrimethamine 500mg/25mg (Fansidar [®]) can be used if pyrimethamine is not available. However, sulfadoxine/pyrimethamine is associated with higher incidence of adverse drug reactions such as sulfa allergy. *Requires special medicine approval
Suppressive/Maintenance Therapy	Trimethoprim/sulfamethoxazole (80/400mg) 2 tablets PO q12h	*Pyrimethamine 25-50mg PO q24h	Discontinuation of secondary prophylaxis:

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
		PLUS Folinic acid 10-25mg PO q24h PLUS Clindamycin 600mg PO q8h	Consider in patients with CD4>200 cells/ μ L and undetectable HIV viral load in response to successful ART. *Requires special medicine approval
Primary Prophylaxis Indications: ToxolG +ve with CD4<100	Trimethoprim/sulfamethoxazole (80/400mg) 2 tabs PO q24h	Dapsone 50mg PO q24h PLUS *Pyrimethamine 50mg PO once weekly PLUS Folinic acid 10-25mg PO once weekly OR Dapsone 200mg PO weekly PLUS *Pyrimethamine 75mg PO once weekly PLUS Folinic Acid 25mg PO once weekly	The recommended regimens for primary prophylaxis of toxoplasmosis are also effective in preventing PCP. Discontinuation of primary prophylaxis: Consider in patients with CD4>100 cells/ μ L and undetectable HIV viral load in response to successful ART. If CD4>100 cells/ μ L but <200 cells/ μ L, patients need to be maintained on PCP prophylaxis with TMP-SMX 80/400mg 1 tablet PO q24hr or dapsone 100mg PO q24hr. *Requires special medicine approval
<i>Mycobacterium avium</i> Complex (MAC) Disease			
Treatment	(At Least 2 Drugs as Initial Therapy) Clarithromycin 500mg PO q12h PLUS	Azithromycin 500mg PO q24h PLUS Ethambutol 15mg/kg PO q24h *PLUS	*Addition of 3rd/4th drug should be considered for patients with CD4 count <50, high mycobacterial loads (>2 log CFU/mL of blood), or in the absence of effective ART.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	Ethambutol 15mg/kg PO q24h *PLUS 3rd or 4th drug: Amikacin 10-15mg/kg IV q24h OR Streptomycin 1g IM q24h OR Levofloxacin 500mg PO q24h	3rd or 4th drug: Amikacin 10-15mg/kg IV q24h OR Streptomycin 1g IM q24h OR Levofloxacin 500mg PO q24h	Discontinuation: Consider if patient is on ART and viral load well suppressed, CD4 > 100 cells/ μ L for $\geq$ 6 months, asymptomatic of MAC, and has completed > 12 months of therapy.
Maintenance/ Secondary Prophylaxis	Same as the treatment regimen		Secondary prophylaxis restarted when CD4<100 cells/ μ L ³
Primary Prophylaxis Indications: CD4 < 50 cells/mm ³ Ruled out active MAC and TB	Azithromycin 1250mg PO once weekly	Clarithromycin 500mg PO q12h	Discontinuation: Consider if patient is on ART and viral load well suppressed, CD4 > 100 cells/ μ L for $\geq$ 3 months.
Cytomegalovirus (CMV) Disease			
CMV Retinitis Immediate Sight-Threatening Lesions (Adjacent to the Optic Nerve or Fovea)	Intravitreal injections of Ganciclovir (2mg/injection) biweekly until scarring PLUS Ganciclovir 5mg/kg IV q12h followed by maintenance therapy*	Intravitreal Foscarnet (2.4mg/ injection) biweekly until scarring PLUS Valganciclovir 900mg PO q12h followed by maintenance therapy*	Duration of initial therapy: 14-21 days *Maintenance therapy involving long duration IV Ganciclovir or high-cost Valganciclovir may not be feasible in resource-limited settings.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
CMV Retinitis For Small Peripheral Lesions	Ganciclovir 5mg/kg IV q12h for 14 days followed by maintenance therapy*	Valganciclovir 900mg PO q12h followed by maintenance therapy*	Given that early immune recovery is essential in preventing CMV relapses, ART should be started within 2 weeks of CMV treatment and clinical improvement.
Extraocular CMV diseases (Oesophagitis, colitis, interstitial pneumonitis, neurological)	Ganciclovir 5mg/kg IV q12h followed by maintenance therapy*	May consider switch to Valganciclovir 900mg PO q12h once patient can absorb and tolerate orally (in CMV oesophagitis and colitis only) followed by maintenance therapy*	Duration of initial therapy: 21-42 days or until resolution of signs and symptoms. *Maintenance therapy involving long duration IV Ganciclovir or high-cost Valganciclovir may not be feasible in resource-limited settings. Early immune recovery in response to successful ART is essential in preventing CMV relapses.
Maintenance therapy/ Secondary prophylaxis (CD4 count <100 cells/mm ³)	Ganciclovir 5mg/kg IV q24h 5–7 times weekly	Valganciclovir 900mg PO q24h	Discontinuation: Consider in patients with CD4>100 cells/μL and undetectable HIV viral load in response to successful ART. Maintenance therapy involving long duration IV Ganciclovir or high-cost Valganciclovir may not be feasible in resource-limited settings.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			Early immune recovery in response to successful ART is essential in preventing CMV relapses.
Bacterial Enteric Infections			
Salmonellosis <i>Salmonella</i> non-typhi	Ciprofloxacin 500-750mg PO q12h OR 400mg IV q12h	Trimethoprim/sulfamethoxazole 160/800mg IV/PO q12h OR Ceftriaxone 2g IV q24h OR Ampicillin 2g IV q4-6h	Duration: <i>Without bacteremia:</i> CD4≥200 cells/μL: 7-14 days. CD4 <200 cells/μL: 2-6 weeks <i>With bacteremia:</i> CD4≥200 cells/μL: At least 14 days or longer duration if persistent bacteremia or complicated with metastatic foci of infections. CD4<200 cells/μL: 2-6 weeks Secondary prophylaxis should be considered in patients with recurrent salmonella bacteremia. Effective ART may decrease the risk of recurrence of salmonella infections
Herpes Simplex Virus (HSV) Infections			
Refer to section (Sexually Transmitted Infection)			
Varicella-Zoster Virus Diseases			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Refer to section (Skin and Soft Tissue Infection (SSTI))			
Histoplasmosis (<i>Histoplasma capsulatum</i>)			
Moderate-to-severe disseminated disease	Induction therapy Amphotericin *0.7-1 mg/kg IV q24h for at least 2 weeks or clinical improvement, followed by Maintenance therapy Itraconazole 200mg PO q8h for 3 days, then q12h for at least 12 months		All the triazole antifungals have the potential to interact with certain ARV agents and other anti-infective agents. *The lipid formulations of amphotericin B may be used instead if available
Less severe disseminated disease	Itraconazole 200mg PO q8h for 3 days, then 200mg PO q12h for at least 12 months.	Fluconazole 800 mg PO q24h OR Voriconazole 400mg PO q12h on day 1, then 200mg PO q12h	Itraconazole oral solution is preferred over capsule because of improved absorption, but is less well tolerated. However, this formulation may not be necessary if itraconazole concentration is increased by concomitant use of a CYP3A4 inhibitor such as ritonavir-boosted PIs.
Chronic Suppressive therapy (Secondary prophylaxis) Indication: - severe disseminated or CNS infection after completion of at least 12 months of treatment	Itraconazole 200mg PO q24h	Fluconazole 400mg PO q24h	Discontinuation: • Received azole for >1 year, and • Negative fungal blood cultures, and • CD4 count >150 cells/μL and viral load suppressed in response to ART.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
- relapsed despite appropriate initial therapy CD4<150			
<i>Cystoisospora belli</i> infection			
Initial Therapy	Trimethoprim/sulfamethoxazole 160/800mg PO/IV q6h	*Pyrimethamine 50-75mg PO q24h PLUS Folinic acid 5-10mg PO q24h OR Ciprofloxacin 500mg PO q12h	Duration: 10 days *Requires special medicine approval
Penicilliosis (<i>Talaromyces marneffe</i> infection)			
Acute infection	*Amphotericin B 0.6-0.7mg/kg IV for 2 weeks (induction therapy), followed by consolidation therapy Itraconazole 200mg PO q12h for 10 weeks	<i>If Amphotericin B is not available or not tolerated by patients:</i> Induction therapy: Voriconazole 6 mg/kg IV q12h for 1 day, then voriconazole 4 mg/kg IV q12h for 2 weeks OR Voriconazole 600 mg PO q12h on day 1, then Voriconazole 400 mg PO q12h for 2 weeks, followed by 10 weeks of consolidation therapy with: Voriconazole 200 mg PO q12h OR	*The lipid formulations of amphotericin B may be used instead if available. Patients must be on maintenance therapy after treatment of acute infections.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
		Itraconazole 200 mg PO q12h	
Maintenance therapy/ Secondary prophylaxis	Itraconazole 200mg PO q24h	Fluconazole 400mg PO q24h	Discontinuation: CD4 count >100 cells/mm ³ and HIV viral load suppressed in response to ART.
Progressive Multifocal Leukoencephalopathy (PML)			
<i>Polyoma virus JC virus (JCV)</i>	No effective therapy exists		With ART, some patients improve and others stabilise. Few may deteriorate due to immune reconstitution.
Cryptosporidiosis			
<i>Cryptosporidium</i> sp.	Symptomatic treatment of diarrhoea		Effective ART (to increase CD4 >100 cells/ μ L) can result in complete, sustained clinical, microbiological and histologic resolution.
Rhodococcus infection			
Induction Treatment	*Vancomycin 15-20mg/kg (actual body weight) IV q8-12h PLUS (antibiotics with intracellular activity) Azithromycin 500mg stat and then 250mg IV/PO q24h	Addition of 4th drug should be considered in patients who are not responding clinically or with CNS involvement. 4th drug: Imipenem/cilastatin 500mg IV q6h	*Consider Vancomycin loading dose 25-30mg/kg for critically ill or septic patients. Aim trough level 15-20mcg/mL Once susceptibility data are available, two active agents should be continued (at least one of these should be macrolide or quinolone).

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	PLUS Ciprofloxacin 500-750mg PO q12h or 400 mg IV q8-12h		For patients with CNS involvement, the second agent has to be with good CNS penetration (eg. Imipenem/cilastatin, vancomycin or rifampicin) since macrolide and quinolone have poor penetration into CSF. Duration of treatment: 4-8 weeks (Longer courses should be administered to patients who have persistent clinical or radiographic evidence of infection).
Maintenance/ Secondary prophylaxis	Azithromycin 250mg PO q24h PLUS Ciprofloxacin 500-750mg PO 12h	*Rifampicin 600mg PO 24h PLUS Azithromycin 250mg PO 24h OR Ciprofloxacin 500-750mg PO 12h	*Alternative regime can be used if concomitant tuberculosis can be ruled out Discontinuation: Consider in patients with CD4>200 cells/μL and undetectable HIV viral load in response to successful ART.
Reference: 1) Guidelines for the Prevention and Treatment of Opportunistic Infections in Adults and Adolescents with HIV. National Institutes of Health, Centers for Disease Control and Prevention, and the HIV Medicine Association of the Infectious Disease Society of America NAG (updated June 2022)			

OBSTETRICS & GYNAECOLOGICAL INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Septic Abortion Common organisms: ● <i>Bacteroides</i> sp. ● <i>Streptococcus</i> sp. ● <i>Enterobacterales</i> ● <i>Chlamydia trachomatis</i> ● <i>Ureaplasma urealyticum</i>	Ampicillin 2g IV q4- 6h PLUS Gentamicin 5mg/kg IV q24h PLUS Metronidazole 500mg IV q8h	Ampicillin/sulbactam 3g IV q6h PLUS Doxycycline 100mg PO q12h OR Clindamycin 900mg IV q8h PLUS Gentamicin 5mg/kg IV q24h	Intravenous antibiotics are administered until the patient has improved and been afebrile for 48 hours, then are typically followed by oral antibiotics to complete a 10- to 14-day course.
Intrapartum antibiotic prophylaxis (IAP) for Group B <i>Streptococcus</i> (GBS) positive mothers Indications of IAP: ● Previous infant with invasive GBS disease ● Preterm labour ● GBS carriage in previous pregnancy ● PPROM with known GBS carrier ● GBS carriage in current pregnancy	Benzylpenicillin 5MU IV initial dose, then 2.5-3MU IV q4h until delivery	Ampicillin 2g IV initial dose, then 1g IV q4h until delivery. Mild penicillin allergy: Cefazolin 2g IV initial dose, then 1g q8h until delivery OR Cefuroxime 1.5g IV stat and 750mg IV q8h until delivery Severe penicillin allergy: Vancomycin 15-20mg/kg IV q8-12h until delivery OR Clindamycin 900mg IV q8h until delivery	Prophylaxis begins at hospital admission for labour or rupture of membrane and continues every four hours until the infant is delivered. Treatment is NOT INDICATED if C-section is performed before onset of labour with intact membrane (please use standard surgical prophylaxis). Antenatal treatment is NOT RECOMMENDED for GBS cultured from a vaginal or rectal swab.
Preterm Premature Rupture of Membranes (PPROM)	If non-GBS carrier:		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	Erythromycin ethylsuccinate 400mg PO q6h for 7-10 days If GBS carrier: Ampicillin 2g IV q6h for 48 hours, followed by Amoxicillin 500mg PO q8h for an additional 5-7 days or until delivery, whichever comes first PLUS One dose of Azithromycin 1g PO upon admission (to cover for <i>Ureaplasma</i> – important cause of chorioamnionitis & chlamydia)		
Chorioamnionitis	Ampicillin 2g IV q6h PLUS Gentamicin 5mg/kg IV q24h If the patient is undergoing a caesarean delivery: PLUS Metronidazole 500mg IV q8h	Ampicillin/sulbactam 3g IV q6h Mild penicillin allergy: Cefazolin 2g IV q8h PLUS Gentamicin 5mg/kg IV q24h Severe penicillin allergy: Clindamycin 900mg IV q8h	Antibiotic regimen is continued postpartum until the patient is afebrile and asymptomatic for AT LEAST 48 HOURS. There is NO evidence that continuation with oral antibiotics is beneficial after discontinuation of parenteral therapy.
Pelvic Inflammatory Disease Common organisms: • <i>Neisseria gonorrhoeae</i> • <i>Chlamydia trachomatis</i> • <i>Bacteroides</i> sp.	Outpatient regimen (Mild-moderate): Ceftriaxone 500mg IM in a single dose OR Cefotaxime 1g IM in a single dose PLUS		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
• <i>Enterobacterales</i> • <i>Haemophilus influenzae</i> • <i>Streptococcus</i> sp. especially <i>Streptococcus agalactiae</i> (GBS) • <i>Gardnerella vaginalis</i> • <i>Ureaplasma urealyticum</i> • <i>Mycoplasma hominis</i>	Metronidazole 400mg PO q12h for 14 days PLUS Doxycycline 100mg PO q12h for 14 days OR Azithromycin 1g PO once a week for 2 weeks		
	Inpatient regimen (Moderate-severe): Cefuroxime 1.5g IV q8h OR Ceftriaxone 2g IV q24h PLUS Doxycycline 100mg PO q12h PLUS Metronidazole 500mg IV/PO q8h Duration of treatment: 14 days	Ampicillin/sulbactam 3g IV q6h PLUS Doxycycline 100mg PO q12h	Tubo-ovarian abscess: Surgical intervention for source control may be required. May need to consider tuberculosis if not responding to standard treatment.
Endometritis	Non-pregnancy: Follow antibiotic guide for severe PID		
	Postpartum endometritis: Clindamycin 900mg IV q8h PLUS *Gentamicin 5mg/kg q24h OR Cefotaxime 1g IV q8h	Amoxicillin/clavulanate 1.2g IV q8h OR Ampicillin/sulbactam 3g IV q6h	Duration of treatment: 10-14 days *TDM for gentamicin is required

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	PLUS Metronidazole 500mg IV q8h PLUS *Gentamicin 5mg/kg IV x 1 dose		
Vaginitis Bacterial vaginosis	Metronidazole 400mg PO q8h for 7 days	Clindamycin 300mg PO q12h for 7 days	Metronidazole can be used in any stage of pregnancy
Vaginal Candidiasis <i>Candida albicans</i>	Uncomplicated infection: Clotrimazole 500mg as a single vaginal pessary (stat dose) OR Clotrimazole 200mg as vaginal pessary for 3 nights	Fluconazole 200mg PO for one dose	Pregnancy: If indicated, treatment with topical therapy as oral therapy is CONTRAINDICATED.
	Complicated infections: <u>Severe vaginitis symptoms</u> Fluconazole 150-200mg PO q72h for 2 or 3 doses <u>Recurrent vulvovaginal candidiasis</u> Fluconazole 200mg PO q72h for 3 doses then weekly for 6 months	Clotrimazole 500mg vaginal suppository once weekly for 6 months	
Trichomoniasis <i>Trichomonas vaginalis</i>	Metronidazole 400mg PO q8h for 7 days OR Metronidazole 2g PO as single dose		Metronidazole can be used in any stage of pregnancy. If postpartum and breastfeeding, it is not advisable to breastfeed during treatment. May resume breastfeeding after 24 hours of the last dose.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Cervicitis	Azithromycin 1g single dose	Doxycycline 100mg PO q12h for 7 days	
Postpartum mastitis Common organisms: • <i>Staphylococcus aureus</i> (MSSA) • <i>Streptococcus pyogenes</i> (Grp A, B) • <i>Escherichia coli</i> • <i>Bacteroides</i> sp • <i>Corynebacterium</i> sp • CoNS	Outpatient: Cephalexin 500mg PO q6h for 5-7 Days		Duration of therapy for 5-7 days may be adequate but if poor response consider extending to 10-14 days.
	Inpatient: Cloxacillin 2g IV q6h	Cefazolin 1-2g IV q8h	Milk culture for less severe infection. If severe infection (hemodynamic instability), blood culture required.
Post Lower Segment Cesarean Section (LSCS) infection	In mild surgical site infections (SSI), antibiotics are generally not indicated. Appropriate dressing is the primary treatment.		
	Cloxacillin 2g q6h OR Cefazolin 1-2g IV q8h	Risk of Gram-negative or anaerobic infection (eg: diabetes): Ampicillin/sulbactam 3g IV q6-8h	
References: 1. Savaris RF, de Moraes GS, Cristovam RA, Braun RD. Are antibiotics necessary after 48 hours of improvement in infected/septic abortions? A randomised controlled trial followed by cohort study. <i>Am J ObstetGynecol</i>.2011;204(4):301.e1.Epub 2010 Dec31 2. Mackeen A, Packard RE, Ota E, Speer L. Antibiotic Regimens for postpartum endometritis. <i>Cochrane Database Syst Rev</i> 2:2015 3. Prevention of early-onset Group B Streptococcal Disease in newborns. Committee Opinion No. 485.American College of Obstetricians and Gynecologists. <i>ObstetGynecol</i> 2011;117:1019-27 4. Hughes RG, Brocklehurst P, Steer PJ, Heath P, Stenson BM on behalf of the Royal College of Obstetricians and Gynaecologists. Prevention of 5. Early-onset neonatal group B streptococcal disease. Green-top Guideline No. 36. <i>BJOG</i> 2017;124:e280–e305. 6. NICE Guideline.Premature Labor and birth.https://www.nice.org.uk/guidance/ng25/resources/preterm-labour-and-birth			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
7. Hemsell DL, Little BB, Faro S et al. Comparison of three regimens recommended by the Centers for Disease Control and Prevention for the treatment of women hospitalised with acute pelvic inflammatory disease. <i>Clin Infect Dis</i> 1994;19(4):720-727 8. UK National Guideline for Management of Pelvic Inflammatory Disease 2011. Clinical Effectiveness Group British Association for Sexual Health and HIV. 2011 9. World Health Organization; Mastitis: causes and management; Geneva, Switzerland; accessed21/3/2018 10. Workowski KA, Bolan GA, Centers for Disease Control and Prevention: Sexually transmitted diseases treatment guidelines, 2015. <i>MMWR Recomm Rep</i> 64:1, 2015 11. Antibiotic prophylaxis for prevention of postpartum perineal wound complications: a randomised controlled trial, 2008 12. WHO guidelines for the management of postpartum haemorrhage and retained placenta, 2009			

OCULAR INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Dose alteration may be needed for systemic and intravitreal antibiotics in paediatric patients.			
Blepharitis <i>Staph. aureus</i> <i>Staph. epidermidis</i>	Eyelid hygiene/scrubs is the mainstay of therapy Topical antibiotics are not indicated as an initial therapy	Oxytetracycline with Polymyxin B eye ointment applied q12h to the lid margin OR Fusidic Acid 1% eye ointment applied q12h to the lid margin	
Meibomian gland dysfunction	Warm compression and massage Systemic therapy is not indicated as an initial therapy	In resistant cases, Doxycycline 100mg PO q12h for 4-6 weeks OR Azithromycin 500mg PO q24h for 3 days	Tetracyclines are contraindicated in children < 8 years old
Internal Hordeolum with Secondary Infection <i>Staph. aureus</i> Systemic antibiotics are indicated in the presence of superficial cellulitis or abscess	Warm compresses Cloxacillin 500mg PO q6h for 5 days	- Amoxicillin/clavulanate 625mg PO q8h for 5 days	
External Hordeolum (Stye) <i>Staph. aureus</i>	Epilation of affected eye lash and warm compresses No antibiotic recommended as condition is self-limiting		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
In the presence of superficial cellulitis or abscess	Cloxacillin 500mg PO q6h for 5 days	Amoxicillin/clavulanate 625mg PO q8h for 5 days	
Bacterial Conjunctivitis <i>Staph. aureus</i> <i>Strep. pneumonia</i> <i>Haemophilus influenzae</i>	Chloramphenicol 0.5% eye drop q4h	Levofloxacin 0.5% eye drop q6h OR Moxifloxacin 0.5% eye drop q6h OR Ciprofloxacin 0.3% eye drop q6h	
Gonococcal Conjunctivitis (including neonates) <i>Neisseria gonorrhoeae</i>	Requires systemic therapy. Refer to Sexually Transmitted Infections		Copious irrigation with topical saline drops or artificial tears every 30-60 minutes. Topical antibiotic may considered as ancillary therapy.
Bacterial Keratitis No Growth	Ciprofloxacin 0.3% eye drop q1-2h OR Moxifloxacin 0.5% eye drop q1-2h OR Levofloxacin 0.5% eye drop q1-2h	*Cefuroxime 5% eye drop q1-2h PLUS *Gentamicin 0.9% or 1.4% eye drop q1-2h	Commence a loading dose of one drop every 15 minutes for 3 hours followed by hourly drops around the clock. Taper based on clinical response. *Prepared ready to use extemporaneous by using injectable forms
Contact Lens Related Bacterial Keratitis No Growth	Ciprofloxacin 0.3% eye drop q1-2h OR Levofloxacin 0.5% eye drop q1-2h	*Gentamicin 0.9% or 1.4% eye drop q1-2h Plus *Ceftazidime 5% eye drop q1-2h	Commence a loading dose of one drop every 15 minutes for 3 hours followed by hourly drops around the

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			clock. Taper based on clinical response. *Prepared ready to use extemporaneous by using injectable forms
Bacterial Keratitis Gram-positive cocci Gram-negative rods	Moxifloxacin 0.5% eye drop q1-2h Ciprofloxacin 0.3% eye drop q1-2h OR Levofloxacin 0.5% eye drop q1-2h	*Cefuroxime 5% eye drop q1-2h For MRSA * Vancomycin 5% eye drop q1-2h *Gentamicin 0.9% or 1.4% eye drop q1-2h PLUS *Ceftazidime 5% eye drop q1-2h	Commence a loading dose of one drop every 15 minutes for 3 hours followed by hourly drops around the clock. Taper based on clinical response. *Prepared ready to use extemporaneous by using injectable forms
Acanthamoeba Keratitis <i>Acanthamoeba</i> sp.	*Chlorhexidine 0.02% eye drop q1-2h	**Propamidine isethionate 0.1% q1-2h	Topical therapy tapered with response over a duration of 6-12 months. *Prepared ready to use extemporaneous by using injectable forms **Requires special medicine approval

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Fungal Keratitis Filamentous Fungi/Yeast	*Fluconazole 0.2% eye drop q1-2h OR *Amphotericin B 0.15-0.2% eye drop q1-2h	*/**Voriconazole 1% eye drop q1-2h OR **Natamycin 5% eye drop q1-2h <u>Oral Therapy:</u> May be considered in the absence of contraindications Ketoconazole 200mg PO q24h OR Fluconazole 200mg PO q24h	Natamycin is the choice therapy for <i>Fusarium</i> . Amphotericin B is the choice therapy for candida. In severe fungal keratitis, combination therapy may be used. *Prepare ready to use extemporaneous **Requires special medicine approval
Herpes simplex Keratitis Herpes simplex Type 1 & 2 Epithelial Keratitis In the presence of stromal or endothelial disease Recurrent Herpes Simplex Stromal Keratitis	Acyclovir 3% eye ointment 5 times/day Acyclovir 400mg PO 5 times/day for 7-14 days Prophylaxis: Acyclovir 400mg PO q12h for 12 months		For stromal and endotheliitis: Combination with intensive topical steroid is required.
Herpes zoster Ophthalmicus Herpes zoster virus	Needs systemic therapy. Refer to Skin & Soft Tissue Infections Section		
Ocular Toxoplasmosis <i>Toxoplasma gondii</i>	Trimethoprim/sulfamethoxazole 160/800mg PO q12h for at least 6 weeks	*Pyrimethamine 25-50mg PO q24h PLUS	Pregnancy : May consider Intravitreal Clindamycin 1.0mg /0.1ml

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
		Folinic acid 10-25mg PO q24h PLUS *Sulfadiazine 1g PO q6h OR Azithromycin 500mg PO q24h OR Clindamycin 300mg PO q6h for 3-4 weeks, then 150mg q6h PO for 3-4 weeks	Systemic steroids are usually indicated in immunocompetent patients *Requires special medicine approval Prophylaxis for recurrent lesions: Trimethoprim/ulfamethoxazole 80/400mg q12h PO 3 times a week
Acute Retinal Necrosis Herpes simplex virus	Acyclovir 10mg/kg/dose (not more than 800mg) IV q8h for 10-14 day Followed by Acyclovir 800mg PO 5 times/day for 6 weeks	*Valacyclovir 1g PO q8h for 6 weeks	*Requires special medicine approval Systemic steroid is indicated depending on location or severity of the infection.
CMV Retinitis <i>Cytomegalovirus</i>	Systemic therapy Ganciclovir 5mg/kg IV q12h for 2-3 weeks Intravitreal therapy Intravitreal Ganciclovir 2mg/0.1ml biweekly	Systemic therapy Valganciclovir 900mg PO q12h for 3 weeks (induction) Followed by 900mg PO q24h (maintenance) Intravitreal therapy Intravitreal *Foscarnet 2.4mg/0.1ml (1-2 weekly)	Systemic therapy is indicated in all cases. Maintenance may need to continue until CD4 count is > 150 cells/mm ³ for 3 consecutive months. Intravitreal therapy is indicated in zone 1 and 2 lesions. Intravitreal to be tapered according to clinical response.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			Ganciclovir implant: 4.5g an option for prolonged use of intravitreal Ganciclovir *Requires special medicine approval
Ocular Syphilis <i>Treponema pallidum</i>	Ocular Syphilis (syphilitic uveitis) should be treated as Neurosyphilis. Refer to Sexually Transmitted Infections Section		Referral to Dermatologist/ID Physician
Ocular Tuberculosis <i>Mycobacterium tuberculosis</i>	Needs systemic therapy Refer to MOH CPG on Management of Tuberculosis (Extra-pulmonary TB) Section *Ethambutol may cause optic neuropathy and should avoided depending on the case		Ocular TB: presents as a unilateral/ bilateral infective uveitis characterized by multifocal choroiditis/ granuloma and there may be supportive FFA findings of occlusive vasculitis. The diagnosis may be clinical as vitreous sampling for AFB or TB PCR may not be very sensitive due to small sample size and sensitivity of the tests. Clinical response to anti-TB is often diagnostic. Uveitis secondary to TB Hypersensitivity is an immune response to acid fast bacilli in the eye and manifests predominantly as an inflammatory uveitis. Treatment includes anti-TB in combination with an immunosuppressive dose of systemic steroids for at least 6-9 months.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			Systemic steroid may be indicated but is only for - non-active systemic TB - immunocompetent patients -severe ocular inflammation developing after starting anti-TB treatment - vision threatening condition Systemic steroid should NOT be started alone without anti-TB treatment
Post-operative Bacterial Endophthalmitis <i>Staphylococcus epidermidis</i> <i>Staphylococcus aureus</i> <i>Pseudomonas aeruginosa</i> , <i>Bacteroides</i> Species <i>Streptococcus pneumoniae</i> , <i>Alpha-Haemolytic streptococci</i>	Intravitreal antibiotic injections Vancomycin 1-2mg/0.1ml PLUS Ceftazidime 2mg/0.1ml If suspicious of fungal endophthalmitis: ADD Intravitreal Amphotericin B 0.005mg/0.1ml OR Topical treatment-options: (monotherapy or combination) *Ceftazidime 5% eye drop *Vancomycin 5% eye drop *Gentamicin 1.4% eye drop *Cefuroxime 5% eye drop	Intravitreal antibiotic injections: Vancomycin 1-2mg/0.1ml PLUS Amikacin 0.4mg/0.1ml	Endogenous endophthalmitis: Treatment is based on primary infection (bacterial/fungal etc) and culture and sensitivity results. Systemic antibiotics are indicated in severe, virulent endophthalmitis. Repeat intravitreal antibiotics after 48 to 72 hours if indicated. *Prepared ready to use extemporaneous by using injectable forms

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	Moxifloxacin 0.5% eye drop Levofloxacin 0.5% eye drop OR Systemic treatment Ciprofloxacin 750mg PO q12h for 10 days For culture negative cases, ADD Clarithromycin 250-500mg PO q12h for 7-14 days	Systemic treatment Vancomycin 15-20mg/kg IV q8-12h (not to exceed 2g per dose) PLUS Ceftazidime 1-2g IV q8h	
Post-operative Fungal Endophthalmitis	Intravitreal Amphotericin B 0.005mg/0.1ml OR Systemic Therapy Fluconazole 200mg PO q24h for 6 weeks (minimum)	*Intravitreal Miconazole 0.01mg/0.1ml OR *Intravitreal Voriconazole 50-100 ug/0.1ml OR Systemic Therapy *Voriconazole 200mg PO q12h	Intravitreal and Systemic therapy are indicated in all cases *Requires special medicine approval
Endogenous Endophthalmitis	Systemic treatment Ciprofloxacin 750mg PO q12h for 10 days For culture negative cases: ADD Clarithromycin 250-500mg PO q12h for 7-14 days	Systemic treatment Vancomycin 15-20mg/kg IV q8-12h (not to exceed 2g per dose) PLUS Ceftazidime 1-2g IV q8h	Treatment is based on primary infection (bacterial/fungal etc) and culture and sensitivity results. All cases require systemic therapy. Intravitreal injection is indicated in cases with vitreous involvement and sight threatening choroidal lesions.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	OR Topical treatment-options (monotherapy or combination) *Ceftazidime 5% eye drop *Vancomycin 5% eye drop *Cefuroxime 5% eye drop Gentamicin 0.3% eye drop Moxifloxacin 0.5% eye drop Levofloxacin 0.5% eye drop OR Intravitreal antibiotic injections: Vancomycin 1-2mg/0.1ml PLUS Ceftazidime 2mg/0.1ml If suspicious of fungal endophthalmitis, ADD Intravitreal Amphotericin B 0.005mg/0.1ml	Intravitreal antibiotic injections: Vancomycin 1-2mg/0.1ml PLUS Amikacin 0.4mg/0.1ml	Topical therapy may supplement therapy. Not to use systemic steroids in these cases Review antibiotic regimen after microbiology results. Repeat intravitreal antibiotics after 48 to 72 hours if indicated *Prepared ready to use extemporaneous by using injectable forms
Dacryocystitis <i>Strep pneumoniae</i> <i>Staph aureus</i> Gram-positive anaerobes	Cefuroxime 250mg PO q12h for 7 days	Amoxicillin/clavulanate 625mg PO q8h for 7 days	Consider intravenous antibiotics in severe infections
Preseptal Cellulitis <i>Strep pneumoniae</i> <i>Staph aureus</i> <i>Streptococcus</i> sp.	Cloxacillin 500mg-1g PO q6h for 5 days	Amoxicillin/clavulanate 625mg PO q8h for 7 days OR Ceftriaxone 1-2g IV q24h	Consider intravenous antibiotics in severe infections

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Orbital Cellulitis/abscess <i>Strep pneumoniae</i> <i>Staph aureus</i> <i>Streptococcus</i> sp. Gram-positive anaerobes	Amoxicillin/clavulanate 1.2g q8h IV for 7-10 days	Ceftriaxone 1-2g q24h IV for 7-10 days If anaerobes suspected: ADD Metronidazole 500mg IV q8h for 7-10 days	
References: 1. Sobrin L, Kump L, Foster CS. Intravitreal clindamycin for toxoplasmicretinochoroiditis Retina 2007. Sep; 27(7): 952-7. 2. Patrick MKT, Claire Y H, Susan L. Antiviral selection in the management of acute retinal necrosis. Clinical Ophthalmology 2010;4 11–20 3. Peter R, Jost H, Livia G, et al. Virus Diagnostics and Antiviral Therapy in Acute Retinal Necrosis (ARN). Antiviral Drugs – Aspects of Clinical Use and Recent Advances. Intechopen. 4. MN Muthiah, M Michaelides, CS Child, et al. Acute retinal necrosis: a national population-based study to assess the incidence, methods of diagnosis, treatment strategies and outcomes in the UK. Br J Ophthalmol2007;91:1452–1455 5. Simon RJT, Robin H, Claire YH, Sue Lightman. Valacyclovir in the treatment of acute retinal necrosis. BMC Ophthalmology 2012, 12:48. 6. Robert WW, Emmett TC et al. Diagnosing and Managing Acute Retinal Necrosis. Retinal Physician. 7. Helm CJ, Holland GN. Ocular tuberculosis. SurvOphthalmol. 1993 Nov-Dec;38(3):229-56 8. Bodaghi B1, LeHoang P. Ocular tuberculosis. CurrOpinOphthalmol. 2000 Dec;11(6):443-8 9. CPG for Management of Post- Operative Endophthalmitis, Ministry of Health Malaysia, August 2006 10. Periorbital and orbital cellulitis: A 10 year review of Hospitalized children. <i>Eur J Ophthalmol</i> 2010;20(6): 1066-1072 11. Microbiology and Antibiotic Management of Orbital Cellulitis <i>Pediatrics</i> 2011;127:e566			

ORAL/DENTAL INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
A. ANTIMICROBIAL USE FOR BACTERIAL INFECTIONS			
1. Infections of the Teeth and Supporting Structures			
Reversible/ Irreversible Pulpitis	Systemic antibiotic use not recommended	Systemic antibiotic use not recommended	
Localized Dentoalveolar Abscess	Superficial Systemic antibiotic use not recommended (unless medically compromised) Deep Infection/Medically Compromised Amoxicillin 500mg PO q8h PLUS Metronidazole 400mg PO q8h OR Amoxicillin/clavulanate 625mg PO q8h	Systemic antibiotic use not recommended (unless medically compromised) Penicillin Allergy: Clindamycin 300mg PO q6h	
Dry Socket	Systemic antibiotic use not Recommended	Systemic antibiotic use not recommended	Local treatment with saline irrigation and antiseptic/ analgesic dressings and symptomatic relief of pain
Localized Pericoronitis	Systemic antibiotic use not recommended in absence of regional or systemic signs and symptoms	Systemic antibiotic use not recommended in absence of regional or systemic signs and symptoms	Local treatment with antiseptic irrigation and mouthwash and symptomatic relief of pain

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Chronic Gingivitis	Systemic antibiotic use not recommended		First line treatment: Mechanical and chemical plaque control. 0.2% Aqueous Chlorhexidine Gluconate not be used alone but as an adjunct to mechanical debridement. Second line treatment-Antimicrobial mouth rinse
Chronic Periodontitis Antibiotic can be considered in cases of: 1. Unresponsive to conventional mechanical therapy 2. Acute infection with systemic manifestation 3. Medically compromised	Systemic antibiotic use is generally not recommended. Amoxicillin 500mg PO q8h PLUS Metronidazole 400mg PO q8h OR Amoxicillin/clavulanate 625mg PO q8h	Systemic antibiotic use is generally not recommended. Penicillin Allergy: Clindamycin 300mg PO q6h	First line treatment: Mechanical plaque control
Aggressive Periodontitis <i>Aggregatibacter actinomycetemcomitans</i> <i>Porphyromonas gingivalis</i> <i>Tannerella forsythensis</i> <i>Prevotella intermedia</i> <i>Spirochaetes</i>	Amoxicillin 500mg PO q8h PLUS Metronidazole 400mg PO q8h	Penicillin Allergy: Clindamycin 300mg PO q6h	Antibiotics are not used alone but are used as an adjunct to scaling and root debridement.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Local missed Periodontal Abscess	Systemic antibiotic use not recommended		Incision and Drainage Management of cause of abscess and symptomatic relief of pain
2. Infections of the Jaws			
Osteomyelitis of the jaws of dental origin Different organisms may be involved	For acute cases , start with: Amoxicillin 500mg PO q8h PLUS Metronidazole 400mg PO q8h OR Amoxicillin/clavulanate 625mg PO q8h	Penicillin Allergy: Clindamycin 300-450mg PO/IV q6h	Culture and sensitivity is necessary to guide the antibiotic. For chronic cases , start with surgical treatment first. Antibiotics only when causative organisms are identified. Duration of antibiotic therapy can be 4-6 weeks depending on patient response/microbiological clearance of the pathogen.
3. Spreading Infections and Infections of Fascial Spaces (with/without Systemic Signs)			
Cellulitis ± Abscess of dental origin Viridans Streptococci <i>Staphylococci</i> <i>Prevotella</i> <i>Peptostreptococcus</i> <i>Fusobacterium nucleatum</i> <i>Clostridium</i> sp	Benzylpenicillin 2-4MU IV q4-6h PLUS Metronidazole 500mg IV q8h OR Amoxicillin/clavulanate 1.2g IV q6-8h (not more than 1.2g in a single dose- max 7.2g daily) OR	Penicillin Allergy: Clindamycin 300-450mg PO/IV q6h	Empirical antibiotics are started. Incision and drainage is advised and antibiotics is changed in accordance with result of culture and sensitivity.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Surgical site infection & Traumatic wound infection (Infection is usually by endogenous organisms rather than exogenous) <i>Viridans Streptococci</i> <i>Staphylococci</i> <i>Prevotella</i> <i>Peptostreptococcus</i> , <i>Eubacterium</i> , <i>Fusobacterium nucleatum</i>	Cefuroxime 750mg-1.5g IV q8h PLUS Metronidazole 500mg IV q8h OR If not responding to above antibiotics, Ceftriaxone 1-2g IV q24h PLUS Metronidazole 500mg IV q8h <u>Step Down/Oral therapy</u> Amoxicillin 250-750mg PO q8h PLUS Metronidazole 400mg PO q8-12h OR Amoxicillin/clavulanate 625mg PO q8h OR Cefuroxime 250-500mg PO q12h PLUS Metronidazole 400mg PO q8-12h	Penicillin Allergy: Clindamycin 300-450mg PO q6h	
Traumatic wound involving skin/Infection of skin origin	Cloxacillin 500mg-1g IV q6h (in skin involvement - if <i>Staph.</i> expected) OR Amoxicillin 250-750mg PO q8h PLUS Metronidazole 400mg PO q8-12h	Penicillin Allergy: Clindamycin 300-450mg IV/PO q6h	

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
4. Post-implant Infections (“Peri-implantitis”)			
Causative Organisms: <i>Actinomyces sp.</i> <i>Eubacterium sp.</i> <i>Propionibacterium sp.</i> <i>Lactobacillus sp.</i> <i>Veillonella sp.</i> <i>Porphyromonas gingivalis</i> <i>Prevotella intermedia</i> <i>Fusobacterium nucleatum</i>	Amoxicillin/clavulanate 625mg PO q8h OR Amoxicillin 500mg PO q8h PLUS Metronidazole 400mg PO q8h	Penicillin Allergy: Doxycycline 100mg PO q12h OR Clindamycin 300mg PO q6h	Bacteria associated with periimplantitis are extremely resistant to antibiotics. Antibiotics are not used alone but are used as an adjunct to local mechanical and chemical debridement. Also irrigation with chlorhexidine and optimal oral hygiene by patient. Locally delivered antibiotics is preferred compared to systemic administration. Currently there is no reliable study to suggest most effective antibiotic therapy.
References: 1. Antibiotics and The Treatment of Endodontic Infections. Endodontics colleagues for Excellence 2006; American Association of Endodontics 2. JCan Dent Assoc 2003 Nov 69 (10):660 Clin.Microbiol.Rev.2013,26(2):255 3. Med Oral Patol Oral Cir Bucal 2005; 10:77-85 4. JClinMicrobiol.2003;41(12):5794-7 Journal of the Irish Dental Association 2009; 55 (4): 190 – 192 5. Clinical Periodontology-12thed.2014 6. Clinical Periodontology-9thed.2002 7. Periodontology 2000, Vol. 62, 2013, 218-231 CPG Management of chronic periodontitis Nov 2012 MOH,Malaysia 8. JClin Periodontol.2012;39:284-294 9. Clin Periodontol.2011;38:43-49 10. J Clin Periodontol 2008; 35: 696–704			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
11. J Periodont Res 2012; 47: 137–148 12. Periodontology 2000. Jun2014, Vol. 65 Issue 1, p149-177. 29p. Malaysian Dental Journal (2008) 29(2) 154-157 13. CPG Management of periodontal abscess- MOH,Malaysia 2003 14. Eur J Oral Implantol 2012; 5 (Suppl): S21-S41 15. Clin Oral Impl Res 2012 (23): 205-210 16. Int.J Oral Maxillofac Implants 2014 (29): 325-345 17. Clin Oral Impl Res 2000:11(suppl): 146-155			
B. ANTIMICROBIAL USE FOR FUNGAL INFECTIONS			
Oral Candidiasis			
Oropharyngeal Candidiasis (Oral Thrush)	Refer to section [Infections in Immunocompromised patients – HIV]		
Acute Pseudomembranous Candidiasis Hyperplastic Candidiasis (Candidal Leukoplakia) Mild disease Moderate to severe disease Fluconazole-refractory disease	Topical antifungal Fluconazole 100-200mg PO q24h for 7-14 days Itraconazole solution 200mg PO q24h	Nystatin (oral suspension) 400,000-600,000units q6h for 7-14 days Voriconazole 200mg PO q12h	
Chronic Erythematous Candidiasis	Local measures - denture cleansers, remove dentures at night		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
(<i>Candida</i> -associated denture stomatitis with and without angular cheilitis)	Soak dentures in chlorhexidine mouthwash 2% Topical antifungals if local measures fail: Nystatin (oral suspension) 500,000-1,000,000units q6h-8h (to continue for 2 days after perioral symptoms disappeared or cultures show eradication of candida sp.)		
References: 1. Am Fam Physician. 2008;78(&):845-852 Journal of Oral Microbiology 2011,3:5771-DOI: 10.3402/jom.v3i0.5771 2. Med Oral Patol Oral Cir Bucal. 2011 Mar 1;16(2):e1 39-43 Australia Dental Journal 2010; 55:(1 suppl):48 3. Clinical Practice Guideline for the management of candidiasis: 2016 update by IDSA			
C. ANTIMICROBIAL USE FOR VIRAL INFECTIONS			
Herpes simplex virus type 1 (HSV-1) -Primary herpetic gingivostomatitis -Herpes labialis Herpes simplex virus type 2 (HSV-2) Epstein-Barr virus E.g.: Infectious mononucleosis, oral hairy leukoplakia	Symptomatic treatment in most cases. Can also consider: 1) Topical Acyclovir 5% cream q4h for 5-10 days in prodromal phase for recurrent herpes labialis 2) Systemic antiviral Acyclovir 400-800mg PO 5 times daily for 7-14 days		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Varicella-zoster virus Coxsackie virus -Herpangina -Hand, foot and mouth disease			
Severe Varicella/Herpes Infections or in immunocompromised patients	Refer to section [Skin & Soft Tissue Infections – Viral Infections]		Varicella Zoster Infections: Advisable to start treatment for early within 48 hours of Varicella Infections Herpes infection: Topical antiviral treatment is not recommended. Advisable to start treatment early within 48-72 hours and for all immunocompromised patient or for immunocompetent with following criteria: - > 50 years years of age - Have moderate to severe pain - Have moderate to severe rash - Have non-truncal involvement
Viral Encephalitis Herpes simplex Varicella zoster	Refer to section [Central nervous system (CNS) Infections – Viral Encephalitis]		Duration: 10-21 days

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
References: 1. Aust Dent J 2005;50 Suppl 2: S31- S35 2. Centers for Disease Control and Prevention (CDC) 2010			

OTORHINOLARYNGOLOGY INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks																																			
	Preferred	Alternative																																				
1. General Sore Throat																																						
The Mslsaac Centor score can be used to help physicians decide which patients need no testing, throat culture/rapid antigen detection testing, or empiric antibiotic therapy. The cumulative score determines the likelihood of streptococcal pharyngitis and the need for antibiotics <table><tr><td>Criteria</td><td>Score</td><td>Age</td><td>Score</td></tr><tr><td>Absence of cough</td><td>1</td><td>3 to 14 years</td><td>1</td></tr><tr><td>Swollen and tender anterior cervical lymph nodes</td><td>1</td><td>15 to 44 years</td><td>0</td></tr><tr><td>Temperature > 100.4° F (38° C)</td><td>1</td><td>45 years and older</td><td>-1</td></tr><tr><td>Tonsillar exudates or swelling</td><td>1</td><td></td><td></td></tr></table> <table><tr><td>Cumulative score:</td><td></td><td></td></tr><tr><td>Total score</td><td>Risk</td><td>Comment</td></tr><tr><td>0 or 1</td><td>Low risk</td><td>Do not require testing or antibiotic therapy</td></tr><tr><td>2 or 3</td><td></td><td>Testing recommended. Positive results warrants antibiotics. If test not available, antibiotics may be considered</td></tr><tr><td>≥4</td><td>High risk</td><td>Perform throat culture. Treat with antibiotic only if culture positive</td></tr></table>				Criteria	Score	Age	Score	Absence of cough	1	3 to 14 years	1	Swollen and tender anterior cervical lymph nodes	1	15 to 44 years	0	Temperature > 100.4° F (38° C)	1	45 years and older	-1	Tonsillar exudates or swelling	1			Cumulative score:			Total score	Risk	Comment	0 or 1	Low risk	Do not require testing or antibiotic therapy	2 or 3		Testing recommended. Positive results warrants antibiotics. If test not available, antibiotics may be considered	≥4	High risk	Perform throat culture. Treat with antibiotic only if culture positive
Criteria	Score	Age	Score																																			
Absence of cough	1	3 to 14 years	1																																			
Swollen and tender anterior cervical lymph nodes	1	15 to 44 years	0																																			
Temperature > 100.4° F (38° C)	1	45 years and older	-1																																			
Tonsillar exudates or swelling	1																																					
Cumulative score:																																						
Total score	Risk	Comment																																				
0 or 1	Low risk	Do not require testing or antibiotic therapy																																				
2 or 3		Testing recommended. Positive results warrants antibiotics. If test not available, antibiotics may be considered																																				
≥4	High risk	Perform throat culture. Treat with antibiotic only if culture positive																																				
References: 1. Empirical Validation of Guidelines for the Management of Pharyngitis in Children and Adults. JAMA. April 7, 2004; Vol 291 No 13: 1588-1595 2. The Infectious Disease Society of America (IDSA) 2012 Guidelines																																						
2. Throat and Upper Respiratory																																						

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks							
	Preferred	Alternative								
Tonsillitis/Pharyngitis Group A <i>Streptococcus</i>	Phenoxymethypenicillin (Penicillin V) 500mg PO q6h or 1g PO q12h for 5-10 days If not tolerated orally: Amoxicillin/clavulanate 1.2g IV q8h OR Ampicillin/sulbactam 3g IV q6h OR Cefuroxime 750mg IV q8h	Amoxicillin 500mg PO q8h for 5-10 days Penicillin Allergy: Erythromycin Ethylsuccinate 800mg PO q12h for 5-10 days								
Acute Peritonsillar Abscess Group A <i>Streptococcus</i> <i>Staphylococcus aureus</i> <i>Haemophilus influenzae</i> <i>Fusobacterium necrophorum</i>	Ampicillin/sulbactam 3g IV q6h OR Amoxicillin/clavulanate 1.2g IV q8h Penicillin Allergy: Clindamycin 600mg IV q8h	Amoxicillin/clavulanate 625 mg PO q8h Penicillin Allergy: Clindamycin 300-450mg PO q6h	Abscess to be drained							
Diphtheria <i>Corynebacterium diphtheriae</i>	Antitoxin PLUS Benzylpenicillin 50,000 units/kg to a maximum of 1.2 MU IV q12h, followed by Phenoxymethylpenicillin (Penicillin V) 250mg PO q6h for total of 14 days		Diphtheria antitoxin: <table><tr><td>Pharyngeal/ laryngeal disease of 2 days duration</td><td>20,000-40,000 units</td></tr><tr><td>Nasopharyngeal disease</td><td>40,000-60,000 units</td></tr><tr><td>Systemic disease of ≥3 days or any patient with diffuse neck swelling</td><td>80,000-120,000 units</td></tr></table>		Pharyngeal/ laryngeal disease of 2 days duration	20,000-40,000 units	Nasopharyngeal disease	40,000-60,000 units	Systemic disease of ≥3 days or any patient with diffuse neck swelling	80,000-120,000 units
Pharyngeal/ laryngeal disease of 2 days duration	20,000-40,000 units									
Nasopharyngeal disease	40,000-60,000 units									
Systemic disease of ≥3 days or any patient with diffuse neck swelling	80,000-120,000 units									

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			Administer over 60 minutes to inactivate toxin rapidly
Acute Epiglottitis <i>Haemophilus influenzae</i> Type B virus, <i>Streptococcus pneumoniae</i>	Ceftriaxone 2g IV q24h PLUS Metronidazole 500mg IV q8h OR Ampicillin/sulbactam 3g IV q6h <u>Oral step down therapy:</u> Amoxicillin/clavulanate 625mg PO q8h for 7–14 days	Penicillin Allergy: Clindamycin 600-900mg IV q8h PLUS Ciprofloxacin 400mg IV q12h	Urgent hospitalisation. May present with life threatening upper airway obstruction, especially in paediatrics.
Deep Neck Space Abscess <i>Streptococcus pyogenes</i> <i>Staphylococcus aureus</i> <i>Fusobacterium necrophorum</i>	Ampicillin/sulbactam 3g IV q6h OR Ceftriaxone 2g IV q24h PLUS Metronidazole 500mg IV q6h		Duration: 10-14 days
3. Rhinology			
Acute Bacterial Rhinosinusitis (ABRS) <i>Streptococcus pneumoniae</i> , <i>Haemophilus influenzae</i> , <i>Moraxella catarrhalis</i>	Ampicillin/sulbactam 375mg PO q12h OR Amoxicillin/clavulanate 625mg PO q8h Duration: 5-7 days	Penicillin allergy: Doxycycline 100mg PO q12h for 5-7 days <u>Pregnant patients with penicillin allergy:</u> Azithromycin 500mg PO q24h for 3 days	Consider antibiotic if present at least 3 of below: - Purulent/ greenish nasal discharge - Severe local pain (VAS 8-10) - Fever - Elevated ESR/CRP

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	If no improvement after 3 days of oral antibiotics, refer to ENT.		- Double sickening (becoming worse after initial recovery) VAS: Visual Analogue Score
Chronic Rhinosinusitis (CRS) (with or without nasal polyps) <i>Staphylococcus aureus</i> <i>Enterobacterales</i> spp <i>Pseudomonas</i> spp	There is limited evidence that antibiotics, as a single therapy, are beneficial in the treatment of CRS. For most patients, antibiotics represent one component of comprehensive medical management. It is important to note that antibiotics should be used only in patients who meet the diagnostic criteria for CRS. In refractory patients, some suggest a 2-weeks course (may up to 6 weeks in selected patients) of macrolide therapy. The data are mixed and the potential for adverse effects may outweigh any potential benefit: Clarithromycin 500mg PO q24h for 2 weeks, then 250mg PO q24h (up to 6 weeks in selected patients)		Randomised controlled trials on the efficacy of long-term antibiotics in large and well defined populations with CRS are lacking A more likely patient to respond has signs and symptoms dominated by a neutrophilic inflammation such as purulence, normal IgE and a low CT score. As a precaution, macrolides should be avoided in patients with a long QT interval.
4. Otology			
Acute otitis media (AOM) <i>Streptococcus pneumoniae</i> , <i>Haemophilus influenzae</i> , <i>Moraxella catarrhalis</i>	Amoxicillin/clavulanate 625mg PO q8h for 5 days	Penicillin Allergy: Azithromycin 500mg PO on day 1, followed by 250mg PO q24h until day 5	If symptoms not resolving after 48-72 hours, refer ENT
Malignant Otitis Externa/ Necrotizing Otitis Externa <i>Pseudomonas aeruginosa</i>	Ciprofloxacin 400mg IV q8h for 2 weeks OR Ceftazidime 2g IV q8h for 2 weeks		Prolong treatment of malignant otitis externa for 6-8 weeks is generally recommended, as indicated for osteomyelitis.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	followed by (once showing clinical response, consider switching to oral therapy): Ciprofloxacin 750mg PO q12h for 6 weeks		
Acute Diffuse Otitis Externa <i>Pseudomonas aeruginosa</i> <i>Staphylococcus aureus</i>	Ofloxacin 0.3% otic solution Instil 10 drops into affected ear(s) q24h for 7 days	Hydrocortisone 10mg, neomycin 3.5mg, polymyxin B 10,000U per ml ear drops Instil 3-4 drops into affected ear(s) q6-8h for max 10 days	Aural toileting required in discharging ears Not to use ear drop in perforated ear drum
Chronic Suppurative Otitis Media <i>P. aeruginosa</i> <i>Staphylococcus aureus</i>	Ofloxacin 0.3% otic solution Instil 10 drops into affected ear(s) q12h for 10-14 days		
Otomycosis <i>Aspergillus</i> sp.	Clotrimazole 1% ear solution, applied q12h for 10-14 days		Aural toileting required. Not to use ear drop in perforated ear drum
References: 1. Diagnosis and Management of Group A Streptococcal Pharyngitis: 2012 Update by Infectious Diseases Society of America, Clin Infect Dis 2012;55:86-102 2. Use of Diphtheria Antitoxin (DAT) for Suspected Diphtheria Cases, CDC 2016 3. Fokkens WJ, Lund VJ, Mullol J et al. EPOS 2012. European position paper on rhinosinusitis and nasal polyps 2012. A summary for otorhinolaryngologists. Rhinology 2012 Mar; 50(1): 1 4. Sore Throat (Acute): Antimicrobial Prescribing (NG84), NICE 2018 5. Stanford T. Shulman, Alan L. Bisno, Herbert W. Clegg, Michael A. Gerber, Edward L. Kaplan, Grace Lee. et al. Clinical Practice Guidelines for the 6. Grandis JR. Malignant (necrotizing) external otitis. Uptodate assessed 13/10/22			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
		*Levofloxacin 500-750 mg IV/PO q24h for 5-7 days	
2. Viral Pneumonia			
i. Influenza	Oseltamivir 75mg PO q12h for 5 days		
ii. <i>Varicella zoster</i>	Acyclovir 10mg/kg IV q8h for 7 days		
3. Lung Abscess and Empyema			
Empirical	Amoxicillin/clavulanate 1.2g IV q6-8h	Ceftriaxone 2g IV q24h PLUS *Metronidazole 500 mg IV q8h <u>Penicillin Allergy</u> Clindamycin 600 mg IV/PO q6H	Duration of treatment: -Drained abscess/empyema may require 2-4 weeks of antibiotics -Undrained abscess/empyema may require 4-6 weeks of antibiotics Lung empyema: Attempts should be made to drain the collection May change to oral regime once clinical improvement seen *Metronidazole: in cases of lung abscess when aspiration is suspected.
<i>Staphylococcus aureus</i>	Cloxacillin 2g IV q4-6h	Cefazolin 2g IV q8h	Duration: 4-6 weeks, depending on clinical response. In rare cases (slow response to antibiotics) may need prolonged therapy.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			May change to oral therapy (e.g. Amoxicillin/clavulanate 625mg PO q8h) to complete the duration once patient stabilised and improved
4. Infective Exacerbation of Chronic Obstructive Pulmonary Disease (COPD)			
Antibiotics only considered if there is increased purulence in sputum AND one of the following: - Increased sputum volume - Increased dyspnoea OR Patient intubated			
Outpatient	Amoxicillin/clavulanate 625mg PO q8h for 5-7 days	Cefuroxime 500mg PO q12h for 5-7 days OR Doxycycline 100mg PO q12h for 5-7 days	
Inpatient	Amoxicillin/clavulanate 1.2g IV q8h for 5-7 days PLUS/MINUS Azithromycin 500mg IV/PO q24h for 3-5 days	Ceftriaxone 2g IV q24h for 5-7 days PLUS/MINUS Azithromycin 500 mg IV/PO q24h for 3-5 days	
*If suspect <i>Pseudomonas</i> infection	**Piperacillin/tazobactam 4.5g IV q6-8h OR Cefepime 2g IV q8h PLUS/MINUS Azithromycin 500 mg IV/PO q24h for 3-5 day	Ceftazidime 2g IV q8h PLUS/MINUS Azithromycin 500 mg IV/PO q24h for 3-5 day	* <i>Pseudomonas</i> sp risk factors: 1. Frequent exacerbation 2. Severe airflow limitation 3. Exacerbation requiring mechanical ventilation

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			**Piperacillin/tazobactam: If given as q8h, to be given as extended infusion (over 3-4 hours).
5. Hospital-acquired Pneumonia (HAP/VAP)			
Risk factors for multidrug resistance (MDR) organisms: 1. Prior intravenous antibiotic use within 90 days 2. More than 5 days of hospitalization in high risk ward (ICU, HDU) 3. Previous colonization with MDR pathogens Risk of MDR organisms is lower with early onset HAP/VAP.			
Early Onset HAP/VAP (2-4 days of admission/intubation)	Amoxicillin/clavulanate 1.2g IV q8h for 5-7 days	Ceftriaxone 2g IV q24h for 5-7 days	Need to adjust to local antibiogram/ prevalent organisms
Late Onset HAP/VAP (5 days or more of admission/ intubation) Causative organisms are determined by local prevalence.	*Piperacillin/tazobactam 4.5g IV q6-8h for 7 days OR Cefepime 2g IV q8h for 7 days	Imipenem/cilastatin 500mg IV q6h for 7 days OR Meropenem 1g IV q8h for 7 days	Ideal empirical antibiotic coverage depends on local prevalence of organisms. Duration of antibiotics could be shortened to 7 days even for MDR <i>Pseudomonas aeruginosa</i> and <i>Acinetobacter baumannii</i> infections. Longer duration may be indicated depending upon clinical, radiological and laboratory parameters. To de-escalate antibiotics according to culture and sensitivity results.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			*Piperacillin/tazobactam: If given as q8h, to be given as extended infusion (over 3-4 hours).
6. Aspiration Pneumonia			
	Amoxicillin/clavulanate 1.2g IV q8h	Ceftriaxone 2g IV q24h PLUS Metronidazole 500mg IV q8h	Duration: 7- 10 days To switch to oral therapy when clinical condition improves and patient is able to tolerate orally. Antibiotics are not indicated for aspiration (chemical) pneumonitis.
References: 1. National Antimicrobial Guideline 2019.			

SEXUALLY TRANSMITTED INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Syphilis (<i>Treponema pallidum</i> Infection)			
Early Syphilis (< 2 years) • Primary Syphilis • Secondary Syphilis • Early Latent Syphilis (History of syphilis infection in the last 2 years)	Benzathine Penicillin 2.4MU IM stat OR Procaine Penicillin G 600,000units IM q24h for 10 days	Penicillin Allergy: Ceftriaxone 1g IM/IV q24h for 10 days (if no anaphylaxis to penicillin) OR Doxycycline 100mg PO q12h for 14 days	If drug administration is interrupted for ≥ 1 day at any point during the treatment course, it is recommended that the entire course is restarted. Contact tracing: Examine and investigate sex partner and treat when indicated Follow up: Clinical and serologic evaluation should be performed at 6 and 12 months after treatment; more frequent evaluation might be prudent if opportunity for follow-up is uncertain or if repeat infection is a clinical concern. Jarisch-Herxheimer reaction: To treat with Paracetamol
Late Syphilis (> 2 years) • Late Latent Syphilis • Tertiary syphilis (Gumma) • *Cardiovascular syphilis	Benzathine Penicillin 2.4MU IM weekly for 3 weeks (Day 1, 8 & 15) OR Procaine Penicillin G 600,000units IM q24h for 14 days	Penicillin Allergy: Doxycycline 100mg PO q12h for 28 days	Contact tracing: Examine and investigate sex partner and treat when indicated Follow up: 6 monthly for 2 years and yearly until RPR negative or stable at a low titre RPR 1:4 or less

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			If a patient defaults Benzathine Penicillin treatment by $\geq$ two weeks in between the weekly doses, the whole regime need to be restarted. *For cardiovascular syphilis: Consider Prednisolone 40-60mg PO q24h for 3 days commencing 24h before the antibiotics.
Neurosyphilis	Benzylpenicillin 4MU q4h IV for 10-14 days OR Procaine penicillin G 2.4MU IM q24h PLUS Probenecid 500mg PO q6h (both for 10–14 days)	Penicillin Allergy: Doxycycline 200mg PO q12h for 28 days OR Ceftriaxone 2g IM q24h (lidocaine as diluent) for 10-14 days (if no anaphylaxis to penicillin) OR Ceftriaxone 2g IV q24h (WFI as diluent) for 10-14 days (if no anaphylaxis to penicillin)	Contact tracing: Examine and investigate sex partner and treat when indicated CSF examination should be done in 1. Patients with neurological and/or ocular symptoms or signs. 2. Nontreponemal test titres do not decrease by fourfold within 12 months of therapy All persons who have neurosyphilis, ocular syphilis, or otosyphilis should be tested for HIV at the time of diagnosis. Consider corticosteroid cover to prevent the Jarisch-Herxheimer reaction: Prednisolone 40-60mg PO q24h for 3 days commencing one day prior to syphilis treatment

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Syphilis in HIV Primary, secondary, early and late latent, neurosyphilis, and of unknown duration	Treat as appropriate stage for infection		CSF examination should be done and repeat at 3, 6, 9, 12 and 24 months after therapy to detect any treatment failure (not reduced by 4-fold within 1 year). Perform full neurological examination.
Syphilis in Pregnancy (primary, secondary, early latent)	First and second trimester: Benzathine Penicillin 2.4MU IM stat Third trimester: Benzathine Penicillin 2.4MU IM weekly for 2 weeks (Day 1 & 8) OR (ALL THREE TRIMESTERS) Procaine penicillin G 600,000units IM q24h for 10 days	Penicillin Allergy: (All three trimesters) Ceftriaxone 500mg IM q24h for 10 days OR Azithromycin 500mg PO q24h for 10 days OR *Erythromycin Ethylsuccinate 800mg PO q6h for 14 days	Tetracycline & Doxycycline are contraindicated in pregnancy *If macrolide therapy, neonate require assessment and treatment at birth by Paediatrician Pregnant ladies with syphilis and history of penicillin allergy to be desensitised only in tertiary centres. Women who are treated in the second half of pregnancy are at risk of premature labour and/or fetal distress if their treatment precipitates a Jarisch-Herxheimer reaction.
Syphilis in Pregnancy (Late latent, gummatous, cardiovascular, neurosyphilis)	Treat as for non-pregnant patients (DO NOT USE DOXYCYCLINE in pregnancy)		
Gonorrhoea (<i>Neisseria gonorrhoeae</i> Infection)			
Uncomplicated (anogenital, pharyngeal gonorrhoea)	Ceftriaxone 500mg IM stat PLUS *Azithromycin 1g PO stat	Beta-lactam Allergy: Gentamicin 240mg IM stat PLUS *Azithromycin 2g PO stat	Contact tracing Sexual partners should be treated for gonorrhoea even though they are asymptomatic.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
		<u>Pregnancy & breastfeeding:</u> Ceftriaxone 500mg IM stat PLUS *Azithromycin 1g PO stat	*Azithromycin: Dual therapy to treat for coexisting <i>Chlamydia trachomatis</i> infection (35-40%), synergistic effect and reduce cephalosporin resistance.
Gonococcal Conjunctivitis	Ceftraxone 1g IM stat	Anaphylaxis to Penicillin or established allergy to Cephalosporin: Azithromycin 2g PO stat PLUS Doxycycline 100mg PO q12h for 7 days PLUS Ciprofloxacin 250mg PO q24h for 3 days	
Disseminated Gonorrhoea	Ceftriaxone 1-2g IM/IV q24h for 7 days	Cefotaxime 1g IV q8h for 7 days	When treating for the arthritis-dermatitis syndrome, the provider can switch to an oral agent guided by antimicrobial susceptibility testing 24–48 hours after substantial clinical improvement, for a total treatment course of >7 days. Length of treatment should be determined based on clinical presentation. May be switched to Ciprofloxacin 500mg PO q12h 24-48hrs after symptoms improve.
Gonococcal Epididymitis/ Epididymo-orchitis Caused by gonorrhoea and chlamydia	Ceftriaxone 500mg IM stat PLUS Azithromycin 1g PO stat		Contact tracing

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
STI related but unlikely gonorrhoea Non-STI related (Enteric organisms)	PLUS Doxycycline 100mg PO q12h for 14 days Doxycycline 100mg PO q12h for 14 days Ciprofloxacin 500mg PO q12h for 10 days		
Gonococcal Meningitis	Ceftriaxone 2g IV q12h for 10- 14 days		
Gonococcal Endocarditis	Ceftriaxone 2g IV q24h		Parenteral antimicrobial therapy for endocarditis should be administered for >4 weeks.
Chlamydia Infections (<i>Chlamydia trachomatis</i>)			
Uncomplicated • Non-specific Urethritis • Non-specific Genital Tract Infection	Doxycycline 100mg PO q12h for 7 days	Azithromycin 1g PO stat, then 500mg PO q24h for 2 days	Contact tracing
Chlamydia in Pregnancy • Non-specific Urethritis • Non-specific Genital Tract Infection	Azithromycin 1g PO stat, then 500mg PO q24h for 2 days	Amoxicillin 500mg PO q8h for 7 days OR Erythromycin Ethylsuccinate 800mg PO q6h for 7 days	Doxycycline is contraindicated in pregnancy
Non-Gonococcal Urethritis (NGU)			
First episode	Doxycycline 100mg PO q12h for 7 days	Azithromycin 500mg PO stat then 250mg q24h for 4 days	
Recurrent and persistent If treated with doxycycline first line	Azithromycin 500mg PO stat, then 250mg PO for 4 days PLUS Metronidazole 400mg PO q12h for 5 days		Most common cause: <i>Mycoplasma genitalium</i>

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
If treated with azithromycin first line	Moxifloxacin 400mg PO q12h for 10-14 days PLUS Metronidazole 400mg PO q24h for 5 days		
Herpes Simplex Virus Type-1 and 2 (HSV-1 & 2) Infections			
Herpes Genitalis			
First episode:	Acyclovir 400mg PO q8h for 7-10 days	*Valaciclovir 1g PO q12h day for 7-10 days	Acyclovir 200 mg orally five times/day is also effective but is not recommended because of the frequency of dosing. (CDC) *Requires special medicine approval Duration up to one year, then reassess
Recurrent episodes:	<u>Short Course</u> Acyclovir 800mg q8h PO for 2 days OR <u>5 days regime</u> Acyclovir 400mg q8h PO for 5 days	*Valaciclovir 500mg PO q12h for 3-5 days	
Suppressive therapy: (may be indicated if > 6 recurrences per year)	Acyclovir 400mg PO q12h	*Valaciclovir 500mg PO q24h	
Herpes Genitalis in HIV			
Primary:	**Acyclovir 400-800mg PO q8-12h for 5-10 days	*Valaciclovir 1gm PO q12h for 5- 10 days	*Requires special medicine approval
Severe:	Acyclovir 5-10mg/kg IV q8h for 2-7 days and then followed by Acyclovir PO (minimum 10 days)		**Treatment courses might need to be extended for lesion resolution
Recurrent:	Acyclovir 400mg PO q8h for 5–10 days	*Valaciclovir 1g PO q12h for 5–10 days	Physical supportive measures: saline bathing, analgesic local anaesthetics and psychological support.
Suppressive:	***Acyclovir 400mg-800mg PO q8-12h	*/***Valaciclovir 500mg PO q12h	Oral antiviral drugs indicated within 5 days of the start of the episode and while new lesions are still forming.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			Topical antivirals are less effective than oral agents and not recommended, due to the association with acyclovir resistant strain. Addition of topical antivirals to oral treatment is of no benefit ***Duration: All for up to 1 year, then reassess If breakthrough recurrences occur, dosage should be increased (refer recurrent dose)
Herpes Genitalis in pregnancy First episode Suppressive therapy for recurrent infections	<u>First or second trimester acquisitions</u> Acyclovir 400mg PO q8h for 5 days <u>Third trimester (from 28 weeks)</u> Acyclovir 400mg PO q8h for 5 days Acyclovir 400mg PO q8h (Treatment recommended starting at 36 weeks of gestation until delivery)	 *Valaciclovir 500m PO q12h day for 5 days *Valaciclovir 500mg PO q12h for 5 days	Do not delay treatment whilst awaiting results (HSV PCR recommended) <u>Third trimester acquisition:</u> -No additional monitoring of the pregnancy is required -Continue daily suppressive Acyclovir 400 mg PO q8h until delivery *Requires special medicine approval
Other Sexually Transmitted Infections			
Chancroid <i>Haemophilus ducreyi</i>	Azithromycin 1g PO stat OR Ceftriaxone 250mg IM stat OR		*preferred in HIV positive patients, but not preferred in pregnant/lactating patients

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	*Ciprofloxacin 500mg q12h for 3 days		Contact tracing Sexual contacts within 10 days before onset of patient's symptoms should be examined and treated even in the absence of symptoms Avoid unprotected sexual intercourse until they and their partner(s) have completed treatment and follow-up. Patients should be re-examined 3 to 7 days after initiation of therapy. Successful therapy: ulcers improved symptomatically within 3 days and substantial re-epithelization within 7 days after onset of therapy
Lymphogranuloma Venereum <i>Chlamydia trachomatis</i> <i>Serovar L1, 2, 3</i>	Doxycycline 100mg PO q12h for 21 days	Azithromycin 1g PO weekly for 3 weeks	Contact tracing Persons who have had sexual contact with a patient who has LGV within the 60 days before onset of the patient's symptoms should be evaluated, examined, and tested for chlamydial infection, depending on anatomic site of exposure. Asymptomatic partners should be presumptively treated with a chlamydia regimen (doxycycline 100 mg orally 2 times/day for 7 days)

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			All persons who have been treated for LGV should be retested for chlamydia approximately 3 months after treatment. If retesting at 3 months is not possible, providers should retest at the patient's next visit for medical care within the 12-month period after initial treatment.
Granuloma Inguinale <i>Klebsiella granulomatis</i>	Azithromycin 1g PO weekly or 500mg PO q24h PLUS/MINUS Gentamicin 1mg/kg IV/IM q8h (in patients whose lesion do not respond in the first few days to other agents)	Doxycycline 100mg PO q12h OR Trimethoprim/sulfamethoxazole 80/400mg 2 tabs PO q12h OR Ciprofloxacin 750 mg PO q12h PLUS/MINUS Gentamicin 1mg/kg IV/IM q8h (in patients whose lesion do not respond in the first few days to other agents)	Duration of treatment should be until lesions have healed. Healing times vary greatly between patients. A minimum of 3 weeks treatment is generally recommended Contact tracing In the absence of any reliable screening test and the long incubation period, all sexual contacts in the last 6 months should be examined for possible lesions by clinical examination.
Trichomoniasis <i>Trichomonas vaginalis</i> Treatment failure (second regimen):	Metronidazole 2g PO stat OR 400mg PO q12h for 5 days Metronidazole 400mg PO q12h for 7 days		Sexual contact(s) should be treated simultaneously and patients should be advised to abstain for at least one week until they and their partner(s) have completed treatment and follow-up. Any partners within the 4 weeks prior to presentation should be screened

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			for the full range of STIs and treated for TV. TOC is only recommended if the patient remains symptomatic following treatment, or if symptoms recur. Higher-dose of metronidazole is required if failing the second regimen.
Bacterial vaginosis <i>Gardnerella vaginalis</i> , Anaerobes	Metronidazole 400mg PO q12h for 5-7 days OR 2g PO stat	Clindamycin 300mg PO q12h for 7 days	Not an STI but frequently detected during STI screening
Reference: 1. MMWR Sexually Transmitted Infections Guidelines 2021, CDC 2. National Antimicrobial Guideline 2019. 3. Prevention, Diagnosis & Management Of Infective Endocarditis 2017			

SKIN AND SOFT TISSUE INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
PURULENT SKIN AND SOFT TISSUE INFECTION			
Localized Impetigo (<i>S. aureus</i> , <i>S. pyogenes</i>)	*Topical 2% fusidic acid q8-12h for 5 days (Outpatient use only) OR Cloxacillin 500-1000mg PO q6h for 5-7 days OR Cephalexin 250-500mg PO q6h for 5-7 days		*Only can be used by Dermatologist
Generalised Impetigo/Ecthyma	Cephalexin 250-500mg PO q6h <u>Penicillin allergy:</u> Erythromycin Ethylsuccinate 800mg PO q12h	Amoxicillin/clavulanate 625mg PO q8h <u>Other alternative/ in case of CA-MRSA:</u> Clindamycin 600mg PO q8h OR Trimethoprim/sulfamethoxazole 160/800mg PO q12h	Duration : 5-7 days
Ecthyma gangrenosum <i>Pseudomonas sp.</i>	Ciprofloxacin 500mg PO q12h OR *Piperacillin-Tazobactam 4.5g IV q6-8h	Ceftazidime 2gm IV q8h OR Cefepime 2gm IV q8h	Consider adding aminoglycoside in selected cases such as in immunocompromised/neutropenic and septic shock patients. Use synergistic combination therapy with aminoglycosides until susceptibilities are known. Antibiotics need to be tailored according to culture results.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			*Piperacillin/tazobactam: If given as q8h, to be given as extended infusion (over 3-4 hours).
NON-PURULENT SKIN & SOFT TISSUE INFECTION			
Furuncles	Cloxacillin 500mg PO q6h	Amoxicillin/clavulanate 625mg PO q8h	For 5-7 days
Carbuncles <i>S. aureus</i>	Cloxacillin 1-2gm IV q6h	Amoxicillin/clavulanate 1.2gm IV q8h OR Cefazolin 1gm IV q8h	Surgical drainage is the mainstay of treatment. Duration : 7-10 days
Erysipelas <i>Strep. pyogenes</i> MRSA	Phenoxymethylpenicillin 500mg PO q6h OR Amoxicillin 500mg PO q8h If severe: Benzylpenicillin 2-4MU IV q4-6h Vancomycin 25mg/kg loading dose then 15-20mg/kg q8-12h; not to exceed 2gm/dose	Cephalexin 500mg PO q6h If severe: Cefazolin 1gm IV q8h OR Cefuroxime 750mg IV q8h	Duration : 7-10 days
Diabetic Foot Infections	Refer to Bone & Joint Infections Section		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Gas Gangrene/ Myonecrosis/ Necrotizing Fasciitis	Refer to Bone & Joint Infections Section		
Yaws <i>Treponema pertenue</i>	Benzathine Penicillin 1.2 MU IM single dose	Doxycycline 100mg PO q12h for 15 days OR Azithromycin 30mg/kg (max 2g) single dose Penicillin Allergy: Erythromycin Ethylsuccinate 800mg PO q12h for 15 days	
CELLULITIS			
Mild <i>Staph. aureus</i> <i>Strep. Pyogenes</i>	Cephalexin 500mg PO q6h	Cefuroxime 250-500mg PO q12h OR Amoxicillin/clavulanate 625mg PO q8h	Duration: 5-10 days according to clinical response Change to oral once the condition improved Gram negative coverage may be necessary in the following circumstances: 1. Potential relation of the cellulitis to a decubitus ulcer 2. Crepitant cellulitis 3. Prominent skin necrosis/ gangrene 4. Location (perioral/perirectal cellulitis)
Moderate <i>Staph. aureus</i> <i>Strep. Pyogenes</i>	Cloxacillin 1-2gm IV q6h	Cefazolin 1-2gm IV q8h	
Severe <i>Staph. aureus</i> <i>Strep. Pyogenes</i>	Ampicillin/sulbactam 3gm IV q6-8h PLUS/MINUS Clindamycin 600mg IV q6h (Deescalate once cultures are available/ Necrotizing fasciitis ruled out)	*Piperacillin/tazobactam 4.5gm IV q6-8h PLUS/MINUS Clindamycin 600mg IV q6h (Deescalate once cultures are available/ Necrotizing fasciitis ruled out)	
**Consider alternative organisms in the following circumstances:			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Dog/cat bite <i>Pasteurella multocida</i> <i>Capnocytophaga canimorsus</i>	Amoxicillin/clavulanate 625mg PO q8h		5. Clinical condition: (Septicaemic shock/Suspecting necrotizing fasciitis) 6. Immunocompromised patients 7. Specific exposures** *Piperacillin/tazobactam: If given as q8h, to be given as extended infusion (over 3-4 hours) ***Consider adding 3rd-generation Cephalosporin in severe infections
Cat scratch disease <i>Bartonella henselae</i>	Azithromycin 500mg PO on Day 1, then 250mg PO q24h for 4 days		
Human bite <i>Eikenella corrodens</i> <i>Anaerobes</i> <i>Staph. aureus</i>	Amoxicillin/clavulanate 625mg PO q8h		
Salt water exposure <i>Vibrio sp.</i>	Doxycycline 200mg stat then 100mg PO q12h PLUS/MINUS ***Ceftriaxone 2gm IV q24h		
Fresh or brackish water exposure <i>Aeromonas sp.</i> <i>Plesiomonas</i>	Ciprofloxacin 400mg IV q12h OR Ciprofloxacin 750mg PO q12h		
Neutropenic patients <i>Pseudomonas aeruginosa</i> other Gram negatives	*Piperacillin/tazobactam 4.5gm IV q6-8h	Ceftazidime 2gm IV q8h OR Cefepime 2gm IV q8h	

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
MRSA	Vancomycin 25mg/kg loading dose then 15-20mg/kg IV q8-12h In severe infections: To load with Vancomycin 25-30mg/kg IV, followed by 15-20mg/kg (actual body weight) IV q8-12h; not exceeding 2g/dose	Linezolid 600mg IV/PO q12h	****Consider CA-MRSA if : 1. Outbreaks of known CA-MRSA 2. If non-resolving cellulitis
**** If CA-MRSA suspected	Clindamycin 300-450mg IV/PO q8h OR Doxycycline 100mg PO q12h OR Trimethoprim/sulfamethoxazole 160/800mg PO q12h		
PERIPHERAL PHLEBITIS/THROMBOPHLEBITIS			
<i>Staphylococcus aureus</i> <i>Coagulase negative Staphylococcus</i> <i>Gram negative rods</i>	Early stage phlebitis: Remove the intravenous cannula Medium and advanced stage phlebitis or thrombophlebitis: Remove the intravenous cannula and take blood culture Can consider empirical treatment if persistent fever: Cephalexin 500mg PO q6h OR Cloxacillin 1-2gm IV q6h		Peripheral intravenous catheters with associated pain, induration, erythema, or exudate should be removed.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
BED SORE/PRESSURE SORE/DECUBITUS ULCER			
	Local treatment is preferred. If there is surrounding cellulitis/signs of bacteremia/fasciitis/surrounding intramuscular abscess/OM changes: Ampicillin/sulbactam 3gm IV q6-8h		
MYCOBACTERIAL INFECTIONS			
Hansen’s Disease (Leprosy) <i>Mycobacterium Leprae</i>	Paucibacillary Rifampicin 600mg PO monthly PLUS Dapsone 100mg PO q24h Duration: 6 months (Completion of 6 doses within 9 months) Surveillance: 5 years Multibacillary Rifampicin 600mg PO monthly PLUS Clofazimine 300mg PO monthly and 50-100mg PO q24h	*Bacterial resistance or hypersensitivity to first line Can be substituted with one of the following: Minocycline 100mg PO q24h OR Ofloxacin 400mg PO q24h OR Clarithromycin 500mg PO q24h OR Ethionamide 250mg PO q24h	*Second line can only be initiated by a dermatologist

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	PLUS Dapsone 100mg PO q24h Duration: 1 year (if initial BI<4) or 2 years (if BI≥4) Completion of 12 doses within 18 months (BI<4) Completion of 18 doses within 36 months (BI≥4) Surveillance: 15 years		
Hansen’s Disease (Leprosy) in HIV	Same as non HIV patients		
NON-TUBERCULOUS MYCOBACTERIAL INFECTIONS			
<i>Mycobacterium marinum</i>	Clarithromycin 500mg PO q12h PLUS Minocycline/Doxycycline 100mg PO q12h At least 2 months of treatment until clearance	Rifampicin 600mg PO q24h PLUS Ethambutol 15mg/kg PO q24h for 4-6 months, and continue for at least 1 month after lesions have been cleared OR Doxycycline 100mg q12h for 1-2 months after lesion clearance (3-4 months)	Often resistant to isoniazid
<i>Mycobacterium kansasii</i>	Isoniazid 300mg PO q24h PLUS Rifampicin 600mg PO q24h PLUS Ethambutol 15mg/kg PO q24h		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	Duration: 18 months		
<i>Mycobacterium ulcerans</i> (Buruli ulcer)	Rifampicin 10mg/kg PO q24h PLUS Streptomycin 15mg/kg IM q24h Duration: 8 weeks	Rifampicin 10mg/kg PO q24h PLUS Streptomycin 15mg/kg IM q24h Duration: 4 weeks Followed by Rifampicin 10mg/kg PO q24h PLUS Clarithromycin 7.5mg/kg PO q12h Duration: 4 weeks (Total duration of 8 weeks)	Wide surgical excision and debridement are important.
<i>Mycobacterium fortuitum/ chelonae</i>	Combination therapy (2 of the following): Clarithromycin 500mg PO q12h OR Doxycycline/Minocycline 100mg PO q12h OR Ciprofloxacin 500-750mg PO q12h PLUS/MINUS *Amikacin 15mg/kg IV q24h		*Amikacin: Started for severe infection until clinical improvement (together with 2 oral agents), then continue with just 2 oral agents.
FUNGAL INFECTIONS			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Tinea capitis <i>Trichophyton</i> <i>Microsporum</i>	Griseofulvin 500mg q12h for 6 to 12 weeks or longer till fungal cultures are negative OR Terbinafine 250mg PO q24h PLUS 2.5% Selenium sulphide shampoo OR 2% ketoconazole shampoo, 2 – 3 times per week for 2 weeks	Itraconazole 200mg PO q24h OR Fluconazole 6mg/kg PO q24h Duration is based on mycological agent: - <i>Trichophyton</i> spp : 2-4 weeks - <i>Microsporum</i> spp : 8-12 weeks	Other recommendations: 1)Kerion: Griseofulvin should be considered as the first line unless <i>Trichophyton</i> has been cultured as the pathogen. Duration of treatment may be longer. 2) Contacts of patient may be treated with 2% ketoconazole shampoo 2 – 3 times per week for 2 weeks. 3) Surgical excision is to be avoided 4)Topical therapy alone is not recommended for the management of tinea capitis. 5)Consider adding oral prednisolone in selected cases.
Tinea barbae	Same as treatment of Tinea capitis		
Tinea corporis /Tinea cruris/Tinea faciei <i>Trichophyton</i> <i>Microsporum</i> , <i>Epidermophyton</i>	Mild Infections: Topical imidazoles/allylamines cream/lotion: Clotrimazole 1% OR Miconazole 2% OR Ketoconazole OR Terbinafine Duration: till clinical clearance with additional 2 weeks		Recommendations: 1. In patients with renal or hepatic impairment, caution should be exercised while prescribing systemic antifungals. 2. Terbinafine clearance significantly reduced in patients with renal impairment. Other systemic antifungals are preferred in these patients. 3. Topical Nystatin should not be used in dermatophytosis as they are not effective against dermatophytes

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	Extensive infections: Terbinafine 250mg PO q24h for 2 weeks OR Itraconazole 200mg PO q24h for 2 weeks OR Griseofulvin 500mg PO q12h or q24h for 4-6 weeks	Fluconazole 200-300mg per week PO for 3-4 weeks	
Tinea manuum/ Tinea pedis <i>Trichophyton, Microsporum, Epidermophyton</i>	First line: Topical antifungals as mentioned in tinea corporis for 4-8 weeks Resistant cases: Terbinafine 250mg PO q24h for 2-4 weeks OR Itraconazole 200mg PO q24h for 2-4 weeks OR Griseofulvin 500mg PO q12h for 6-12 weeks	Fluconazole 200mg/week PO for 4 weeks	Recommendations: 1. Topical keratolytic agents can be used in conjunction with antifungals for hyperkeratotic type of tinea pedis/manuum. 2. KMnO ₄ in 1:10,000 dilution wet dressings, applied for 20 min 2-3 times/day, may be helpful if vesiculation or maceration is present. 3. Systemic antifungals can be prescribed as first line treatment in severe moccasin-type tinea pedis or severe recurrent tinea with blisters.
Tinea unguium <i>Trichophyton, Microsporum, Epidermophyton</i>	Amorolfine 5% Nail Lacquer weekly application Duration: For 6 months (fingernails) For 12 months (toenails) OR	Griseofulvin 500mg PO q12h Duration: For 6 months (fingernails) For 12 months (toenails) OR Fluconazole 200mg PO once weekly Duration:	Amorolfine 5% Nail Lacquer is not indicated for children less than 12 years old. Patients with contraindications to systemic agents may consider topical antifungal agents.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	Pulse Itraconazole 200mg PO q12h for 1 week per month Duration: For 2 months (fingernails) For 3 months (toenails) OR Terbinafine 250mg PO q24h Duration: For 6 weeks (fingernails) For 12 weeks (toenails)	For ≥ 3 months (fingernails) For 6-12 months (toenails)	Diagnosis of onychomycosis should be confirmed with a KOH preparation, culture, or PAS Stain. Empirical treatment is not recommended.
Tinea versicolor <i>Malassezia Furfur</i> <i>Pityrosporum Orbiculare</i>	First line: Topical treatment only Selenium Sulphide 2% Shampoo Apply to affected areas 10 minutes before bathing OR dilute to 1:1 with water, apply and leave overnight (treat for 1-2 weeks) OR 2% Ketoconazole shampoo apply to affected areas 10 minutes before bathing For face: Topical Imidazole for 4-6 weeks e.g: Miconazole 2% cream OR Clotrimazole 1% cream For recurrent or resistant cases: Itraconazole 200mg PO q24h for 1 week	Sulfur PLUS salicylic solution	Recommendations: Ketoconazole shampoo or Selenium sulphide shampoo to be used once every two to four weeks for approximately six months in order to try and prevent recurrence.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	OR Fluconazole 200-300mg PO weekly dose for 2 to 4 weeks		
Candidiasis <i>Candida albicans</i>	Mild cutaneous candidiasis: Topical Imidazole q12h till clear e.g, Miconazole 2% cream OR Clotrimazole 1% cream		Treatment of sexual partners is advisable in case of recurrent infection. *Itraconazole: Absorption depends on gut acidity. Take capsule with food and acidic beverage (e.g.: Cola drinks). Avoid PPIs and H2 blockers.
	Extensive cutaneous candidiasis: *Itraconazole 200mg PO q24h for 1 week	In severe and immunocompromised patients: Fluconazole 100mg PO q24h for 1 week	
Subcutaneous Fungal Infections Lymphocutaneous and Cutaneous Sporotrichosis	*Itraconazole 200mg PO 12h until all lesions have resolved (usually for a total of 3-6 months)	For patients not able to tolerate Itraconazole: Terbinafine 250mg PO q12h OR Fluconazole 400-800mg q24h	In some immunocompromised condition such as AIDS, longer treatment may be necessary. Refer to Opportunistic Infections in HIV Patients *Itraconazole: Absorption depends on gut acidity. Take capsule with food and acidic beverage (e.g.: Cola drinks). Avoid PPIs and H2 blockers.
Systemic sporotrichosis (pulmonary osteoarticular, meningeal, or disseminated sporotrichosis)	Amphotericin B deoxycholate 0.7-1mg/kg q24h for 2 weeks followed by *Itraconazole 200mg PO q12-24h for 12 months		
Sporotrichosis In Pregnancy*	Terbinafine 250mg PO q24h	Amphotericin B deoxycholate 0.7-1mg/kg q24h	*Avoid azole in pregnancy.
Cutaneous fungal infection in immunocompromised patients	Refer to treatment of disseminated fungal infection in immunocompromised/HIV patients		Skin biopsy for HPE and culture are advised before commencing treatment.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
<i>Aspergillus</i> sp, <i>Scedosporium apiospermum</i>, and <i>Fusarium</i> sp. infection	Voriconazole 6mg/kg IV q12h for 2 doses, followed by 4mg/kg IV q12h	Amphotericin B deoxycholate 0.7-1mg/kg q24h OR *Amphotericin B lipid formulation 3-5mg/kg q24h	*Only by ID prescriber
Cryptococcal infections 1.Mild 2.Life threatening	Fluconazole 100mg-400mg PO q24h Refer to Treatment of disseminated fungal infection in Immunocompromised/HIV patients		
Penicilliosis and life threatening acute severe disseminated Histoplasmosis	Refer to Treatment of disseminated fungal infection in immunocompromised/HIV patients		
VIRAL INFECTIONS			
Herpes Simplex Infections	i) Mild infection: Acyclovir 400mg PO q8h for 5 days ii) Severe life threatening: Acyclovir 5-10mg/kg IV q8h for 5 days or until able to take orally, then change to oral iii) Genitalia: Refer to Sexually Transmitted Infections section		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Chickenpox (Varicella Zoster)	i) Immunocompetent Acyclovir 800mg PO 5 times daily for 7 days ii) Immunocompromised Acyclovir 10mg/kg IV q8h for 7 days (change to oral once there is an improvement)		Advisable to start treatment early within 48 hours.
Herpes Zoster	Refer to varicella zoster treatment above		Topical antiviral treatment is not recommended for Herpes Zoster. Systemic antiviral treatment is recommended for all immunocompromised patients or for immunocompetent patients with following criteria: (1) >50 years of age (2) have moderate or severe pain (3) have moderate or severe rash (4) have non-truncal involvement Advisable to start treatment early within 48-72 hours
PARASITIC INFESTATION			
Scabies <i>Sarcoptes scabiei</i>	Benzyl Benzoate emulsion 25% (EBB) apply from neck down and leave for 24 hours for 2-3 days OR		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	Permethrin 5% lotion/cream apply and leave for 8 hours Repeat application after 1 week In pregnancy/ Immunocompromised: Permethrin 5% lotion/cream apply and leave for 8 hours Repeat application after 1 week		
Head Lice <i>Pediculus humanus capitis</i>	Permethrin 1% lotion apply to scalp for 10 min and wash off		
Body Lice/pubic Lice <i>Pediculus humanus</i>	Permethrin 1% cream apply to affected area for 10 min and wash off		
References: 1. Stevens et al. Clinical Infectious Diseases 2014;59(2) 2. Sarkar S et al. Indian Dermatol Online J 2016; 7:36. 3. Reich HL et.al. J Am Acad Dermatol. 2004;50 4. Morton N Swartz. N Engl J Med 2004; 350:904-12. 5. CREST guidelines. 2005 6. Begier EM et al. Clin Infect Dis 2004; 39:1446. 7. Malaysian Clinical practice Guideline on Management of leprosy 2014 8. Primary Care Dermatology Society UK 2013 9. L.C. Fuller,1 R.C. Barton,2 M.F. Mohd Mustapa,3 L.E. Proudfoot,4 S.P. Punjabi5 and E.M. Higgins, British Association of Dermatologists' guidelines for the management of tinea capitis 2014 10. RxFiles Newsletter : Antifungal newsletter (April 2010) Canadian : Bugs and Drugs 11. Alok Kumar Sahoo et al, indian journal of dermatology 2016, Management of tinea corporis, tinea cruris, and tinea pedis: A comprehensive review 12. Craig G Burkhart et al. Tinea Versicolor Treatment & Management.medscape. updated Dec 2013 13. Clinical Practice Guidelines for the Management of Sporotrichosis: 2007 Update by the Infectious Diseases Society of America 14. Centers for Disease Control and Prevention (CDC) 2010			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
15. IDSA Guidelines for Intravascular Catheter-Related Infection • CID 2009:49 16. ESPID Reports and Review : The Pediatric Infectious Disease Journal 2014 17. Rook Textbook Dermatology 4th edition (www.dermnetnz.org)			

SURGICAL INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
A. GENERAL SURGERY			
Acute Pancreatitis			
Mild to moderate	No antibiotics		Antibiotics should be given for extra-pancreatic infections, such as cholangitis, catheter-acquired infections, bacteremia, urinary tract infections and pneumonia.
Severe Antibiotics are mainly indicated for infected pancreatic necrosis. Possible causative organisms: Enterobacterales Enterococci <i>Staphylococcus aureus</i> <i>Streptococcus</i> <i>Staphylococcus epidermidis</i> Anaerobes <i>Candida</i> sp. (rarely)	*Piperacillin/tazobactam 4.5g IV q6-8h	Cefoperazone 1-2g IV q12h PLUS Metronidazole 500mg IV q8h	Modify antibiotics once culture and sensitivity are available. Reserve carbapenem for infections caused by resistant pathogens. *Piperacillin/tazobactam: If given as q8h, to be given as extended infusion (over 3-4 hours).
Diverticulitis Antibiotics considered for patients with the following: Fever, elevated WBC, patients who have failed to respond to conservative management			
Diverticulitis	Amoxicillin/clavulanate 625mg PO q8h for 5 days	Non-severe penicillin allergy: Cefuroxime 1.5g IV q8h	

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
(Not undergoing a source control procedure)	OR Ampicillin/sulbactam 3g IV q6h	PLUS Metronidazole 500mg IV q8h	
Diverticulitis (Severe infection/life threatening infection)	*Piperacillin/tazobactam 4.5g IV q6-8h for 7 days	Severe penicillin allergy: Ciprofloxacin 400mg IV q12h PLUS Metronidazole 500mg IV q8h	*Piperacillin/tazobactam: If given as q8h, to be given as extended infusion (over 3-4 hours).
Breast abscess/Mastitis Common organism: <i>Staphylococcus aureus</i>	Cloxacillin 1-2g IV q6h OR Cefazolin 1-2g IV q8h	Amoxicillin/clavulanate 625mg PO q8h OR Ampicillin/sulbactam 750mg PO q12h Penicillin allergy: Clindamycin 600mg IV/PO q8h	Drainage may be required for abscess. For lactating mastitis: Consider sending breast milk for C&S if not responding after 48 hours of initial antibiotic therapy or recurring mastitis. Duration: 10 to 14 days but shorter courses (5 to 7 days) can be used if the response to therapy is rapid and complete.
Hernia repair with mesh	Refer to Chemoprophylaxis-Surgical		
Appendicitis Common organisms: Enterobacterales Enterococci Bacteroides	Cefuroxime 1.5g IV q8h PLUS Metronidazole 500mg IV q8h	Ampicillin/sulbactam 1.5g IV q6-8h OR Amoxicillin/clavulanate 1.2g IV q8h	Acute appendicitis without evidence of perforation, abscess, or local peritonitis; treatment should be discontinued within 24 hours. For patients with various forms of appendicitis not undergoing a source

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			control procedure, change to early oral therapy. Duration: 4-7 days
Perforated Appendix/Appendicular Mass	Cefuroxime 1.5g IV q8h OR Cefoperazone 1-2g IV q12h PLUS Metronidazole 500mg IV q8h	Ampicillin/sulbactam 1.5-3g IV q6-8h OR Amoxicillin/clavulanate 1.2g IV q8h	Duration: 4-7 days
Perforated Viscus Peritonitis	Cefuroxime 1.5g IV q8h OR Cefoperazone 1-2g IV q12h PLUS Metronidazole 500mg IV q8h	Ampicillin/sulbactam 1.5-3g IV q6-8h OR Amoxicillin/clavulanate 1.2g IV q8h	Duration: 4-7 days
Abdominal trauma Stab wound Suspected bowel or solid organ injury Common organisms: Gram-negative enteric aerobes and anaerobes	Amoxicillin/clavulanate 1.2g IV q8h	Cefuroxime 1.5g IV q8h PLUS Metronidazole 500mg IV q8h	Duration: 4-7 days (if adequate source control, no delay in surgical intervention and patient has rapid recovery) *Piperacillin/tazobactam: if given as q8h, to be given as extended infusion (over 3-4 hours).
	Severe/Infected wound: Cefazolin 2g IV q8h PLUS Metronidazole 500mg IV q8h OR *Piperacillin/tazobactam 4.5g IV q6-8h	Severe/Infected wound: Ciprofloxacin 400mg IV q12h PLUS Clindamycin 450-600mg IV q8h	

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Anal/Rectal abscess	Amoxicillin/clavulanate 1.2g IV or 625mg PO q8h		Drainage is required. Duration: 4-7 days (if adequate source control, no delay in surgical intervention and patient has rapid clinical recovery) Routine antibiotics are not recommended in otherwise healthy patients.
Burn wound sepsis Likely organisms: <i>Streptococcus pyogenes</i> <i>Staphylococcus aureus</i> <i>Enterobacter</i> spp. <i>Staphylococcus epidermidis</i> <i>Enterococcus faecalis</i> <i>Pseudomonas aeruginosa</i>	*Piperacillin/tazobactam 4.5g IV q6-8h	Cefepime 1-2g IV q8h	<i>S. aureus</i> tends to remain localised to burn wounds, if toxic, consider toxic shock syndrome. <i>Candida</i> sp. colonise seldom invades. Once C&S results are available, antibiotic therapy should be adjusted accordingly. *Piperacillin/tazobactam: if given as q8h, to be given as extended infusion (over 3-4 hours).
VASCULAR			
Mycotic aneurysm (Initial Treatment) Vascular prosthesis infection Common organisms:	Ceftriaxone 2g IV q24h *Step down therapy: Ciprofloxacin 250mg PO q12h	*Piperacillin/tazobactam 4.5g IV q6-8h	Duration: At least six weeks (IV then oral based on clinical response and cultures)

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
<i>Streptococcus pneumoniae</i> <i>Staphylococcus aureus</i> (30%) <i>Salmonella</i> sp. (50%)	OR Amoxicillin/clavulanate 625mg PO q8h		*Piperacillin/tazobactam: if given as q8h, to be given as extended infusion (over 3-4 hours). Consider adding Vancomycin if suspecting MRSA/CoNS or vascular prosthesis infection *CRP monitoring upon follow-up
Ischaemic limb ulcers with infection	Ampicillin/sulbactam 1.5-3g IV q6-8h for 7 days (to continue until C&S available)	Amoxicillin/clavulanate 1.2g IV q8h for 7 days (to continue until C&S available)	Duration depends on the extent of the infection (longer if bone is involved)
BITES (penetrating injuries)			
Animal bite Common organisms: <i>Staphylococcus aureus</i> <i>Streptococcus</i> Gram-negative Bacilli Anaerobes <i>Pasturella</i> (50% dog bites and 75% cat bites) <i>Eikenella corrodens</i> <i>Pseudomonas</i> sp.	Amoxicillin/clavulanate 625mg PO q8h	Doxycycline 100mg PO q12h PLUS Clindamycin 300mg PO q6h If severe/life threatening: Ampicillin/sulbactam 1.5-3g IV q6-8h OR Piperacillin/tazobactam 4.5g IV q8h	Prophylactic duration: 3-5 days -Associated crush injury -In the hands or proximity to a joint -Associated edema If wound is infected: 10 days or longer is recommended *Piperacillin/tazobactam: if given as q8h, to be given as extended infusion (over 3-4 hours).
	If severe/life threatening: Ampicillin/sulbactam 1.5-3g IV q6-8h	If severe/life threatening: *Piperacillin/tazobactam 4.5g IV q6-8h	
Human bite Common organisms: <i>Staphylococcus aureus</i>	Amoxicillin/clavulanate 625mg PO q8h	Penicillin Allergy: Clindamycin 300mg PO q6h PLUS Ciprofloxacin 500-750mg PO q12h	

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Anaerobes <i>Eikenella corrodens</i> <i>Streptococcus</i> (esp <i>viridans</i>)		OR Trimethoprim/sulfamethoxazole 160/800mg PO q12h	
References: 1. Mazuski JE, Tessier JM, May AK, Sawyer RG, Nadler EP, Rosengart MR, et al. The Surgical Infection Society Revised Guidelines on the Management of Intra-Abdominal Infection. <i>Surgical Infections</i>. 2017;8(1). 2. Bradley E. A Clinically Based Classification System for Acute Pancreatitis. Summary of the International Symposium on Acute Pancreatitis, Atlanta, Ga, September 11 Through 13, 1992. <i>Arch Surg</i>. 1993;128(5):586-90. 3. Neil Stollman WS, Ikuo Hirano et al. American Gastroenterological Association Institute Guideline on the Management of Acute Diverticulitis. <i>Gastroenterology</i>. 2015;149(7):1944–194. 4. Banks PA, Bollen TL, Dervenis C, Gooszen HG, Johnson CD, Sarr MG, et al. Classification of acute pancreatitis--2012: revision of the Atlanta classification and definitions by international consensus. <i>Gut</i>. 2013;62(1):102-11. 5. Perez A WE, Brooks DC, et al. Is severity of necrotizing pancreatitis increased in extended necrosis and infected necrosis? <i>Pancreas</i>. 2002;25(3):229-33. 6. Australian Clinical Practice Guidelines. Therapeutic guidelines antibiotic version 15. Australian Government, National Health and Medical Research Council 2014. 7. Joseph A, Weston V, Catt L. Antimicrobial Prescribing Guidelines for Primary Care 2017. National Health Services (NHS) Nottinghamshire Area Prescribing Committee. 2017. 8. Dennis L. Stevens, Alan L. Bisno, Henry F. Chambers, E. Patchen Dellinger, Ellie J. C. Goldstein, Gorbach SL, et al. Practice Guidelines for the Diagnosis and Management of Skin and Soft Tissue Infections: 2014 Update by the Infectious Diseases Society of America. <i>Clinical Infectious Diseases</i> 2014;59(2):e10–52. 2014;59(2):e10-52. 9. University Malaya Medical Centre Antibiotic Guidelines Edition 2017. 2017. 10. Wilson B. Necrotizing fasciitis. <i>Am Surg</i>. 1952;18(4):416-31. 11. Burpee JF EP. Fournier's gangrene. <i>J Urol</i>. 1972;107(5):812-4. 12. Currie B. Melioidosis: The 2014 Revised RDH Guideline. <i>The Northern Territory Disease Control Bulletin</i> 2014;21(2) 13. IDSA Practice Guidelines, April 2014.			
B. BONE AND JOINT INFECTIONS			
OSTEOMYELITIS			
Acute Osteomyelitis	Empirical coverage:	Penicillin allergy: Cefazolin 2g IV q6-8h	Duration: Initial IV therapy for 2-4 weeks followed by oral therapy.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Common organisms: <i>Staphylococcus aureus</i> (80%) Group A <i>Streptococcus pyogenes</i> Rarely Gram-negative bacilli	Cloxacillin 2g IV q6h To tailor antibiotics according to definitive cultures		Minimum 6 weeks. Modify according to clinical response. A shorter duration of antibiotic can be considered if the osteomyelitis is fully resected (e.g. amputation with a clear margin): - No surrounding soft tissue infection: 5 days. - Evidence of soft tissue infection: 10-14 days.
Chronic Osteomyelitis Definition: - Relapsing infection despite adequate duration of appropriate antibiotic - Chronic pain/swelling/bone tenderness associated with tissue necrosis, increased drainage or persistent sinus tracts. - Presence of bone destruction and presence of sequestra on imaging Common organism: <i>Staphylococcus aureus</i>	Empirical treatment before taking adequate cultures is not recommended Choice of antibiotic depends on C&S result from tissue/bone as <u>swab</u> culture <u>NOT reliable</u>. Thorough surgical debridement required (Removal of deadbone/orthopaedic hardware)		Duration: 6 weeks but usually > 3 months Treat until inflammatory parameters are normal

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Vertebral Osteomyelitis Epidural Abscess Common organisms: <i>Staphylococcus aureus</i> (main) <i>Brucella</i> <i>Salmonella</i> Gram-negative Bacilli	Empirical therapy should be withheld unless the patient is septic or in patients with neurologic compromise. Cloxacillin 2g IV q4h To tailor antibiotics according to definitive cultures.	Cefazolin 2g IV q6-8h	Empiric Gram-negative should be covered if patient had: -Recent spinal hardware inserted or surgery -Intra-abdominal infections -Coexisting or synchronous genitourinary infection -HIV infection
	Duration: Minimum 6 weeks. Minimum 8 weeks if undrained paravertebral abscess(es) and/or infection due to drug-resistant organisms Up to 12 weeks if extensive bone destruction.		Surgical therapy is necessary in: -Spinal cord compression/instability -Persistence of epidural abscess despite adequate antibiotic -Considering TB spine/MDR organisms
SEPTIC ARTHRITIS			
Acute monoarticular In person who do not have any risk factors for STD (<i>Staphylococcus/Streptococcus</i>)	Cloxacillin 2g IV q4-6h Duration: IV therapy 2-4 weeks Oral therapy to complete total 4-6 weeks	Penicillin allergy: Cefazolin 2g IV q6-8h OR Clindamycin 600mg IV q6h, followed by oral therapy (same dose) OR *Vancomycin 15-20mg/kg (actual body weight) IV q8-12h; not to exceed 2g/dose Duration: IV therapy 2-4 weeks Oral therapy to complete total 4-6 weeks	Drainage, debridement and washout the infected joint is important to limit further damage. A shorter duration of therapy is possible in immunocompromised patients who have had adequate surgical drainage. *Vancomycin: If suspected/confirmed MRSA, consider loading dose 25-30mg/kg for critically ill/ septic patients to achieve faster steady state.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			Empirical therapy wherever possible should be directed by the result of Gram stain of the joint aspirate.
Acute monoarticular In person who have risk factors for STD (Gonorrhea/ <i>Streptococcus</i> / <i>Staphylococcus</i> / Gram-negative Bacilli)	Ceftriaxone 2g IV q24h for 1-2 weeks PLUS Azithromycin 1g PO stat OR Doxycycline 100mg PO q12h for 7 days	Cefotaxime 2g IV q8h for 1-2 weeks PLUS Azithromycin 1g PO stat OR Doxycycline 100mg PO q12h for 7 days	
Polyarticular Gonorrhoea	Ceftriaxone 2g IV q24h for 7 days		
References: 1. Katie A. et.al. Curr Rheumatol Rep (2013) 15:332. 2. Mathews CJ et al Lancet 2010 375(9717): 846. 3. Barberi et al . CID2015. 4. George Varghese et al. BMJ 2010;341:c5470. 5. Michales pentali. EFORT Open Rev 2016.			
PROSTHETIC JOINT INFECTION			
Prosthetic Joint Infections (Empirical) Early: < 3 months after surgery <i>Staphylococcus aureus</i> Gram-negative Bacilli Delayed onset: from 3 to 12 months after surgery			Treatment concepts: 1. Empirical therapy is not recommended 2. Treatment is based on C&S 3. Rifampicin should never be used alone and should be started only after the clearance of bacteremia.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Less purulent organisms: CoNS/<i>Enterococcus</i>/ anaerobes Late onset: >12 months after surgery <i>Staphylococcus aureus</i> Enterobacterales Beta-hemolytic <i>Streptococcus</i> Anaerobes			4. Treatment strategy and duration of treatment depends on surgical strategy.
Definitive Prosthetic Joint infection treatment Methicillin-sensitive <i>Staphylococcus aureus</i> (MSSA)	Initial Treatment: Cloxacillin 2g IV q4-6h OR Cefazolin 2g IV q6-8h PLUS Rifampicin 600mg PO q24h or 450mg PO q12h		Duration: 2-6 weeks (according to treatment strategy) Followed by an oral combination therapy according to susceptibility. Rifampicin should be included if the implant is <i>in situ</i>.
Definitive Prosthetic Joint Infection treatment Methicillin-resistant <i>Staphylococcus aureus</i> (MRSA)	Initial treatment: Vancomycin 15-20mg/kg (actual body weight) IV q8-12h; not to exceed 2g/dose PLUS Rifampicin 300-450mg PO q12h		Duration: 2-6 weeks (according to treatment strategy) Followed by an oral combination therapy according to susceptibility. Rifampicin should be included if the implant is <i>in situ</i>.
DIABETIC FOOT INFECTIONS			
Antibiotics should not be used unless there are local or systemic symptoms of infection. Local treatment including surgical debridement is important. Antibiotic selection should be based on the most recent culture and sensitivity report.			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Mild Infections: a. Local infection involving skin & subcutaneous tissues b. Erythema, less than 2 cm around the ulcer c. No systemic signs of infection	Amoxicillin/clavulanate 625mg PO q8h OR Ampicillin/sulbactam 375-750mg PO q12h	Cephalexin 500mg PO q6h PLUS Metronidazole 400mg PO q8h	Duration: 5-7 days
Moderate Infection: a. Deep tissue infection b. Erythema more than 2 cm around ulcer c. No SIRS	Ampicillin/sulbactam 3g IV q6-8h If <i>Pseudomonas</i> is suspected: *Piperacillin/tazobactam 4.5g IV q6-8h	Cefazolin 2g IV q8h PLUS Metronidazole 500mg IV q8h Penicillin allergy: Ciprofloxacin 400mg IV q8-12h PLUS Clindamycin 600mg IV q8h	Duration: 7-14 days Modify according to clinical response. If proven osteomyelitis or margin of resection is inadequate: at least 4-6 weeks. However, a shorter duration (1-2 weeks) is sufficient if the margin of surgical resection is adequate. *Piperacillin/tazobactam: If given as q8h, to be given as extended infusion (over 3-4 hours).
Severe Infections: All of the above 2 or more SIRS -History of previous antibiotic exposure - Recurrent admission -Risk of pseudomonas infection	*Piperacillin/tazobactam 4.5g IV q6-8h If given as q8h, to be given as extended infusion (over 3-4 hours)	Cefepime 2g IV q8h PLUS Metronidazole 500mg IV q8h	Surgical debridement is URGENT. Based on intraoperative C&S, antibiotics should be streamlined. Duration: 7-14 days (subject to clinical improvement)

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
-Immunocompromised			If proven osteomyelitis or margin of resection is inadequate: at least 4-6 weeks A shorter duration of antibiotics can be considered if the osteomyelitis is fully resected (i.e. amputation with a clear margin): -No surrounding soft tissue infection: 5 days -Evidence of soft tissue infection: 10-14 days
NECROTIZING FASCIITIS			
Type 1 Polymicrobial infection Primarily occurs in patients who are immunocompromised or have certain chronic diseases such as diabetes	*Piperacillin/tazobactam 4.5g IV q6-8h PLUS/MINUS **Clindamycin 600-900mg IV q8h	Cefotaxime 2g IV q8h PLUS Metronidazole 500mg IV q8h OR **Clindamycin 600-900mg IV q8h OR Ampicillin/sulbactam 3g IV 6-8h PLUS/MINUS **Clindamycin 600-900mg IV q8h	*Piperacillin/tazobactam: If given as q8h, to be given as extended infusion (over 3-4 hours). **Clindamycin: Only necessary if risk of Group A <i>Streptococcus</i>/presence of gas crepitus or suspected toxic shock syndrome. Immediate aggressive surgical debridement is the primary treatment modality Repeated surgical debridement for source control is normally necessary.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			Urgent Gram stain Based on intraoperative culture & sensitivity, antibiotics should be streamlined.
Type 2 Monomicrobial infection Group A <i>streptococcus</i> (most common)	Benzylpenicillin 2-4MU IV q4h PLUS *Clindamycin 600-900mg IV q8h		*Clindamycin: Only necessary if risk of Group A <i>Streptococcus</i> /presence of gas crepitus or suspected toxic shock syndrome. Duration: 7-14 days (subject to clinical assessment)
<i>Vibrio vulnificus</i> <i>Aeromonas hydrophila</i> Consider in water related injuries and patients with liver cirrhosis and ingestion of raw oysters	Ceftriaxone 1g IV q12h PLUS Doxycycline 100mg PO q12h	Ciprofloxacin 400mg IV q8h	
Fournier's Gangrene Common organisms: <i>Eschericia coli</i> <i>Klebsiella</i> <i>Proteus</i> <i>Enterococcus</i> <i>Pseudomonas</i> Anaerobes	Ampicillin/sulbactam 3g IV q6-8h	*Piperacillin/tazobactam 4.5g IV q6-8h OR Ceftriaxone 2g IV q24h PLUS Metronidazole 500mg IV q8h	Aggressive surgical debridement is necessary to remove all necrotic tissues. *Piperacillin/tazobactam: if given as q8h, to be given as extended infusion (over 3-4 hours)

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
SOFT TISSUE INFECTION SECONDARY TO GAS PRODUCING ORGANISM			
Common organisms: <i>Clostridium</i> sp Gram-negative organisms	BenzyIpenicillin 4MU IV q4h PLUS Clindamycin 600-900mg IV q6h PLUS/ MINUS *Gentamicin 5mg/kg IV q24h	Cefotaxime 2g IV q8h PLUS Clindamycin 600-900mg IV q6h	Duration: 10-28 days *Gentamicin: if suspected Gram-negative infection. early aggressive surgical debridement is essential.
SUPPURATIVE WOUND INFECTIONS, SURGICAL OR TRAUMATIC			
Suppurative wound infections, surgical or traumatic	If there is surrounding cellulitis and/or systemic symptoms are present: Cloxacillin 500mg PO/IV q6h PLUS/MINUS Gentamicin 5mg/kg IV q24h (if gram negative organisms suspected or known to be involved) OR As a monotherapy: Cefuroxime 1.5g IV q8h		Change antibiotics accordingly after C&S results are available. Topical antibiotics are not recommended for treatment of wound infections as it may result in the emergence of resistant organisms. Patient tetanus immunisation status should be assessed in all cases.
MUSCULAR, SKELETAL AND SOFT TISSUE TRAUMA, CRUSH INJURIES AND STAB WOUNDS			
Muscular, skeletal and soft tissue trauma, crush injuries and stab wounds	Cloxacillin 2g IV q6h PLUS/MINUS *Metronidazole 500mg IV q6h PLUS/MINUS **Gentamicin 5mg/kg IV q24h	Cefazolin 2g IV q6-8h OR Cefuroxime 1.5g IV as a loading dose, followed by 750mg IV q8h PLUS	*Metronidazole: in soil/rust contamination or heavy machinery. **Gentamicin: if there is extensive skin and soft tissue involvement

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
		*Metronidazole 500mg IV q8h	Thorough surgical debridement, soft tissue and fracture stabilization. For severe penetrating injuries, especially those involving joints and/or tendons, antibiotics must be given for at least 5 days.
COMPOUND FRACTURES/OPEN FRACTURES			
Compound fractures: Antibiotics are administered as prophylaxis within 3 hours of injury.			
Gustilo I & II fractures	Cefazolin 1-2g IV q8h OR Cefuroxime 1.5g IV q8h	Amoxicillin/clavulanate 1.2g IV q8h	Pre-debridement and post-debridement cultures are not representative of actual infection.
Gustilo III fractures Mostly nosocomial and Gram-positive	As per Gustilo 1 & 2 fractures PLUS *Gentamicin 3-5mg/kg IV stat dose PLUS/MINUS **Metronidazole 500mg IV q8h		Duration of antibiotic for open fractures classification: - Gustilo type 1: stop after 24 hrs - Gustilo type II: stop after 24 hrs to 48 hrs - Gustilo type III: 24 hrs after wound closure or up to a maximum of 72 hrs (whichever is earlier) *Gentamicin: if initial debridement is expected to last more than 2 hours, a higher dose of gentamicin 5mg/kg IV stat is needed. **Metronidazole: in soil/rust contamination or heavy machinery.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			If soft tissue injury is of concern, to follow antibiotic guidelines for soft tissue injury.
References: 1. Zimmerli et al. NEJM 2004; 14;351;1645. 2. Del Pozo JL. NEJM.2009 361(8): 787 3. IDSA guidelines, Clinical Infectious Diseases ; 2012 ; 54 : 132-173 2012 4. Luca L.et al.International Journal of Infectious Diseases (2005) 9, 127 5. Michealis et al.. EFORT Open Rev 2016;1:128 6. Dennis L et al. N Engl J Med 377;23 7. Nayagam S. et al. British Orthopaedic Association Standards for Trauma. 2009			
C. NEUROSURGERY			
Antibiotic usage is not recommended for: - Traumatic CSF fistula - Post-surgical CSF leak - Basal skull fractures However, it should be highlighted that these groups of patients have increased risks of meningitis due to increased possibility of bacteria from the paranasal sinuses, nasopharynx and the ear canal making direct contact with the CNS as well as cerebrospinal fluid (CSF) leak, rendering antibiotic use needed in some situations. CSF leak would resolve spontaneously on day 5-7, or with the help of external lumbar drain. Patients with persistent CSF leak should have CSF cultures to guide antibiotic therapy. Patients with clinical presentation consistent with meningitis such as fever/altered consciousness/photophobia/neck stiffness should be treated with empiric antibiotics until culture results are available.			
Compound depressed skull fractures/Open skull fracture	Cefuroxime 1.5g IV q8h PLUS Metronidazole 500mg IV q8h		Duration: 5-7 days Review tetanus status of patients and consider vaccination.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			Surgical debridement within 12-24 hours is important to remove the source of infection.
Penetrating craniocerebral injuries	Ceftriaxone 2g IV q12h PLUS Metronidazole 400mg PO q8h		Duration: 2 weeks initially and then review with microbiology
References: 1. Salford Royal, NHS. Antibiotic Prophylaxis in Cranial Neurosurgery Antibiotic Guidelines, Unique ID: 144TD(C)25(F4) Issue number: 6, 2018.			

TROPICAL INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
1. Typhoid Fever (<i>Salmonella enterica</i> serovar Typhi and Paratyphi)			
Uncomplicated	Ceftriaxone 50-75mg/kg/24h (2-4 g/day) IV q12-24h for 5-7 days	Ciprofloxacin 500mg PO q12h for 5-7 days OR Azithromycin 1g PO stat, then 500mg PO q24h for 5-7 days	Ceftriaxone is recommended for quinolone resistance. Recommend for early IV to oral switch once symptoms improved or stabilised.
Complicated/Severe (patients with systemic toxicity, depressed consciousness, prolonged fever, organ system dysfunction or other features that prompt hospitalization)	Ceftriaxone 50-75mg/kg/24h (2-4 g/day) IV q12-24h for 10-14 days OR Cefotaxime 40-80mg/kg/24h (2-6 g/day) IV q8-12h for 10-14 days PLUS/MINUS *Dexamethasone 3mg/kg IV followed by 1mg/kg IV q6h for 2 days	Ciprofloxacin 400mg IV q12h for 10-14 days OR **Meropenem 1-2g IV q8h for 10-14 days PLUS/MINUS *Dexamethasone 3mg/kg IV followed by 1mg/kg IV q6h for 2 days	*Indication of dexamethasone (adjunctive): I. Typhoid psychosis II. Septic shock and other indications (discuss with ID physician) **Carbapenems should be reserved for patients who have severe or complicated infection with a suspected extensively drug-resistant strain. Recommend for early IV to oral switch once symptoms improved or stabilised
2. Cholera (<i>Vibrio cholerae</i>)			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Non-Tetracycline resistance	Doxycycline 300mg PO stat	Ciprofloxacin 1g PO stat	Rehydration is the mainstay of cholera treatment. Avoid antidiarrheal agents e.g. Diphenoxylate/atropine (Lomotil®) or Loperamide Indication for antibiotics: i) Severely ill patients especially who are severely or moderately dehydrated and continue to pass a large volume of stool during rehydration treatment. ii) All patients who are hospitalised and moderate to severe cases. *Azithromycin/Erythromycin: Recommended alternatives for pregnant women
Tetracycline resistance	Ciprofloxacin 1g PO stat OR *Azithromycin 1g PO stat	*Erythromycin ethylsuccinate 800mg PO q12h for 3 days	
3. Scrub Typhus (<i>Orientia tsutsugamushi</i> (<i>Rickettsia tsutsugamushi</i>))			
Uncomplicated	Doxycycline 100mg PO q12h for 7 days	*Azithromycin 500mg PO stat	*Azithromycin: Recommended for pregnant women
Complicated (ARDS, septic shock, myocarditis, meningoencephalitis, hepatitis, renal failure)	*Azithromycin 500mg IV q24h for 5 days	Rifampicin 600mg PO q24h for 5 days (if not responding to azithromycin)	*Recommend for early IV to oral switch once symptoms improve or stable
4. Brucellosis			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
<i>Brucella melitensis</i> , <i>Brucella abortus</i> , <i>Brucella suis</i> and <i>Brucella canis</i>	Doxycycline 100mg PO q12h for 6 weeks PLUS Streptomycin 1g (15mg/kg) IM q24h for 2 - 3 weeks	Doxycycline 100mg PO q12h for 6 weeks PLUS Gentamicin 5mg/kg/24h IV for 7 days OR Doxycycline 100mg PO q12h for 6 weeks PLUS Rifampicin 600-900mg (15mg/kg) PO q24h for 6 weeks OR *Rifampicin 600-900mg (15mg/kg) PO q24h for 6 weeks PLUS *Trimethoprim/sulfamethoxazole 160/800mg PO q12h for 6 weeks	Longer duration (up to 12 weeks) may be required in complicated cases e.g. spondylitis, neurobrucellosis, infective endocarditis, localised suppurative lesions *Recommended alternative for pregnant women. Optimal therapy for pregnant women has not been determined. Trimethoprim/sulfamethoxazole has been used with success. Another alternative is rifampicin monotherapy for at least 45 days.
5. Leptospirosis (<i>Leptospira</i> sp.)			
Mild to Moderate disease	Doxycycline 100mg PO q12h for 5-7 days	*Azithromycin 500mg PO q24h for 3 days	*Recommended alternative for pregnant women
Severe disease (Leptospiral pulmonary syndrome, multiorgan involvement, sepsis)	Benzylpenicillin 1.5MU IV q6h OR Ceftriaxone 1-2g IV q24h for 7 days		May consider Methylprednisolone 500-1000 mg IV for 3 days if pulmonary haemorrhage is present. However, there is insufficient evidence to support the routine use of corticosteroid.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
6. Tetanus			
<i>Clostridium tetani</i>	Metronidazole 500mg IV q6h-q8h for 7-10 days PLUS Human Tetanus Immunoglobulin 3000- 6000IU IM stat PLUS Anti-tetanus toxoid vaccine IM (initiate age appropriate active immunization at a different site)	Benzylpenicillin 100,000-200,000 unit/kg/24h IV q6h for 7-10 days PLUS Human Tetanus Immunoglobulin 3000- 6000IU IM stat PLUS Anti-tetanus toxoid vaccine IM (initiate age appropriate active immunization at a different site)	Human Tetanus Immunoglobulin 500IU might be as effective as higher doses of 3,000 to 6,000IU and causes less discomfort. All patients with tetanus should undergo wound debridement to eradicate spores and necrotic tissues.
7. Melioidosis (<i>Burkholderia pseudomallei</i>)			
Intensive Therapy (uncomplicated)	Ceftazidime 100-120mg/kg/24h IV q6h-q8h (usual dose: 2g IV q6h) PLUS/MINUS *Trimethoprim/sulfamethoxazole • < 40kg: 160/800mg IV/PO q12h • 40-60kg: 240/1200mg IV/PO q12h • >60kg: 320/1600mg IV/PO q12h		*Add on trimethoprim/sulfamethoxazole in eye, neurologic, testicular, prostatic, pericardium, bone and joint melioidosis. Drainage of abscesses should be attempted where ever appropriate such as prostatic, empyema and pericardium.
Intensive Therapy (complicated)	Meropenem 75mg/kg/24h IV q8h (with CNS involvement: 2g IV q8h; without CNS involvement: 1g IV q8h) OR Imipenem 50mg/kg/24h IV q6h (usual dose: 500-1000mg q6h) PLUS/MINUS *Trimethoprim/sulfamethoxazole • < 40kg: 160/800mg IV/PO q12h		Duration of intensive therapy: • Skin, bacteraemia with no foci, mild pneumonia: 2 weeks • Complicated pneumonia, prostatic, deep-seated foci, septic arthritis: 4 weeks • Osteomyelitis: 6 weeks

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	● 40-60kg: 240/1200mg IV/PO q12h ● >60kg: 320/1600mg IV/PO q12h (to deescalate to Ceftazidime once symptoms improve/stable)		● Neurologic/CNS: 8 weeks Folic Acid 5mg PO q24h to be given for patients on trimethoprim/sulfamethoxazole.
Eradication/Maintenance Therapy	Trimethoprim/sulfamethoxazole ● < 40kg: 160/800mg PO q12h ● 40-60kg: 240/1200mg PO q12h ● >60kg: 320/1600mg PO q12h	Amoxicillin/clavulanate ● ≤ 60kg: 1250mg (2 tabs of 625mg) PO q8h ● > 60kg: 1875mg (3 tabs of 625mg) PO q8h	Duration of eradication therapy: ● Osteomyelitis, Neurologic/CNS: 24 weeks ● Others: minimum 12 weeks
8. Malaria			
WHO recommends the use of Artemisinin Combination Therapy (ACT) as the standard treatment for malaria and discourages the prescription of monotherapy or substandard ACT as this will promote resistance. <u>Features of severe/complicated Malaria includes at least one of the following clinical or laboratory features:</u> <u>Clinical manifestation:</u> Impaired consciousness or unrousable coma Prostration (generalised weakness so that the patient is unable to walk or sit up without assistance) Failure to feed/not tolerating orally Convulsion Deep breathing, respiratory distress (acidotic breathing) Circulatory collapse or shock Clinical jaundice and evidence of other vital organ dysfunction Haemoglobinuria Abnormal spontaneous bleeding Pulmonary oedema (radiological) <u>Laboratory test:</u>			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Hypoglycaemia, metabolic acidosis, severe normocytic anaemia, haemoglobinuria, hyperparasitaemia (>20,000 parasites/microL for <i>P. knowlesi</i> or >100,000 parasites/microL for other <i>Plasmodium</i> species), hyperlactatemia or renal impairment.			
<i>Plasmodium falciparum</i> A. Non-complicated i) New Infection	Riamet® (1 tablet: Artemether/lumefantrine 20/120mg) The patient should receive an initial dose, followed by a second dose 8 hours later, then 1 dose q12h for the following 2 days. <15kg: 1 tab per dose 15 - <25kg: 2 tab per dose 25 - <35kg: 3 tab per dose ≥35kg: 4 tab per dose (Riamet should be taken with a fatty meal e.g milk to increase bioavailability)		Primaquine 0.75mg/kg (max: 45mg) to be given on Day 1 as a single dose (gametocytocide) except in pregnant/lactating women. If G6PD status unknown before use, to use 0.25mg/kg. Pregnancy: -Pregnant women with uncomplicated <i>P. falciparum</i> malaria should be treated with artemether-lumefantrine during the first trimester. -Exposure of Artemisinin derivatives during the second and third trimester has shown no adverse effects on the mother or fetus. Lactating women: Do not use Primaquine and Tetracycline
<i>Plasmodium falciparum</i> B. Complicated/Severe	Artesunate 2.4mg/kg IV at 0 hour, 12 hour, 24 hour and q24h till day 7 PLUS Doxycycline 100mg PO q12h for 7 days (given together with IV Artesunate)		Primaquine 0.75mg/kg (max 45mg) to be given on Day 1 of treatment (gametocytocide) in addition to Artemisinin regime and Quinine except for pregnant women. If possible, check G6PD status before use.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	Parenteral artesunate should be given for a minimum of 24 hours (3 doses) or until the patient can tolerate orally then it can be switched to a complete course of oral ACT regime.		Pregnancy (all stage): Artesunate IV as for normal adults
<i>Plasmodium vivax</i> <i>Plasmodium ovale</i>	Riamet® (dosing as per <i>Plasmodium falciparum</i> treatment) PLUS *Primaquine 0.5mg/kg (max 30mg) PO q24h for 14 days		*G6PD deficiency: Primaquine 0.75mg/kg PO once a week for 8 weeks. If significant haemolysis occurs, it should be stopped. Pregnancy: Full course Riamet® to be given, followed by chloroquine 300mg PO once a week till delivery. Full course of primaquine only to be given post- delivery. In severe <i>P. vivax</i> or <i>P. ovale</i> infections, the treatment is as severe as <i>P. falciparum</i> .
<i>Plasmodium malariae</i> <i>Plasmodium knowlesi</i>	Riamet® (dosing as per <i>Plasmodium falciparum</i> treatment)		In severe <i>P. malariae</i> or <i>P. knowlesi</i> infections, the treatment is as severe as <i>P. falciparum</i> . No radical cure with primaquine is required because no hypnozoites are formed in infection with these species.
Mixed Infection	Treat as <i>Plasmodium falciparum</i>		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Chemoprophylaxis	Doxycycline 100mg PO q24h Start: 1-2 days before departure Stop: 4 weeks after travel Maximum duration: 2 years OR *Atovaquone/proguanil (Malarone®) 100/250mg q24h Start: 1-2 days before departure Stop: 7 days after travel	**Mefloquine 250mg PO once a week Start: 2 weeks before departure Stop: 4 weeks after travel Maximum duration: 1 year	Pregnancy: Only Mefloquine can be used Refer to the drug resistance pattern and recommended prophylaxis in the travelling destination (CDC website). *Requires special medicine approval **Not available in HRPB Formulary
References: 1. CDC 2022. Recommendations for the Use of Antibiotics for the Treatment of Cholera. https://www.cdc.gov/cholera/treatment/antibiotic-treatment.html 2. WHO Global Task Force on Cholera Control 2020. https://choleraoutbreak.org/book-page/section-7-case-management-treatment-facilities 3. Rahi M, et al. DHR-ICMR Guidelines for Diagnosis & Management of Rickettsial Diseases in India. Indian J Med Res. 2015;141(4):417-22. 4. Kim YS et al. A comparative trial of single dose azithromycin vs doxycycline for treatment of old mild scrub typhus. CID. 2004; 39(9):1329-35 5. WHO. Brucellosis in humans and animals. 2006 6. MOH Malaysia. CPG Brucellosis. 2012 7. CDC Yellow Book 2019: Brucellosis. https://wwwnc.cdc.gov/travel/yellowbook/2020/travel-related-infectious-diseases/brucellosis 8. WHO. Leptospirosis. 2013. 9. CDC Yellow Book 2019: Leptospirosis. https://wwwnc.cdc.gov/travel/yellowbook/2020/travel-related-infectious-diseases/leptospirosis 10. Phimda K, et al. Doxycycline vs azithromycin for treatment of leptospirosis and scrub typhus. Antimicrob Agents Chemother. 2007;51(9):3259-63 11. Panaphut, T et al. Ceftriaxone compared with s.penicillin G for treatment of severe leptospirosis. CID. 2003;36(12), 1507-1 12. Suputtamongkol Y, et al. An open RCT comparing penicillin, doxycycline and cefotaxime for patients with severe leptospirosis. CID 2004;39(10):1417-24 13. Shenoy VV et al. Pulmonary leptospirosis: an excellent response to bolus methylprednisolone. Postgrad Med J. 2006; 82(971): 602–6 14. Rodrigo C et al. High dose corticosteroids in severe leptospirosis: a systematic review. Trans R Soc Trop Med Hyg. 2014;108(12):743-50 15. WHO. Current recommendations for treatment of tetanus. 2010. 16. CDC. Tetanus, Clinical information for clinicians. 2017			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
17. Currie B. Melioidosis: The 2014 Revised RDH Guideline. The Northern Territory Disease Control Bulletin. 2014;21(2): 4-8. 18. Dance D. Treatment and prophylaxis of melioidosis. Int J Antimicrob Agents 2014; 43: 310-8. 19. White NJ, et al. Halving of mortality of severe melioidosis by ceftazidime. Lancet. 1989;2: 697–701 20. Simpson AJ, et al. Comparison of imipenem and ceftazidime as therapy for severe melioidosis. CID. 1999;29:381–7 21. Management Guideline of Malaria in Malaysia 2013 22. WHO Guideline for Malaria 2023			

URINARY TRACT INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
1. CYSTITIS			
Uncomplicated Cystitis Common organisms: <i>Escherichia coli</i> Enterobacterales: <i>Klebsiella</i> , <i>Proteus</i> <i>Staphylococcus saprophyticus</i> In non-pregnant, pre-menopausal women with structurally and functionally normal urinary tract.	*Nitrofurantoin 50-100mg PO q6h (macrocrystals) OR Cephalexin 500mg PO q12h	Cefuroxime 250mg PO q12h OR Amoxicillin/clavulanate 625mg PO q8h OR Ampicillin/sulbactam 375-750mg PO q12h OR **Fosfomycin 3g PO x 1 dose	Urine culture is indicated only if symptoms are unresolved or recur. *Nitrofurantoin should not be used if GFR < 30 mL/min. **Consider Fosfomycin for patients suspected to have MDR Gram-negative infections. Ciprofloxacin and other quinolones are not recommended as empirical treatment in UTIs due to selection of resistance, and potential serious adverse events. Consider use of quinolones only in patients with history of anaphylactic reaction to beta-lactam antibiotics. Duration of treatment: 5-7 days
Cystitis in Pregnancy	*Nitrofurantoin 50-100mg PO q6h (macrocrystals) OR Cephalexin 500mg PO q12h	Cefuroxime 250mg PO q12h OR **Amoxicillin/clavulanate 625mg PO q8h OR	Obtain urine culture before starting treatment and repeat 1-2 weeks after completion of antibiotics to ensure eradication.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
		Ampicillin/sulbactam 375-750mg PO q12h OR ***Fosfomycin 3g PO x 1 dose	Duration of treatment: 5-7 days (treat for 7 days if recurrent) *Avoid Nitrofurantoin at third trimester if other options are available due to small risk of haemolytic anaemia in newborn. **Amoxicillin/clavulanate is generally safe in pregnancy (Category B), but there may be an increased risk of necrotizing enterocolitis associated with use in preterm, premature rupture of membranes. Avoid fluoroquinolones in pregnancy. ***Consider Fosfomycin for patients suspected to have MDR Gram-negative infections.
2. PYELONEPHRITIS			
Uncomplicated Pyelonephritis Common organisms: Enterobacterales Enterococci In non-pregnant, pre-menopausal women	<u>Outpatient treatment:</u> Amoxicillin/clavulanate 625mg PO q8h for 14 days OR Ampicillin/sulbactam 375-750mg PO q12h for 10-14 days		Obtain culture before starting treatment. Perform ultrasound of the upper urinary tract to exclude obstructive pyelonephritis.
	<u>Inpatient treatment:</u>		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
without urological abnormalities or comorbidities	Amoxicillin/clavulanate 1.2g IV q8h for 10-14 days OR Cefuroxime 750mg IV q8h OR Ampicillin/sulbactam 3g IV q8h	Ceftriaxone 1g IV q24h for 10 days	May step down to oral antibiotics guided by culture and sensitivity results once can tolerate orally and afebrile > 48 hours.
Pyelonephritis in other categories (e.g. pregnancy)	Treat as inpatient treatment for uncomplicated pyelonephritis		
3. OTHER URINARY TRACT INFECTIONS (UTIs)			
Complicated UTIs Common organisms: Enterobacterales Enterococci <i>Pseudomonas</i> sp. UTI symptoms in men OR presence of a structural or functional abnormality: ● Urinary tract obstruction ● Chronic kidney disease ● Poorly-controlled type 2 diabetes ● Immunosuppression ● Urinary catheter <i>in situ</i> ● Neurogenic bladder ● Post-menopausal women ● History of recurrent UTIs	Oral Therapy: Amoxicillin/clavulanate 625mg PO q8h for 7 days OR Cephalexin 500mg PO q6h for 7 days		Obtain urine culture before starting treatment. Treat for 10-14 days in patients with upper tract symptoms, delayed response or sepsis.
	Parenteral Therapy: Amoxicillin/clavulanate 1.2g IV q8h OR Ampicillin/sulbactam 3g IV q8h OR Cefuroxime 750mg IV q8h PLUS/MINUS Aminoglycoside	Ceftriaxone 1g IV q24h PLUS/MINUS Aminoglycoside	May step down to oral antibiotics guided by culture and sensitivity results once can tolerate orally and afebrile > 48 hours.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Nephrolithiasis			
Asymptomatic Bacteriuria (ABU) Urine bacterial growth $\geq 10^5$cfu/mL in 2 serial samples in women or a single sample in men without UTI symptoms.	Screening and treatment is ONLY recommended in the following patients: - pregnant women OR - patients undergoing invasive urological procedures i.e. prior to transurethral resection of prostate (TURP) or urological procedures breaching the mucosa (refer “surgical prophylaxis” for treatment) Whenever indicated, treatment should be guided by urine culture and sensitivity results.		Duration of treatment for pregnant women: 5-7 days Antibiotics to consider in pregnant women include amoxicillin, nitrofurantoin and oral cephalosporin.
Catheter-associated UTI (CA-UTI) Urine colony count $\geq 10^3$cfu/mL with at least one sign or symptom compatible with UTI, with no other identifiable source of infection.	Refer to “Complicated UTIs”		Routine screening and treating asymptomatic catheterized patients are not recommended. Pyuria alone in the absence of other symptoms is not diagnostic of CA-UTI. Remove unnecessary catheters. Whenever indicated, change the catheter prior to starting treatment. Treat for 7 days for prompt resolution of symptoms or 10-14 days if delayed response or sepsis.
Prophylaxis for Recurrent Urinary Tract Infections (rUTIs)	Nitrofurantoin 50-100mg PO ON (macrocrystals) OR Cephalexin 250mg PO ON		Antimicrobial prophylaxis is indicated if non-antimicrobial measures fail. Post-coital prophylaxis may be appropriate for sexually-related rUTIs.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
2 episodes/6 months OR > 3 episodes/year			When primary options fail, consider daily antibiotic prophylaxis guided by culture and sensitivity results. Duration: 3-6 months
4. UROLOGY			
Pyonephrosis/Perinephric Abscess/Renal Abscess Common organisms: Enterobacterales Enterococci <i>Staphylococcus aureus</i> <i>Pseudomonas</i> sp.	Amoxicillin/clavulanate 1.2g IV q8h OR Cefuroxime 750mg IV q8h OR Ampicillin/sulbactam 3g IV q6-8h PLUS/MINUS Gentamicin 5mg/kg IV q24h	Ceftriaxone 2g IV q24h PLUS/MINUS Gentamicin 5mg/kg IV q24h	Obtain blood and urine cultures before starting treatment. Send pus for culture and sensitivity. Drainage the mainstay of treatment followed by definitive surgery if warranted. Step down to oral antibiotics guided by culture and sensitivity results once can tolerate orally and afebrile > 48 hours following catheter removal. Duration: 2-3 weeks (of both IV and oral) Longer course of antibiotic if: • Difficult to drain abscess • Slow resolution on follow-up imaging.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Acute Prostatitis Common organisms: <u>Age < 35 years old</u> • <i>N. gonorrhea</i> • <i>C. trachomatis</i> <u>Age ≥ 35 years old</u> • Enterobacterales • Coliforms • Enterococci Fever, chills, malaise, myalgia, dysuria, irritative urinary symptoms (frequency, urgency, urge incontinence), pelvic or perineal pain, and cloudy urine.	<u>Outpatient treatment:</u> Trimethoprim/sulfamethoxazole 160/800mg PO q12h OR Ciprofloxacin 500-750mg PO q12h		Obtain urine cultures before starting treatment. May step down to oral antibiotic once can tolerate orally and afebrile > 48 hours. Duration: 2 weeks or up to 4 weeks in severe illness or concomitant bacteraemia.
	<u>Inpatient treatment:</u> Amoxicillin/clavulanate 1.2g IV q8h OR Ampicillin/sulbactam 3g IV q6- 8h OR Cefuroxime 750mg IV q8h PLUS/MINUS Gentamicin 5mg/kg IV q24h	Ceftriaxone 1-2g IV q24h PLUS/MINUS Gentamicin 5mg/kg IV q24h	
Chronic Bacterial Prostatitis (NIH Type II) Chronic or recurrent urogenital symptoms that persist for at least 3 months. Relapsing UTI with repeated isolation of the same organism from urine is the hallmark.	Ciprofloxacin 500mg PO q12h	Trimethoprim/sulfamethoxazole 160/800mg PO q12h	Reassess after 2 weeks of antimicrobial therapy. Only continue antibiotics if pre-treatment cultures are positive and/or symptoms improve. Duration: 4-6 weeks
Epididymo-orchitis (non-STD related)	Ciprofloxacin 500mg PO q12h for a minimum of 2 weeks		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Common organisms: Enterobacterales Enterococci <i>Pseudomonas</i> sp. Acute onset, usually unilateral scrotal pain swelling with or without fever, rigors, and lower urinary tract symptoms.			
Epididymo-orchitis (STD related)	Refer to Sexually Transmitted Infections Section		
Testicular Abscess Common organisms: Enterobacterales Enterococci <i>Pseudomonas</i> sp.	Amoxicillin/clavulanate 1.2g IV q8h OR Ampicillin/Sulbactam 3g IV q6-8h OR Cefuroxime 750mg IV q8h PLUS/MINUS Gentamicin 5mg/kg IV q24h	Ceftriaxone 2g IV q24h PLUS/MINUS Gentamicin 5mg/kg IV q24h	Drainage is the mainstay of treatment. Send pus for culture and sensitivity.
References: 1. The Sanford Guide To Antimicrobial Therapy 2021. (Version 5.1.13) [Mobile app] 2. Guidelines on Urological Infections, European Association of Urology 2022, Available at: https://uroweb.org/guidelines/urological-infection (Accessed 10th June 2022) 3. Clinical Practice Guideline for the Management of Asymptomatic Bacteriuria: 2019 Update by the Infectious Diseases Society of America, <i>Clinical Infectious Diseases</i>, Volume 68, Issue 10, 15 May 2019, Pages e83–e110, Available at: https://doi.org/10.1093/cid/ciy1121 (Accessed 10th June 2022) 4. International Clinical Practice Guidelines for the Treatment of Acute Uncomplicated Cystitis and Pyelonephritis in Women: A 2010 Update by the IDSA and European Society for Microbiology and Infectious Diseases 2011. 5. Hooton TM. Urinary tract infections and asymptomatic bacteriuria in pregnancy. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA, Available at: https://www.uptodate.com/contents/urinary-tract-infections-and-asymptomatic-bacteriuria-in-pregnancy (Accessed 10th June 2022)			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
6.	Abdelmoteleb, Haitham & Rashed, Fatima & Hawary, Amr. (2017). Management of prostate abscess in the absence of guidelines. International braz j urol : official journal of the Brazilian Society of Urology. 43. 10.1590/S1677-5538.IBJU.2016.0472.		
7.	Reddivari AKR, Mehta P. Prostatic Abscess. [Updated 2022 May 12]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan-. Available at: https://www.ncbi.nlm.nih.gov/books/NBK551663/ (Accessed 10 th June 2022)		
8.	Ackerman A. L., Parameshwar P. S., Anger J. T. (2018). Diagnosis and treatment of patients with prostatic abscess in the post-antibiotic era. <i>Int. J. Urol.</i> 25, 103–110. Available at: https://onlinelibrary.wiley.com/doi/epdf/10.1111/iju.13451 (Accessed 10 th June 2022)		

INFECTIONS IN PATIENTS ON LONG-TERM RENAL REPLACEMENT THERAPY

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
A. HAEMODIALYSIS-RELATED CATHETER-RELATED BLOODSTREAM INFECTION (Renal impairment adjusted doses are used in this section)			
Empirical Therapy	Ceftazidime 2g IV stat and 1g IV q24h PLUS Cefazolin 2g IV stat and 1g IV q24h		Two sets of blood cultures should be obtained prior initiation of antibiotics. Remove catheter (immediately) if: - concurrent exit site infection (redness, pus) - ill patients with septic shock Linezolid IV is not an appropriate choice for empirical therapy
Definitive Therapy			
<i>Staphylococcus aureus</i> MSSA	Cefazolin 2g IV stat and 1g IV q24h		Duration: 4 to 6 weeks. However, if the catheter is removed and bacteraemia resolved within 72h, a shorter duration of 2 weeks may be appropriate. Transthoracic ECHO (TTE) should be pursued in the setting of <i>S. aureus</i> bacteraemia to rule out infective endocarditis.
MRSA*	Vancomycin 15-20mg/kg IV or 1g IV stat (refer TDM for level)		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			Repeat peripheral culture every 48-72h until culture is negative. *In MRSA infection, nasal swab for MRSA and decolonization with mupirocin 2% cream plus chlorhexidine 4% bath may be considered.
<i>Enterococcus</i>	Ampicillin 1-2g IV q12-24h OR Vancomycin 15-20mg/kg IV or 1g IV stat (refer TDM for level)	Linezolid 600mg IV q12h	The choice of antibiotic should be tailored to the culture and sensitivity results.
<i>Coagulase-negative staphylococci (CoNS)</i> Methicillin-sensitive	Cefazolin 2g IV stat and 1g IV q24h		In the absence of endovascular hardware (e.g. cardiac implant, foreign material in the body), only consider antibiotics if fever and/or bacteraemia persists after catheter removal. Duration: - If catheter is removed, for 5 to 7 days - If catheter salvage is necessary, for 14 days and repeat cultures for confirmation
Methicillin-resistant	Vancomycin 15-20mg/kg IV or 1g IV stat (refer TDM for level)		
<i>Corynebacterium, Bacillus, Micrococcus</i>	Vancomycin 15-20mg/kg or 1g IV stat (refer TDM for level)		Duration: 2 weeks The necessity of antibiotic therapy has to correlate with clinical infection.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Gram-negative bacilli Uncomplicated (e.g. <i>Klebsiella pneumoniae</i> , <i>E. coli</i>) <i>Pseudomonas</i> spp.*	Ceftazidime 2g IV stat and 1g IV q24h		Duration: 2 weeks *May consider a combination of antibiotics if suspected <i>Pseudomonas</i> spp. Infection. The choice of antibiotic should be tailored to the culture and sensitivity results.
<i>Burkholderia cepacia</i>	Trimethoprim/sulfamethoxazole 2.5-5mg/kg (TMP component) IV q24h		
<i>Candida</i> species	Fluconazole 200mg IV q24h*		*Except <i>Candida krusei</i> and <i>Candida glabrata</i> Duration: 14 days after first negative culture
Catheter removal is preferred especially if: (i) metastatic infection – septic arthritis, endocarditis (4-6 weeks antibiotics); osteomyelitis (6-8 weeks antibiotics) (ii) <i>S. aureus</i> , MDR organisms or fungal infection (iii) tunnel infection (iv) clinical picture not improving or repeated blood cultures remain positive If catheter removal is undesirable, for guidewire catheter exchange (after 72 hours of appropriate antibiotics)			
References: 1. Mermel LA, Allon M, Bouza E, et al. Clinical practice guidelines for the diagnosis and management of intravascular catheter-related infection: 2009 Update by the Infectious Disease Society of America. Clin Infect Dis. 2009;49:1-45 2. Abad CL, Safdar N. Catheter-related bloodstream infection. IDSE Special edition 2011. 3. Vanholder R, Canaud B, Fluck R <i>et al.</i> Diagnosis, prevention and treatment of haemodialysis catheter-related bloodstream infections (CRBSI): a position statement of European Renal Best Practice (ERBP). NDT Plus 2010;3:234-46. 4. Band JD. Treatment of intravascular catheter-related infections. In: UpToDate. (Accessed on October 18, 2017.)			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
B. PERITONITIS			
CAPD Peritonitis <i>S. aureus</i> CoNS <i>Pseudomonas aeruginosa</i> Gram-negative enterics	<u>First line:</u> <i>For anuric patients:</i> Cefepime 1g IP loading, then 250mg IP q6h <i>For oliguric patients:</i> Cefepime 1.25g IP loading, then 250/250/250/500mg IP q6h PLUS Antifungal prophylaxis: Nystatin suspension 500,000 unit PO q6h If no response, <u>Second line:</u> Continue Cefepime IP, add Amikacin 125mg IP ON PLUS Antifungal prophylaxis: Fluconazole 100-200mg PO q24h <u>Third line:</u> Discontinue Cefepime IP, continue Amikacin IP add Imipenem 500mg IP loading, then 100mg IP q6h add Vancomycin 1.5g IP loading, then 100mg IP q6h (anuric) or 125mg IP q6h (oliguric)	<u>First line:</u> Cloxacillin 1g IP loading, then 250mg IP q6h OR Cefazolin 1g IP loading, then 250mg IP q6h PLUS Ceftazidime 1g IP loading, then 250mg IP q6h PLUS Antifungal prophylaxis: Nystatin suspension 500,000 unit PO q6h	Prolonged courses of intraperitoneal aminoglycosides should be avoided. Adjunctive oral N-acetylcysteine therapy may help prevent aminoglycoside ototoxicity. Stability and compatibility of antibiotics in dextrose-based PD solution under room temperature: - Cefazolin - 8 days (7 days in Icodextrin) - Ceftazidime - 4 days (2 days in Icodextrin) - Vancomycin - 28 days (14 days in Icodextrin) - Cefepime, Amikacin, Imipenem - no data Aminoglycosides and penicillins should not be added to the same bag due to chemical incompatibility.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	PLUS Antifungal prophylaxis: Fluconazole 100-200mg PO q24h		
References: 1. ISPD Peritonitis Guideline Recommendations: 2022 Update on Prevention and Treatment. <i>Peritoneal Dialysis International</i> , Vol. 42(2), pp. 110–153 2. CPG RRT 2017 pg 236-237			

CHEMOPROPHYLAXIS

A. SURGICAL CHEMOPROPHYLAXIS

The goal of antibiotic prophylaxis is to prevent surgical site infection by reducing the burden of microorganisms at the surgical site during the operative procedure.

Single-dose prophylaxis is usually sufficient. If antibiotic prophylaxis is continued post-operatively, the duration should be less than 24 hours (up to 48 hours for cardiac surgery), regardless of the presence of intravascular catheters or indwelling drains.

If there are pre-existing infections (known or suspected), use appropriate treatment regimen instead of prophylactic regimen for procedure. However, re-dosing is required just prior to skin incision.

The optimal time for administration of pre-operative antibiotics is 60 minutes prior to surgical incision. Some agents, such as fluoroquinolones and vancomycin, require administration over one to two hours. Hence, the administration of these agents should begin within 120 minutes before surgical incision.

For patients with penicillin allergy, vancomycin or clindamycin is recommended unless stated otherwise.

An additional dose of prophylactic antibiotic during operation is indicated if:

- ❖ excessive blood loss (>1500 mL)
- ❖ procedures exceed two half-lives of the drug
- ❖ if there are other factors that may shorten the half-life of the prophylactic agent (e.g. extensive burns)

Antimicrobial	Recommended redosing interval with normal renal function (from initiation of preoperative dose), hr
Amoxicillin/clavulanate	3
Ampicillin/sulbactam	2
Benzylpenicillin	2
Cefazolin	4
Cefuroxime	4
Clindamycin	6
Gentamicin	NA
Metronidazole	NA
Vancomycin	NA

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
1. OBSTETRICS & GYNAECOLOGY			
A. Obstetrics			
Caesarean section a. Elective b. Emergency	Cefazolin 2g IV* OR Cefuroxime 1.5g IV PLUS/MINUS Metronidazole 500mg IV	Ampicillin/sulbactam 3g IV In patients with severe penicillin/cephalosporin allergy: Clindamycin 600-900mg IV PLUS Gentamicin 5mg/kg IV (If overweight or obese, based on adjusted body weight)	*Cefazolin 3g IV for patients weighing ≥120 kg Cefazolin 1g IV for patients weighing <80 kg **Clinical judgement is needed to evaluate a different regimen (higher dose, second dose) of prophylactic antibiotics in the presence of risk factors: - High BMI - Prolonged labour - Prolonged duration of surgery - Extensive surgical manipulation - Massive blood loss ***May be omitted if the patient is already receiving other equivalent broad spectrum antibiotics for other indications.
Manual removal of placenta	Cefazolin 2g IV* OR Cefuroxime 1.5g IV PLUS	Ampicillin/sulbactam 3g IV OR Amoxicillin/clavulanate 1.2g IV	As a single dose

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	Metronidazole 500mg IV	In patients with severe penicillin/cephalosporin allergy: Clindamycin 600-900mg IV PLUS Gentamicin 5mg/kg IV	
Repair of perineal tear: - First/second degree tears (including episiotomy repairs)	Antibiotic not indicated		
Repair of perineal tear: - Third or fourth degree tears	Cefazolin 2g IV* OR Cefuroxime 1.5g IV PLUS Metronidazole 500mg IV	Ampicillin/sulbactam 3g IV OR Amoxicillin/clavulanate 1.2g IV In patients with severe penicillin/cephalosporin allergy: Gentamicin 5mg/kg IV PLUS Metronidazole 500mg IV	Duration: 5-7 days
Cervical cerclage	Current evidence insufficient to recommend for or against antibiotics prophylaxis		
Operative vaginal delivery (forceps/vacuum-assisted delivery)	Amoxicillin/clavulanate 1.2g IV		As a single dose
B. Gynaecology			
Laparotomy: Hysterectomy	Cefazolin 2g IV* OR	Ampicillin/sulbactam 3g IV OR	Consider to give second or additional dose for:

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Laparoscopic: Hysterectomy (TLH/LAVH) or when need to access uterine cavity/vagina	PLUS Cefuroxime 1.5g IV Metronidazole 500mg IV	Amoxicillin/clavulanate 1.2g IV In patients with severe penicillin/cephalosporin allergy: Clindamycin 600-900mg IV PLUS Gentamicin 5mg/kg IV	- prolonged procedures (>3 hours) - blood loss >1500ml
Vaginal Hysterectomy			
Laparotomy: lymphadenectomy, omentectomy, cystectomy, oophorectomy, myomectomy, salpingectomy/tubal ligation (without entry into bowel/vagina)	Cefazolin 2g IV* OR Cefuroxime 1.5g IV PLUS/MINUS Metronidazole 500mg IV	Ampicillin/sulbactam 3g IV OR Amoxicillin/clavulanate 1.2g IV In patients with severe penicillin/cephalosporin allergy: Clindamycin 600-900mg IV PLUS Gentamicin 5mg/kg IV	
Laparoscopic surgery – without breaching bowel/uterine/vaginal cavity	Antibiotic not recommended		
Surgical termination of pregnancy	Antibiotic not indicated		
Hysteroscopy	Antibiotic not indicated		
Urogynaecology procedures (urethral sling/TVT, colposuspension, vaginal)	Cefazolin 2g IV* OR Cefuroxime 1.5g IV	Ampicillin/sulbactam 3g IV OR Amoxicillin/clavulanate 1.2g IV	

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
prolapse surgery +/- mesh/SSF)	PLUS Metronidazole 500mg IV	In patients with severe penicillin/cephalosporin allergy: Clindamycin 600-900mg IV PLUS Gentamicin 5mg/kg IV	
Large loop excision of transformation zone/knife cone biopsy	Current evidence insufficient to recommend for or against antibiotics prophylaxis		
Transvaginal oocyte retrieval	Doxycycline 200mg PO		
References: 1. Bratzler DW, Dellinger EP, Olsen KM et al. Clinical practice guidelines for antimicrobial prophylaxis in surgery. <i>Am J Health-Syst Pharm.</i> 2013; 70:195-283 2. Van Eyk N, van Schalkwyk J, Infectious Diseases Committee. J ObstetGynaecol Can. 2012 Apr; 34(4):382-391. 3. Van Schalkwyk J, Van Eyk N et al. No 247 – Antibiotic Prophylaxis in Obstetric Procedures. 4. Journal of Obstetrics and Gynaecology Canada, Volume 39, Issue 9, September 2017, Pages e300-e308. 5. WHO recommendation on prophylactic antibiotics for women undergoing caesarean section. Geneva: World Health Organization; 2021. Licence: CC BY-NC-SA 3.0 IG 6. National Antimicrobial Guideline 2019. Ministry of Health Malaysia 7. The management of third- and fourth-degree perineal tears. Greentop Guideline No 29, June 2015; Royal College of Obstetricians & Gynaecologist 8. ACOG practice bulletin No 195: Prevention of infection after gynecologic procedures. <i>Obstet Gynecol</i> 2018; 131: e172 9. ACOG practice bulletin No. 199: Use of prophylactic antibiotics in labour and delivery. <i>Obstet Gynecol</i> 2018;132: e103 10. WHO guidelines for the management of postpartum haemorrhage and retained placenta, 2009 11. Surgical Antibiotic Prophylaxis Guideline 2020. The Women's, The Royal Women's Hospital, Victoria, Australia. 12. Ekanem EE, Oniya O, Saleh H, Konje JC. Surgical site infection in obstetrics and gynaecology: prevention and management. <i>The Obstetrician & Gynaecologist</i> 2021; 23:124-37. https://doi.org/10.1111/tog.12730 13. HCTM PPUKM. Surgical Prophylaxis Guide 2018 14. Prophylactic antibiotics in the prevention of infection after operative vaginal delivery (ANODE): a multicentre RCT; <i>Lancet</i> 2019;393:2395-403			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
15. Weinreb et al. Should all oocyte donors receive prophylactic antibiotics for retrieval?; Fertility and Sterility Vol 94.No 7.Dec 2010 American Society for Reproductive Medicine.			
2. OTORHINOLARYNGOLOGY SURGERY			
Head and neck			
Clean	Antibiotic not required	Antibiotic not required	
Clean with placement of prosthesis (excludes tympanostomy tubes)	Amoxicillin/clavulanate 1.2g IV OR Cefazolin 2g IV (3g IV for patients weighing ≥120 kg)	Cefuroxime 1.5g IV	Duration: - One dose prior to skin incision - Post-operative duration variable depending on individual cases
Clean-contaminated cancer surgery Other clean-contaminated procedures with the exception of tonsillectomy and functional endoscopic sinus procedures	Amoxicillin/clavulanate 1.2g IV OR Cefazolin 2g IV (3g IV for patients weighing ≥120 kg) PLUS Metronidazole 500mg IV	Cefuroxime 1.5g IV PLUS Metronidazole 500mg IV	Duration: - One dose prior to skin incision - Post-operative duration variable depending on individual cases
References: 1. Simo R, French G. The use of prophylactic antibiotics in head and neck oncological surgery. <i>Curr Opin Otolaryngol Head Neck Surg.</i> 2006; 14:55-61 88 2. Surgical Prophylaxis Guide 2018, HUKM. 3. Vander Poorten V, Uyttebroek S, Robbins KT et al. Perioperative antibiotics in clean-contaminated head and neck surgery: a systematic review and meta-analysis. <i>Adv Ther.</i> 2020;37(4):1360-1380. 4. Iocca O, Copelli C, Ramieri G et al. Antibiotic prophylaxis in head and neck cancer surgery: systematic review and Bayesian network meta-analysis. <i>Head Neck.</i> 2022;44(1):254-261.			
3. ORAL/DENTAL SURGERY			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Clean surgery (Class 1) - Submandibular gland surgery - TMJ surgery - Excision of benign tumours/cysts	Not indicated for most surgeries May be indicated i. if the duration of the surgery is expected to be very long ii. for open reduction and internal fixation of facial bone fractures		Prophylaxis is recommended for all patients with an increased risk of surgical wound infection i.e. in immunocompromised patients *In patients with cardiac conditions with increased risk of infective Endocarditis, chemoprophylaxis is indicated. Please refer to the Non-Surgical Antibiotic Prophylaxis section .
Minor clean-contaminated surgery (Class 2) - Soft tissue surgery - Dentoalveolar surgery* - Periodontal surgery			
Minor Clean-contaminated surgery (Class 2) - insertion of dental implants and use of graft material - high degree of difficulty/long duration	Amoxicillin 1g PO OR Clindamycin 600-900mg IV/PO OR Benzylpenicillin 2MU IV	Amoxicillin/clavulanate 1.2g IV or 1.25g PO OR Cefuroxime 1.5g IV or 500mg PO	
Major clean-contaminated surgery (Class 3) - Orthognathic surgery - Excision / enucleation of large benign tumours/cysts - All oral cancer surgery - Open reduction and internal fixation of facial bone fractures	Benzylpenicillin 2MU IV OR Clindamycin 600-900mg IV	Amoxicillin/clavulanate 1.2g IV OR Cefuroxime 1.5g IV	For oral and maxillofacial fractures, antibiotics are recommended for the immediate post-trauma period and should be discontinued once open reduction and internal fixation are completed.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Reference: 1. Oral Health Division Ministry of Health Malaysia. Antibiotic Prophylaxis in Oral Surgery for Prevention of Surgical Site Infection. Putrajaya: Dental Technology Section Oral Health Division (OHD) Ministry of Health Malaysia; 2015.			
4. PLASTIC SURGERY			
Not indicated: for the majority of clean procedures, unless the patient has risk factors for postoperative infection (e.g. implantation of prosthetic material, prior skin irradiation). The continuation of antibiotics while waiting for non-infected skin grafts or flaps to epithelialize is not evidence-based.			
For clean-contaminated procedures	Cefazolin 2g IV (3g IV for patients weighing ≥120 kg)	Amoxicillin/clavulanate 1.2g IV	
Reference: 1. Antibiotic Expert Groups. Therapeutic guidelines: antibiotic. Version 15. Melbourne: Therapeutic Guidelines Limited; 2014.			
5. VASCULAR SURGERY			
Amputation of ischemic limb Suspected organisms: <i>Staphylococcus</i> spp. & anaerobic organisms	Ampicillin/sulbactam 3g IV	Amoxicillin/clavulanate 1.2gm IV	
Open and endovascular repair of abdominal aneurysm	Amoxicillin/clavulanate 1.2g IV	Penicillin allergy: Vancomycin 1g IV (1.5g IV for patients weighing ≥ 75 kg)	
Bypass surgery	Amoxicillin/clavulanate 1.2g IV	Penicillin allergy: Vancomycin 1g IV (1.5g IV for patients weighing ≥ 75 kg)	

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Arteriovenous graft	Amoxicillin/clavulanate 1.2g IV If high risk for MRSA: Vancomycin 1g IV (1.5g IV for patients weighing ≥ 75 kg)		MRSA risk – defined as history of MRSA colonisation or infection OR inpatient of high-risk hospital or unit (where MRSA is endemic).
Reference: 1. Bratzler DW, Dellinger EP, Olsen KM et al. Clinical practice guidelines for antimicrobial prophylaxis in surgery. Am J Health-Syst Pharm.2013; 70:195-283			
6. GENERAL SURGERY			
Procedures involving entry into lumen of gastrointestinal tract (bariatric, pancreaticoduodenectomy)	Cefazolin 2g IV (3g IV for patients weighing ≥120 kg)	Cefuroxime 1.5g IV	
Procedures without entry into gastrointestinal tract (antireflux, highly selective vagotomy) for high-risk patients	Cefazolin 2g IV (3g IV for patients weighing ≥120 kg)	Cefuroxime 1.5g IV	
Appendectomy for uncomplicated appendicitis Colorectal	Cefazolin 2g IV (3g IV for patients weighing ≥120 kg) PLUS Metronidazole 500mg IV OR Ampicillin/sulbactam 3g IV	Cefuroxime 1.5g IV PLUS Metronidazole 500mg IV Penicillin allergy: Clindamycin 600-900mg IV PLUS Gentamicin 5mg/kg	For patients undergoing elective colorectal surgery, a mechanical bowel preparation combined with neomycin PO PLUS erythromycin PO OR neomycin PO PLUS metronidazole PO should be given along with intravenous antibiotic prophylaxis. Preferred regimen: Neomycin 1g PO PLUS Metronidazole 400mg PO

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			for 2-3 doses in conjunction with mechanical bowel preparation *Preferred timing for 2 doses: 3:00 p.m. and 7:00 p.m. (after administration of oral bowel preparation)
Small intestine	Non-obstructed: Cefazolin 2g IV (3g IV for patients weighing ≥ 120 kg)	Cefuroxime 1.5g IV Penicillin allergy: Clindamycin 600-900mg IV PLUS Gentamicin 5mg/kg IV	
	Obstructed: Cefazolin 2g IV (3g IV for patients weighing ≥ 120 kg) PLUS Metronidazole 500mg IV	Cefuroxime 1.5g IV PLUS Metronidazole 500mg IV Penicillin allergy: Clindamycin 600-900mg IV PLUS Gentamicin 5mg/kg	
Hernia repair with mesh	Cefazolin 2g IV (3g IV for patients weighing ≥ 120 kg)	Amoxicillin/clavulanate 1.2g IV OR Ampicillin/sulbactam 3g IV	Includes laparoscopic repair.
Breast cancer surgery	Cefazolin 2g IV (3g IV for patients weighing ≥ 120 kg)	Amoxicillin/clavulanate 1.2g IV OR Ampicillin/sulbactam 3g IV	The benefits of routine postoperative antibiotic doses in reconstruction surgery are uncertain; there may be a benefit in obese patients or those treated with radiation therapy. The need for postoperative doses should be considered on an individual patient
Breast reshaping procedures	Cefazolin 2g IV (3g IV for patients weighing ≥ 120 kg)	Amoxicillin/clavulanate 1.2g IV OR Ampicillin/sulbactam 3g IV	

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			basis; if used, postoperative prophylaxis should not exceed 24 hours. ²
Breast surgery with implant (reconstructive or aesthetic)	Cefazolin 2g IV (3g IV for patients weighing ≥120 kg)	Amoxicillin/clavulanate 1.2g IV OR Ampicillin/sulbactam 3g IV	
References: 1. Bratzler DW, Dellinger EP, Olsen KM et al. Clinical practice guidelines for antimicrobial prophylaxis in surgery. Am J Health-Syst Pharm. 2013; 70:195-283 2. Antibiotic Expert Groups. Therapeutic guidelines: antibiotic. Version 15. Melbourne: Therapeutic Guidelines Limited; 2014. 3. Rollins KE, Javanmard-Emamghissi H, Acheson AG and Lobo DN. The role of oral antibiotic preparation in elective colorectal surgery. Annals of Surgery. 2019; 270(1), 43–58. 4. Kelleher DC, Sigle GW and Kumar AS. Bowel preparation before elective surgery. Clinics in Colon and Rectal Surgery. 2013; 26(03): 146–152.			
7. ORTHOPAEDIC SURGERY			
Clean operations involving hand, knee, or foot and not involving implantation of foreign materials	None		
Internal fixation of all closed fracture/total joint replacement/spine surgery (with and without instrumentation)/Arthroscopy	Cefazolin 2g IV (3g IV for patients weighing ≥120 kg)	Cefuroxime 1.5g IV Penicillin allergy: Clindamycin 600-900mg IV	The benefits of routine postoperative antibiotics are uncertain. If used, postoperative prophylaxis should not exceed 24 hours.
References: 1. Bratzler DW, Dellinger EP, Olsen KM et al. Clinical practice guidelines for antimicrobial prophylaxis in surgery. Am J Health-Syst Pharm.2013; 70:195-283 2. T.P Ruedi ,R.GBuckley,C.GMarani,AO principle of fracture management. A.H.R.W Simpson, BMJ 2015			
8. UROLOGY SURGERY			
A. Diagnostic Procedures			
Transrectal ultrasound and prostate biopsy	Ciprofloxacin 500mg PO q12h for 3 days (start 24 hours before procedure) PLUS/MINUS	Targeted antibiotic therapy based on preoperative rectal swab results.	Consider povidone-iodine bowel preparation to further decrease infection risk.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Common organisms: <i>Escherichia coli</i> , <i>Klebsiella</i> , <i>Proteus</i> , <i>Enterococcus</i> , <i>Pseudomonas</i>	Gentamicin 80mg IV single dose given 30-60 minutes before procedure.		
Cystoscopy/Urodynamics study	Antibiotic not recommended		Prophylaxis only for high-risk cases (immunocompromised patients, e.g. debilitated patients on long term catheters, patient with prosthetic/heart valves, diabetics, transplant recipients): Cefuroxime 250mg PO stat. If heart valve: Follow recommendation for SBE prophylaxis.
Retrograde pyelogram/Ureteral stenting	Cefuroxime 250mg PO stat		
B. Endourology			
Endourological surgery e.g. PCNL, URS, RIRS, TURP Common organisms: <i>Escherichia coli</i> , <i>Klebsiella</i> , <i>Proteus</i> , <i>Enterococcus</i> , <i>Pseudomonas</i>	Amoxicillin/clavulanate 1.2g IV OR Ampicillin/sulbactam 3g IV	Cefuroxime 1.5g IV OR Ceftazidime 2g IV (if urine grew <i>Pseudomonas</i>)	Antibiotic selection to be determined based on the patient's latest urine culture result.
C. Open Surgery			
Clean operations e.g. orchidectomy, orchidopexy, varicocelectomy, deroofing renal cysts	Antibiotic not required		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Clean-contaminated (with opening of urinary tract) e.g. nephrectomy, prostatectomy, open stone surgery. Common organisms: <i>Escherichia coli</i> , <i>Klebsiella</i> , <i>Proteus</i> , <i>Enterococcus</i> , <i>Pseudomonas</i>	Amoxicillin/clavulanate 1.2g IV q8h for 1 day OR Ampicillin/sulbactam 3g IV q8h for 1 day	Cefoperazone 1g IV q12h for 1 day OR Ceftazidime 2g q8h IV for 1 day (if urine grew <i>Pseudomonas</i>)	
Clean-contaminated (with use of bowel segments) e.g. cystectomy with urinary diversion, cystoplasty. Common organisms: <i>Escherichia coli</i> , <i>Klebsiella</i> , <i>Proteus</i> , <i>Enterococcus</i> , <i>Pseudomonas</i> , anaerobes	Cefoperazone 1g IV q12h PLUS Metronidazole 500mg IV q8h	Gentamicin 1.5mg/kg IV q8h PLUS Metronidazole 500mg IV q8h	For duration of catheter presence.
Implant of prosthetic devices e.g. insertion of penile prosthesis or artificial urinary sphincter, artificial slings Common organism: <i>Staphylococcus aureus</i>	Cefuroxime 1.5g IV q8h for 1 week	Amoxicillin/clavulanate 1.2g IV q8h for 1 week OR Ampicillin/sulbactam 3g IV q8h for 1 week	
Laparoscopic surgery or clean– contaminated	As for open surgery	As for open surgery	Depending on type of procedure performed whether clean
Reference: 1.Pickard R., Bartoletti R., Bjerklund-Johansen TE., Bonkat G., Bruyere F., Cek M. et al. members of the EAU – ESTRO – ESUR –SIOG Urological Infections Guidelines Panel. EAU – ESTRO – ESUR – SIOG Guidelines on Urological Infections. Edn. presented at the EAU Annual Congress London 2017. 978-90-79754-91-5. Publisher: EAU Guidelines Office. Place published: Arnhem, The Netherlands.			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
9. CARDIAC SURGERY			
Coronary artery bypass	Cefazolin 2g IV (3g IV for patients weighing ≥120 kg)	Cefuroxime 1.5g IV	
Cardiac device insertion procedures (e.g. pacemaker implantation)	Cefazolin 2g IV (3g IV for patients weighing ≥120 kg)	Cefuroxime 1.5g IV	
Reference: 1. Bratzler DW, Dellinger EP, Olsen KM et al. Clinical practice guidelines for antimicrobial prophylaxis in surgery. Am J Health-Syst Pharm.2013; 70:195-283			
10. HEPATOBILIARY SURGERY			
Laparoscopic procedures Low risk	Cefazolin 2g (3g IV for patients weighing ≥120 kg)		Optimum antibiotic timing is to complete intravenous infusion ≤60min (optimal window 15 to 45 min) prior to skin incision; to ensure adequate time to reach bactericidal serum and tissue concentration before skin is incised. Repeat intraoperative dosing is recommended in: • Prolonged surgery > 4 hours • Massive blood loss > 1.5 L *Aminoglycosides should not be re-dosed.
Laparoscopic procedures High risk: Stent insertion Biliary obstruction (high direct bilirubin)	Cefazolin 2g (3g IV for patients weighing ≥120 kg) PLUS *Gentamicin 5mg/kg IV (2mg/kg IV single dose if CrCl< 20)		
Open surgery Low risk	Cefazolin 2g IV (3g IV for patients weighing ≥120 kg)		
Open surgery High risk	Cefazolin 2g IV (3g IV for patients weighing ≥120 kg)		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Multiple ERCP (≥ 2) done with stenting Biliary obstruction Biliary infection or surgery within < 30 days	PLUS Gentamicin 5mg/kg IV (2mg/kg IV single dose if CrCl< 20) If high risk ESBL/multi-resistant organisms, e.g. ESBL in the last 3 months but treated Piperacillin/tazobactam 4.5g IV PLUS Gentamicin 5mg/kg IV (2mg/kg IV single dose if CrCl< 20)		
Pre-existing infection before surgery, gallbladder empyema, ascending cholangitis	Initiate antibiotics according to culture results, or refer to treatment guidelines. If patient is at risk of infection with multidrug resistant organism, to discuss with consultant surgeons/ID physicians		
Reference: 1. Bratzler DW, Dellinger EP, Olsen KM et al. Clinical practice guidelines for antimicrobial prophylaxis in surgery. Am J Health-Syst Pharm.2013; 70:195-283			
11. NEUROLOGICAL SURGERY			
Clean wounds (Uninfected operative wounds in which no inflammation is encountered and no viscus is entered during the procedures) - Elective craniotomy or spinal procedures	Cefuroxime 1.5g IV (Given as a single IV dose at induction or within 60 minutes before incision. For prolonged procedures, additional intraoperative doses are given at every 4 hours interval during surgery in patients with normal renal function)	Vancomycin 15-20mg/kg IV (max 2g) (Infusion is started within 60-120 minutes before incision. Additional redose interval is at every 12 hours during surgery in patients with normal renal function)	Situations where the use of vancomycin is appropriate:- ● In hospitals in which MRSA or <i>S. epidermidis</i> is the frequent cause of postoperative wound infection, ● In patients previously colonised with MRSA, or ● Those who are allergic to penicillins or cephalosporins.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			Rapid IV administration of vancomycin may cause hypotension.
Clean wounds with foreign body or instrumentation - CSF shunting procedures, implantation of cranial or spinal implants	Cefuroxime 1.5g IV PLUS Metronidazole 500mg IV (Given as a single IV dose at induction or within 60 minutes before incision. Additional redose interval is at every 4 hours during surgery in patients with normal renal function)	Vancomycin 15-20mg/kg IV (max 2g) PLUS Gentamicin 5mg/kg IV (Given as a single IV dose at induction or within 60 minutes before incision in patients with normal renal function) PLUS Metronidazole 500mg IV	Addition of another drug such as metronidazole and aminoglycoside is appropriate for procedures in which enteric gram-negative bacilli or anaerobes are common pathogens.
Clean-contaminated wounds (Operative wounds in which a viscus is entered and without unusual contaminations) - Procedures that breach air cells or nasal or oral cavity	Cefuroxime 1.5g IV PLUS Metronidazole 500mg IV	Vancomycin 15-20mg/kg IV (max 2g) PLUS Gentamicin 5mg/kg IV PLUS Metronidazole 500mg IV	
Contaminated wounds (Open, fresh accidental wounds, operation with major breaks in sterile technique, or gross spillage from a viscus)	Ceftriaxone 2g IV (Given as a single IV dose at induction or within 60 minutes before incision. Additional redose interval is at every 12 hours during surgery in patients with normal renal function) PLUS Metronidazole 500mg IV	Vancomycin 15-20mg/kg IV (max 2g) PLUS Gentamicin 5mg/kg IV PLUS Metronidazole 500mg IV	

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Dirty wounds (Infected CSF shunt, old traumatic wounds with retained devitalized tissue, foreign bodies or wounds that involve existing clinical infection or perforated viscus)	Ceftriaxone 2g IV PLUS Metronidazole 500mg IV	Vancomycin 15-20mg/kg IV (max 2g) PLUS Gentamicin 5mg/kg IV PLUS Metronidazole 500mg IV	Settings where intraventricular antibiotics (Vancomycin 10mg or Gentamicin 5mg) may be useful: ● Failure to sterilise the CSF with IV therapy ● Poor response to systemic IV antibiotics ● Presence of highly resistant organisms susceptible to only antibiotics with poor CSF penetration ● Circumstances in which shunt devices cannot be removed (including infected Ommaya reservoir)
References: 1. Salford Royal, NHS. Antibiotic Prophylaxis in Cranial Neurosurgery Antibiotic Guidelines, Unique ID: 144TD(C)25(F4) Issue number: 6, 2018 2. SIGN 104 Antibiotic prophylaxis in surgery. July 2008, updated April 2014 3. Surgical Antimicrobial Prophylaxis Clinical Guideline v2.0. Department for Health and Ageing, Government of South Australia. October 2017			
12. OPHTHALMOLOGY			
- The use of povidone iodine 10% to the periorbital skin and 5% to the conjunctival sac as an antiseptic agent for preoperative surgical site preparations are recommended. - Intracameral injection of 1mg Cefuroxime in 0.1ml at the end of cataract surgery is recommended. Careful dilution should be undertaken to prevent potential toxicity. - Topical antibiotics at the end of surgery.			
Reference: 1. Prophylaxis for intraocular surgery-CPG for Management of Post-Operative Endophthalmitis, Ministry of Health Malaysia, August 2006			

B. NON-SURGICAL CHEMOPROPHYLAXIS

Patients with cardiac conditions listed in Table 1 are considered as being at increased risk of developing infective endocarditis (IE) and are indicated for antibiotic prophylaxis prior to certain procedures.

Table 1: Cardiac conditions with increased risk of IE
• Prosthetic cardiac valves or prosthetic material used for cardiac valve repair • Established rheumatic heart disease • Previous history of IE • Unrepaired cyanotic congenital heart disease (CHD), including palliative shunts and conduits • Completely repaired CHD with prosthetic material or devices, for first 6 months after the procedure • Repaired CHD with residual defects at the site or adjacent to the site of the prosthetic device (which inhibit endothelialisation) • Cardiac transplantation recipients who develop cardiac valvulopathy

Dental procedures

For patients considered as high risk (Table 1), antibiotic prophylaxis is recommended for invasive dental procedures involving manipulation of gingival tissue or the periapical region of the teeth or perforation of the gingival mucosa.

Even with high cardiac risk of IE, antibiotic prophylaxis is not recommended for:

- local anaesthetic injections in non-infected tissues
- treatment of superficial caries
- removal of sutures
- dental X-rays
- placement or adjustment of removable prosthodontic or orthodontic appliances or braces
- following the shedding of deciduous teeth
- trauma to the lips and oral mucosa

Non-dental procedures

Table 2: Non-dental procedures and recommendations for antibiotic prophylaxis in patients with increased risk of IE		
Respiratory tract procedures	Gastrointestinal and genitourinary procedures	Dermatological or musculoskeletal procedures
• Antimicrobial prophylaxis is recommended for patients with increased risk of IE (Table 1) who undergo an invasive respiratory tract procedure that involve incision or biopsy of the respiratory mucosa. • Patients who undergo an invasive respiratory tract procedure to treat an established infection, e.g. biopsy drainage of an abscess, should receive an antibiotic prophylaxis which contains an anti-staphylococcal agent.	• Routine pre-procedural antibiotic prophylaxis is no longer recommended for patients undergoing genitourinary or gastrointestinal tract procedures. • However, for patients with increased risk of IE (Table 1) who have an established gastrointestinal or genitourinary infection, or for those who receive antibiotic therapy for surgical reasons, the antibiotic regimen should include an agent active against enterococci, such as ampicillin or vancomycin.	• For patients described in Table 1 undergoing surgical procedures involving infected skin (including local abscesses), skin structure or musculoskeletal tissue, it is reasonable that the therapeutic regimen contains an agent active against staphylococci and beta-hemolytic streptococci. Vancomycin or clindamycin may be used in patients unable to tolerate a β-lactam antibiotic. • If the infection is known or suspected to be caused by MRSA, vancomycin or another suitable agent should be administered.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Prophylactic regimens for high-risk dental procedures in high-risk patients			
Prophylactic regimen	Amoxicillin 2g PO OR Ampicillin 2g IV	Cefazolin 1g IV Penicillin allergy: Clindamycin 600mg IV/PO	See above for antibiotic prophylaxis in patients undergoing invasive surgical procedure to treat an established infection. Antibiotic to be administered as a single dose 30 to 60 minutes before procedure.
Secondary prevention of rheumatic fever			
Secondary prevention of rheumatic fever	Benzathine benzylpenicillin 1.2MU IM every 3 to 4 weeks OR Phenoxymethylpenicillin (Penicillin V) 250mg PO q12h	Penicillin allergy: Erythromycin Ethylsuccinate 800mg PO q12h	<i>Duration of prophylaxis please refer to the table below</i>

Type of Infection	Duration of prophylaxis
Rheumatic fever with carditis and residual heart disease (persistent valvular disease)	10 years or until age 40 years, whichever is longer, sometimes lifelong prophylaxis
Rheumatic fever with carditis but no residual heart disease (no valvular disease)	10 years or until age 21 years, whichever is longer
Rheumatic fever without carditis	5 years or until age 21 years, whichever is longer

References:

1. Ministry of Health Malaysia's Clinical Practice Guidelines For The Prevention, Diagnosis & Management Of Infective Endocarditis 2017
2. ESC Guidelines on Prevention of Infective Endocarditis 2015
3. The 2020 Australian Guideline for Prevention, Diagnosis and Management of Acute Rheumatic Fever and Rheumatic Heart Disease (3.2 Edition, March 2022)

(SECTION B: PAEDIATRICS)

Doses of recommended antimicrobials are not included in this section.

Please refer to relevant references for dosage recommendations

CARDIOVASCULAR INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
ACUTE MYOCARDITIS			
Viral (commonest cause) Enteroviruses Adenovirus Influenza HIV etc	Treatment mainly supportive		Among the viruses implicated are enteroviruses including Coxsackie & EV71. For severe HFMD with cardiopulmonary failure stage, use of IVIG may be considered if not used during CNS involvement or autonomic nervous system dysregulation stage.
ACUTE PERICARDITIS			
Viral (commonest cause) Bacterial: <i>Staphylococcus aureus</i> <i>Haemophilus influenza</i> <i>Salmonella</i> spp. <i>M.Tuberculosis</i>	Treatment mainly supportive Empiric for purulent pericarditis: IV Cloxacillin PLUS IV Cefotaxime	<u>Antibiotic allergy:</u> IV Cefazolin	Need pericardial fluid to differentiate between different etiologic agents and C&S to adjust antibiotics. Consider surgical drainage for tamponade, pre-tamponade & ineffective conservative management. Duration of therapy: 4 weeks
INFECTIVE ENDOCARDITIS			
Empirical Therapy for Infective Endocarditis			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Community-acquired organisms: <i>Streptococcus</i> , <i>Enterococcus</i> HACEK Gram-negative organisms	IV Ampicillin PLUS IV Gentamicin	PLUS/MINUS *IV Cloxacillin	*For acute presentation, need to cover for MSSA since <i>Streptococcus</i> & <i>Enterococcus</i> HACEK presentations are usually sub-acute.
Healthcare-associated organisms: MRSA Non-HACEK Gram-negative organisms <i>Enterococcus sp.</i>	IV Vancomycin PLUS IV Gentamicin PLUS/MINUS *Rifampicin		*Rifampicin is ONLY for prosthetic valve AND added 3-5 days later than vancomycin & gentamicin If non-HACEK Gram-negative organisms like pseudomonas are suspected epidemiologically, add cefepime until cultures are known. Once cultures are available, adjust accordingly.
Specific Organisms:			
1. Infective endocarditis (<i>Streptococcus viridans</i>)			
Strains fully susceptible to penicillin (MIC<0.125 mg/L)	IV Benzylpenicillin	IV Ampicillin OR IV Ceftriaxone OR <u>Antibiotic allergy</u> *IV Vancomycin	Duration: -4 weeks for native valve -6 weeks for prosthetic valve Vancomycin dose adjusted for trough concentration of 10-15mg/ml
Strains with MIC > 0.125 to 2mg/L	IV Benzylpenicillin PLUS		*Vancomycin therapy is recommended only for patients

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	IV Gentamicin for 2 weeks (add to first line regimen of penicillin/ceftriaxone) Do not use ampicillin		with immediate type penicillin hypersensitivity. For this strain (MIC>0.125): Antibiotics of choice are either penicillin with gentamicin or ceftriaxone with gentamicin.
2. Infective Endocarditis (<i>Enterococcus</i> sp.)			
Penicillin-sensitive (MIC ≤ 8mg/L)	IV Ampicillin for *4-6 weeks PLUS IV Gentamicin for *2-6 weeks	IV Ampicillin PLUS IV Ceftriaxone	*Duration: -If symptoms less than 3 months & native valve: ampicillin for 4 weeks & gentamicin for 2 weeks -If symptoms more than 3 months: ampicillin & gentamicin for 6 weeks Ampicillin plus ceftriaxone is preferred for individuals with renal impairment (CrCl ≤ 50ml/min) ONLY. Cannot use ceftriaxone alone since enterococcus is intrinsically resistant to this drug. This combination is NOT ACTIVE against <i>E. faecium</i> . **Maximum dose of vancomycin: 2g/day unless not able to achieve therapeutic range. Aim for serum trough of 10-20mg/L.
Sensitive to penicillin & vancomycin but high-level resistance to gentamicin. (MIC > 500mg/L)	IV Ampicillin PLUS IV Ceftriaxone Duration : 6 weeks		
Resistant to penicillin but susceptible to vancomycin & gentamicin	**IV Vancomycin PLUS IV Gentamicin Duration : 6 weeks		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
3. Infective Endocarditis (<i>Staphylococcus aureus</i>)			
Methicillin-sensitive (left-sided)	IV Cloxacillin for 4-6 weeks	<u>Antibiotic allergy</u> IV Cefazolin for 4-6 weeks OR IV Vancomycin for 4-6 weeks.	If allergy to penicillin but not immediate type hypersensitivity, use cefazolin. Methicillin-sensitive (right sided): can shorten duration to 2 weeks if good response, no metastatic sites, no cardiac or extracardiac complications with size of vegetation less than 20mm.
Methicillin-sensitive (right-sided)	IV Cloxacillin for 4 weeks		
Methicillin-resistant (left & right)	IV Vancomycin for 4-6 weeks	IV Daptomycin for 4-6 weeks	Daptomycin is superior to vancomycin for MRSA bacteremia with MIC >1mg/L.
Methicillin-sensitive (prosthetic valve)	IV Cloxacillin for ≥6 weeks PLUS IV Gentamicin for 2 weeks PLUS *PO Rifampicin ≥6 weeks		*Rifampicin has better penetration but to protect against development of resistance, use only after 3-5 days of cloxacillin and/or bacteremia has been cleared. MRSA (prosthetic valve): Vancomycin & rifampicin for 6 weeks or more.
Methicillin-resistant (prosthetic valve)	IV Vancomycin ≥6 weeks PLUS IV Gentamicin for 2 weeks PLUS *PO Rifampicin ≥6 weeks		
4. Culture-negative endocarditis	IV Ampicillin/sulbactam for 4-6 weeks PLUS IV Gentamicin for 4-6 weeks		Culture-negative endocarditis (CNE) is diagnosed when a child has clinical

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			& echocardiogram evidence of IE but persistent negative cultures. This is in individuals with no prior antimicrobial use. If fungi or fastidious organisms are suspected, need to ask microbiologist to prolong incubation. Patients with culture-negative endocarditis should be treated in consultation with an ID specialist.
References: 1. American Academy of Paediatrics. Committee on Infectious Diseases. Red Book: Report of the Committee on Infectious Diseases (2018) 2. Malaysia Clinical Practice Guideline for the Prevention, Diagnosis and Management of Infective Endocarditis 2017			

CENTRAL NERVOUS INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Meningitis empirical treatment			
Age group: 1-3 months: <i>Group B streptococcus (GBS), E. Coli, S. pneumoniae & N. meningitidis</i> > 3months: <i>S. pneumoniae, Hib, E. coli, Salmonellosis, N. meningitidis</i>	<u>Birth - 3 months old:</u> IV Benzylpenicillin PLUS IV Cefotaxime <u>More than 3 months old:</u> IV Benzylpenicillin PLUS IV Cefotaxime OR IV Ceftriaxone		For children below 3 months of age: Cefotaxime is the preferred third generation cephalosporin since less drug interaction (in terms of interaction with calcium containing infusion and bilirubin displacement) Once organism is known, please refer below to adjust antibiotics
Specific organisms			
<i>Haemophilus influenzae</i>	IV Cefotaxime OR IV Ceftriaxone	IV Ampicillin	Duration 10 days
<i>Neisseria meningitidis</i>	IV Benzylpenicillin	IV Cefotaxime OR IV Ceftriaxone	Duration: 7 days Prophylaxis for all household contacts and healthcare workers involved in unprotected contact during intubation and suctioning of airway or mouth-to-mouth resuscitation.
<i>Streptococcus pneumoniae</i>			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Penicillin-susceptible (MIC≤0.06mg/L)	IV Benzylpenicillin		Duration: 14 days
Penicillin-resistant (MIC ≥0.12mg/L) & cefotaxime-/ceftriaxone-sensitive (MIC ≤0.5mg/mL)	IV Cefotaxime OR IV Ceftriaxone		
Penicillin & cefotaxime-/ceftriaxone - resistant (MIC ≥2.0mcg/ml) (drug-resistant <i>Streptococcus pneumoniae</i> , DRSP)	High dose IV Cefotaxime OR High dose IV Ceftriaxone PLUS IV Vancomycin		Treat in consultation with ID specialist
Cryptococcal meningitis <i>Cryptococcus neoformans</i>	Induction therapy: IV Amphotericin B PLUS/MINUS Oral 5-flucytosine (5-FU) for 2 to 4 weeks Consolidation Therapy: IV/PO Fluconazole for 8 weeks		Duration of induction with 5-FU at least two weeks and until CSF repeat culture is NEGATIVE.
Herpes Simplex encephalitis	4 months to 12 years old IV Acyclovir		Duration: 14 to 21 days (treat in consultation with ID specialist)
Brain abscess	IV Cefotaxime OR IV Ceftriaxone PLUS IV Metronidazole	If secondary to trauma: PLUS IV Cloxacillin (add to 3 rd -generation cephalosporin)	Surgical drainage may be indicated if appropriate. Duration 6-8 weeks, depending on response as seen from neuroimaging & clinical presentation.
References: - Academy of Medicine of Malaysia Clinical Practice Guidelines on Rational Antibiotic Utilisation in Selected Paediatric Conditions April 2004 - The Sanford Guide to Antimicrobial therapy 2018			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
3. Malaysia Paediatric Protocol, 4 th ed. 2018, Kementerian Kesihatan Malaysia			

GASTROINTESTINAL INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Acute gastroenteritis Usually viruses e.g.:rotavirus	Antibiotics not recommended		Oral rehydration is the cornerstone of treatment. Antibiotic therapy may prolong carriage state of salmonellosis.
Dysentery <i>Shigella, E. coli, Campylobacter</i> Mild or uncomplicated Severe illness (hospitalisation, invasive or other complications) or immunocompromised patients	Most are mild infections which resolved spontaneously without antibiotics No treatment required Empiric: IV Ceftriaxone for 5 days (origin of infections: Asia)	*PO Ampicillin for 5-7 days for hospitalized children IV Azithromycin for 3 days	Resistance patterns towards amoxicillin, trimethoprim sulphamethazole, ciprofloxacin & azithromycin are on the rise. Adjust antibiotic once culture & sensitivity (C&S) results are available. For immunocompromised host-treat longer (7-10 days). Reserve fluoroquinolone only for isolate where there is no other antibiotic option available due to its many side effects. *PO Ampicillin not available in HRPB Formulary.
Dysentery <i>Amoebiasis</i>	PO Metronidazole for 7-10 days		Similar dosage for extraintestinal disease
Giardiasis	PO Metronidazole for 5-7 days		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Typhoid fever <i>Salmonella</i> Typhi <i>S. Paratyphi</i> A&B Mild or uncomplicated Severe infection or suspected resistant organism Chronic carrier state(>1year)	Empirical treatment: IV Ceftriaxone for 7-14 days PO Ciprofloxacin for 5-7 days IV Ceftriaxone for 7-14 days *PO Ampicillin for 6 weeks OR PO Amoxicillin for 6 weeks OR PO Trimethoprim/sulfamethoxazole for 6 weeks	PO Chloramphenicol for minimum 14 days PO Ciprofloxacin for 7-10 days PO Ciprofloxacin for 4 weeks OR IV Ampicillin (If oral therapy not tolerated & strain is susceptible)	Adjust antibiotic once C&S results are known. Duration of antibiotics: 7 days (uncomplicated) to 14 days (severe disease or if using ampicillin or trimethoprim/sulfamethoxazole). Choice of antibiotics & duration depends on disease, C&S results & whether oral route is preferred. Fluoroquinolones need to be used with caution in children due to possible arthropathy & rapid development of resistance. There is now increasing data of other side effects e.g hypoglycaemia & neuropsychiatric disorder. Ampicillin & trimethoprim/sulfamethoxazole may be considered for susceptible strain. More strains are now becoming sensitive to these agents except for certain countries. *PO Ampicillin not available in HRPB Formulary
Cholera	PO Azithromycin in a single dose OR		Oral or IV rehydration is the cornerstone of treatment. Prompt

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	PO Erythromycin Ethylsuccinate for 3 days OR PO Doxycycline (children>8 years old) single dose		initiation of antibiotic therapy reduces the volume & duration of diarrhea. Antimicrobials should be considered for people who are moderately to severely ill. Choice dependent on age & pattern of resistance. Monitor antimicrobial sensitivity pattern at beginning of & during the outbreak as it can change Avoid using tetracycline or doxycycline for young children as they can cause staining of the teeth. Use of doxycycline should be considered in an epidemic caused by susceptible isolates. Fluoroquinolones are not approved for children younger than 18 years old for this indication.
Liver abscess (amoebic) <i>Entamoeba Histolytica</i>	PO Metronidazole for 7-10 days		Amoebic abscess tends to be solitary lesion. Consider surgical drainage if needed.
Liver abscess (pyogenic) <i>Klebsiella spp.</i> , <i>E.coli</i> , <i>Streptococcus milleri</i> , other Gram-negative	IV Cefotaxime OR IV Ceftriaxone PLUS	IV Piperacillin/tazobactam <u>ESBL-Klebsiella:</u> IV Ertapenem (above 3 months of age)	Surgical drainage is needed in most cases Duration: 4-6 weeks

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
organisms, anaerobes, <i>S. aureus</i>	IV Metronidazole		
Acute cholangitis Gram-positive & Gram-negative organisms, anaerobes	IV Cefotaxime OR IV Ceftriaxone PLUS IV Metronidazole	IV Piperacillin/tazobactam OR IV Ampicillin/sulbactam	Duration: 7 days. Outcome is similar with less than 7 days to those with longer duration > 7 days in patients treated with percutaneous cholecystectomy. In treatment failure, source control is needed.
Peritonitis Gram positive & Gram-negative organism, anaerobes	Primary/spontaneous bacterial peritonitis IV Cefotaxime Secondary (nosocomial) peritonitis IV Piperacillin/tazobactam If culture proven ESBL: IV Imipenem/cilastatin OR IV Meropenem De-escalate treatment to IV ertapenem once patient is stable	*PO Ampicillin PLUS IV Gentamicin PLUS IV Metronidazole for 7-14 days	May omit metronidazole in primary peritonitis. In immunocompetent patients with mild to moderate peritonitis & source control, suggest 5 days of therapy. Ertapenem is not licenced to be used in infants less than 3 months old *PO Ampicillin not available in HRPB Formulary (Alternative: PO Unasyn)
References: 1. CDC. Antibiotics Resistance Threats in the United States 2013. 2. Dr. Phoebe Williams, Prof James A Berkeley. Dysentery (Shigellosis) Current WHO Guidelines and the WHO Essential Medicine List for Children. November 2016. 3. Frank Shann, Seventeenth edition, 2017. 4. WHO/V&B/03-07 (2003) Background document: the diagnosis, treatment and prevention of typhoid fever. 5. WHO/CDD/SER/91.15 REV.1. Management of Patient with Cholera. 6. Christoph Lübbert, Johannes Wiegand, Thomas Karlas. Viszeralmedizin. 2014 Oct; 30(5): 334–341. 7. Patrick Mosler. Management of Acute Cholangitis. Gastroenterol Hepato (NY) 2011 Feb; 7(2): 121–123.			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
8. Solomkin JS, Mazuski JE, Bradley JS, et al.; Surgical Infection Society; Infectious Diseases Society of America. Diagnosis and management of complicated intra-abdominal infection in adults and children: guidelines by the Surgical Infection Society and the Infectious Diseases Society of America. Clin Infect Dis. 2010;50(2):135.			
9. American Academy of Paediatrics. Committee on Infectious Diseases. Red Book: Report of the Committee on Infectious Diseases (2018).			

INFECTIONS IN IMMUNOCOMPROMISED PATIENTS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Febrile Neutropenia First line: Fever >38°C, neutrophil<500mm3 <i>Enterobacterales (Klebsiella sp., E. coli etc.), Pseudomonas, aerobic Gram-positive (Staphylococci, Streptococci)</i>	IV Piperacillin/tazobactam PLUS IV Gentamicin	IV Cefepime	Need to look at local epidemiological data.
Second line: Persistent fever > 72 hours <i>Enterobacterales (Klebsiella sp, E. coli etc.), Pseudomonas, aerobic Gram-positive (Staphylococci, Streptococci), Enterococci</i> or other resistant organisms *DO NOT MODIFY INITIAL COVERAGE BASED SOLELY ON PERSISTENCE OF FEVER	IV Meropenem PLUS/MINUS IV Vancomycin		Escalate to second line if patient unstable to cover resistant Gram-negative, Gram-positive & anaerobes. Consider adding vancomycin in suspected catheter-related infections, positive blood culture for Gram-positive cocci, hypotensive patients & patients who are known to be colonised with MRSA. In patients responding to initial empiric antibiotic therapy, discontinue double coverage (empirical vancomycin, if initiated) or double gram negative after 24-72 hours if there is no specific microbiologic

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			indication to continue combination therapy.
Third line: Fever > 4-7 days with no identified source of fever <i>Candida</i> sp. <i>Aspergillus</i> sp., <i>Fusarium</i> sp. Viral: Respiratory viruses are the most common, HSV, VZV	IV Imipenem/cilastatin PLUS IV Amphotericin B OR Lipid Formulation of Amphotericin B	IV Imipenem/cilastatin PLUS IV Caspofungin	1/3 of febrile neutropenic patients with persistent fever >1 week have systemic fungal infections ² In patients at high risk of invasive fungal disease with prolonged (≥96 hours) febrile neutropenia unresponsive to broad spectrum antibacterial agents, initiate antifungal. Amphotericin-based antifungal is considered broader spectrum than echinocandin (e.g. Caspofungin).
References: 1. β lactam monotherapy versus β lactam-aminoglycoside combination therapy for fever with neutropenia: systematic review and meta-analysis. BMJ 2003; 326:1111. 2. Clinical practice guideline for the use of antimicrobial agents in neutropenic patients with cancer: 2010 update by the Infectious Diseases Society of America. Freifeld AG, Bow EJ, Sepkowitz KA, Boeckh MJ, Ito JI, Mullen CA, Raad II, Rolston KV, Young JA, Wingard JR, Clin Infect Dis. 2011;52(4):e56. 3. Guideline for the management of fever and neutropenia in children with cancer and/or undergoing hematopoietic stem-cell transplantation recipients. Lehrnbecher T, Robinsn P, Fisher B,, Alexander S, Ammann RA, , Beauchemin M, Carlesse F, Groll AH, Haeusler GM, Santolaya M, Steinbach WJ, Castagnola E, Davis BL, Dupuis LL, Gaur AH, Tissing WJE, Zotis T, Phillips R, Sung L,J Clin Oncol. 2017;35(18):2082-2094. 4. Empiric treatment against invasive fungal diseases in febrile neutropenic patients: a systematic review and network meta-analysis. Ken Chen, Qi Wang, Roy A. Pleasants, Long Ge. Wei Liu, Kanging Peng and Suodi Zhai. BMC Infectious Diseases. 2017; 17;159. 5. A Randomized, Double-Blind, Multicenter Study of Caspofungin versus Liposomal Amphotericin B for Empirical Antifungal Therapy in Pediatric Patients with Persistent Fever and Neutropenia. Maertens JA, Maedro L, Reilly AF, Lehrnbecher T, Groll AH, Jafri HS, Green M, Nania JJ, Kartsonis NA, Chow JW, Arndt CAS, DePauw BE, Walsh T. Pediatr Infect Dis J. 2010; 29:415-420.			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
6. Lehrnbecher et al. Guideline for Management of fever and neutropenia in children with cancer and hemapoietic stem cell transplantation recipients-2017 update. J Clin Oncology 2017 35:18, 2082- 2094.			

NEONATAL INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Congenital & Perinatal Infections			
Meningitis GBS <i>E. coli</i> Listeria other Gram-negative bacilli/rod (GNR)	Empirical therapy: IV Benzylpenicillin OR IV Ampicillin PLUS IV Cefotaxime		Once cultures are known, adjust antibiotics accordingly.
Necrotising enterocolitis (NEC) Klebsiella, <i>E. coli</i> , Clostridia, Coagulase-negative Staphylococci, Enterococci, Bacteroides	IV Ampicillin PLUS IV Gentamicin PLUS IV Metronidazole Duration: 10-14 days		There is insufficient evidence regarding the duration of antibiotic treatment for NEC. This suggested regimen for NEC is empirical. Once culture is known, decisions regarding choice of antibiotics are best guided by culture results. Use vancomycin if CoNS/MRSA is suspected (substitute ampicillin with vancomycin).
Early onset sepsis (<48 hrs) Group B Streptococcus (GBS), Listeria, Streptococcus sp., <i>E. coli</i> , <i>Haemophilus influenza</i> , <i>Klebsiella</i> sp. etc.	IV Benzylpenicillin OR IV Ampicillin PLUS IV Gentamicin		If negative blood culture, initial clinical suspicion not strong & reassuring baby’s condition with low CRP, consider stopping antibiotics at 48 hours. If positive blood culture or strong clinical suspicion of sepsis but negative culture, may give 5-7 days of antibiotics.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			Consider antibiotics for more than 5-7 days if baby not fully recovered & based on pathogen identified on blood culture. In this empiric therapy, meningitis is not a consideration. Once cultures are known; adjust antibiotics accordingly.
Late onset sepsis >48 hours Methicillin-sensitive/ -resistant <i>S. aureus</i> (MSSA/MRSA), Coagulase-negative staphylococci (CONS), Gram-negative rods (dependent on local epidemiological data)	First line: IV Cloxacillin PLUS IV Gentamicin		For late onset sepsis, the most common organisms are predominantly Gram-positive cocci, namely staphylococci, especially CONS, in premature neonates & also neonates with central catheters. Piperacillin/tazobactam is a good second line option in pneumonia & intra- abdominal sepsis (non-CONS sepsis with good coverage against Gram-positive, Gram-negative & anaerobes). There is possibility of Gram-negative rods with inducible β-lactamases & ESBL-producing organisms such as <i>Klebsiella</i>, <i>Serratia</i> & <i>E. coli</i> in some NICU in Malaysia hence need to look at local epidemiology before deciding on suitable second line.
	Second line: IV Piperacillin/tazobactam		
	Other options: IV Cefepime OR IV Meropenem OR IV Imipenem/cilastatin		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			Cefepime is the preferred agent when there are Gram-negative bacteria with extended spectrum cephalosporin resistance due to AmpC-β-lactamases (also termed Class C or Group 1). Studies comparing imipenem/cilastatin & meropenem for indication other than meningitis found no significant differences in efficacy or safety between the two. However, a recent meta-analysis suggested that meropenem may be better than imipenem/cilastatin in efficacy & safety especially due to more cases of seizure with imipenem (WHO Subcommittee Meeting of the Expert Committee on the Selection & Use of Essential Medicine 2008).
Congenital syphilis <i>T. pallidum</i>	IV Benzylpenicillin (Penicillin G) Duration : 10 days <u>If diagnosed with congenital syphilis after one month of age:</u> IV Benzylpenicillin (Penicillin G) for 10-14 days. In infants considered less likely to have syphilis & normal CSF examination including	*IM Procaine penicillin 10 days.	Only severe cases are clinically apparent at birth. Refer to algorithm for diagnosing & evaluation. Re-evaluate & possibly re-treat. Please refer Red Book 2018. *Procaine penicillin not available in HRPB Formulary

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	normal physical examination & long bone radiograph: Benzathine penicillin IM in a single dose can be given		
Congenital toxoplasmosis <i>T. gondii</i>	*Pyrimethamine PLUS *PO Sulfadoxine PLUS PO Folinic Acid for 12 months [*PO Pyrimethamine/sulfadoxine (Fansidar®)]	*PO Pyrimethamine PLUS *PO Sulfadiazine PLUS PO Folinic Acid	Duration: For 12 months Drug regimen is not definitively established. Clinical trials are ongoing. Prednisolone q12h can be added if CSF protein $\geq 1\text{g/dL}$ or active severe chorioretinitis. Steroids given till CSF protein $< 1\text{g/dL}$ or resolution of severe chorioretinitis. Fansidar is currently an “orphan” drug that need special procurement measures to buy. Refer to paediatric ID consultant for treatment and availability of drug. * Requires special medicine approval.
Herpes simplex • Localised skin, eye & mouth (SEM) • Central nervous system (CNS) with or without SEM • Disseminated disease involving multiple organs	IV Acyclovir Duration: -Skin, eyes, mouth: 14 days -CNS/disseminated: minimum of 21 days All infants surviving neonatal HSV infection of any classification should receive oral acyclovir suppression administered 3 times		Screen for other STDs. For CNS disease: Repeat lumbar puncture at end of therapy for HSV PCR. If PCR remains positive, continue IV acyclovir for another one week. Recurrence of HSV can occur & may be a lifelong problem.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	daily for 6 months after completion of parenteral therapy.		
Tetanus neonatorum	IV Metronidazole	IV Benzylpenicillin	Duration: 10 days
Congenital gonococcal ophthalmitis/conjunctivitis	Immediate & frequent saline eye irrigation. Non-disseminated disease: IV Cefotaxime in a single dose May need to continue for 48-72 hours until systemic infection has been ruled out. Disseminated disease: IV Cefotaxime For 7 days, with a duration of 10-14 days, if meningitis is documented.	If penicillin-sensitive, may give IV Benzylpenicillin	Evaluate for signs of disseminated infection (e.g. sepsis, arthritis & meningitis). Screen mother & baby for chlamydial infection. Screen for other STDs. Investigate & treat parents.
<i>Chlamydia trachomatis</i> conjunctivitis	PO Erythromycin Ethylsuccinate Duration : 14 days Local eye toilet until discharge stops.	PO Azithromycin Duration: 3 days	Initial treatment for chlamydial conjunctivitis should be based upon a positive diagnostic test. Re-swab after treatment; 20-30% will need a second course to clear infection.
GBS <i>Streptococcus agalactiae</i>			
Sepsis	IV Benzylpenicillin OR IV Ampicillin PLUS IV Gentamicin		Duration of treatment for GBS: -Uncomplicated: 14 days (bacteremia without a defined focus). -Meningitis: 21 days.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			Gentamicin can be discontinued once the infection is under control.
Meningitis	IV Benzylpenicillin OR IV Ampicillin PLUS IV Gentamicin		Duration for treatment: 21 days. Doses of penicillin for meningitis is higher as recommended by experts (as high as 500,000 unit/kg/day (> 7 days of age)).
<i>E. coli</i> Sepsis/Meningitis	IV Cefotaxime PLUS IV Gentamicin		Duration in bacteremia: 14 days. Duration for meningitis: 21 days. All cases of bacteremia need lumbar puncture to exclude meningitis. Treatment duration of 14 days can be decided on a case-by-case basis if meningitis excluded & good clinical response.
References: 1. Congenital syphilis. 2015 Treatment Guidelines. Available at https://www.cdc.gov/std/tg2015/congenital.htm. 2. Yvonne A. Maldonado, MD, FAAP, Jennifer S. Read. Diagnosis, Treatment, and Prevention of Congenital Toxoplasmosis in the United States. PEDIATRICS. Volume 139, number 2, February 2017:e 20163860. 3. Swetha G. Pinninti, David W. Kimberlin. Neonatal Herpes Simplex Virus Infections. Seminars in Perinatology 42(2018) 168-175. 4. Christina W. Obiero, Anna C. Seale, James E. Berkley. The Pediatric Infectious Disease Journal • Volume 34, Number 6, June 2015. 5. American Academy of Paediatrics. Committee on Infectious Diseases. Red Book: Report of the Committee on Infectious Diseases (2018). 6. The Sanford Guide to Antimicrobial therapy 2018. 7. IBM Micromedex Neofax version number v50_190403210			

ORAL/DENTAL INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
ANTIMICROBIAL USE FOR BACTERIAL INFECTIONS			
Dental Trauma			
Avulsion of permanent teeth or incisors with an open or closed apex	PO Amoxicillin OR PO Phenoxymethylpenicillin (Penicillin V)	Penicillin Allergy: PO Doxycycline	The use of systemic antibiotics is recommended after replantation of an avulsed tooth. Risk of discoloration of permanent teeth must be considered before systemic use of tetracycline or doxycycline. Tetracycline or doxycycline are generally not recommended for children under 8 years of age. Topical antibiotics are not recommended.
Luxations orLuxation injuries in the primary dentition	Systemic antibiotic use not recommended	Systemic antibiotic use not recommended	
Dental Fractures	Systemic antibiotic use not recommended	Systemic antibiotic use not recommended	
References: 1. American Academy of Pediatric Dentistry. Use of anti- biotic therapy for pediatric dental patients. The Reference Manual of Pediatric Dentistry. Chicago, Ill.: American Academy of Pediatric Dentistry; 2021:461-4 2. Fouad AF, Abbott PV, Tsilingaridis G, et al. International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: 2. Avulsion of permanent teeth. <i>Dent Traumatol.</i> 2020;36:331– 342. 3. Day P, Flores MT, O'Connell A, et al. International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: 3. Injuries in the primary dentition. <i>Dent Traumatol.</i> 2020;36:343–359.			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
4. Bourguignon C, Cohenca N, Lauridsen E, et al. International Association of Dental Traumatology guidelines for the management of traumatic dental injuries: 1. Fractures and luxations. <i>Dent Traumatol.</i> 2020;36:314–330. 5. Goel D, Goel, Gaurav K, et al. Antibiotic prescriptions in pediatric dentistry. <i>Journal of Family Medicine and Primary Car.</i> 2020; 9:2:473-480 6. Oral Health Programme Ministry of Health Malaysia. Management of Avulsed Permanent Anterior Teeth. <i>Clinical Practice Guideline 3rd Edition 2019</i> 7. American Academy of Pediatrics (AAP). In: Kimberlin DW, Brady MT, Jackson MA, Long SA, eds. <i>Red Book: 2015 Report of the Committee on Infectious Diseases.</i> 30th ed. Elk Grove Village, IL: American Academy of Pediatrics; 2015. 8. American Academy of Pediatrics (AAP). In: Kimberlin DW, Brady MT, Jackson MA, Long SA, eds. <i>Red Book: 2018 Report of the Committee on Infectious Diseases.</i> 31th ed. Itasca, IL: American Academy of Pediatrics; 2018 9. Lexicomp Online, Pediatric and Neonatal Lexi-Drugs Online. Waltham, MA: UpToDate, Inc.; 8 April, 2022. https://online.lexi.com. Accessed June 1, 2022.			

OTORHINOLARYNGOLOGY INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Tonsillitis/Pharyngitis Group A <i>Streptococcus</i>	PO Phenoxymethylpenicillin (penicillin V) OR PO Amoxicillin	PO Cephalexin OR PO Erythromycin Ethylsuccinate	Duration: 7 days
Rhinosinusitis <i>Streptococcus pneumoniae</i> <i>Haemophilus influenzae</i> <i>Moraxella catarrhalis</i> Group A <i>Streptococcus</i>	PO Amoxicillin for 10 days Second line: PO Amoxicillin/clavulanate Failing amoxicillin/clavulanate: PO Clindamycin AND PO Cefuroxime Inpatient (severe): IV Ampicillin/sulbactam	<u>Penicillin Allergy:</u> PO Clindamycin IV Ceftriaxone	The most common causes are viral infections. Acute bacterial sinusitis is suspected when child with URI presents with: 1. Persistent illness (nasal discharge or daytime cough or both for ≥ 10 days without improvement) 2. Worsening course 3. Severe onset (concurrent fever & purulent discharge for 3 days) For rhinosinusitis, most experts recommend using high dose amoxicillin.
Acute otitis media <i>Streptococcus pneumoniae</i> <i>Haemophilus influenzae</i> <i>Moraxella catarrhalis</i>	PO Amoxicillin <2 years old : 10 days 2-5 years old: 7 days >5 years old : 5 days. For clinical failure, history of using amoxicillin in the last 30 days & has concurrent purulent conjunctivitis: PO Amoxicillin/clavulanate	<u>Penicillin Allergy:</u> PO Erythromycin Ethylsuccinate OR PO Clarithromycin OR PO Azithromycin	
Acute otitis externa <i>Pseudomonas aeruginosa</i> <i>Staphylococcus aureus</i>	Mild to moderate: Topical antibiotic with/without topical steroids.		Ototoxic agents like gentamicin or neomycin should not be used in the presence of tympanostomy tubes or

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	E.g. Gentamicin 0.3% ear drop for 7 days Polymyxin B sulphate, neomycin sulphate & hydrocortisone ear drops for 7 days Ofloxacin 0.3% otic solution for 7 days Indication: for 1-12 years old		perforated tympanic membrane. Clinical response should be seen within 48 to 72 hours but full response may take up to 6 days. Non-response should prompt an evaluation for obstruction, presence of foreign body, non-adherence or an alternative diagnosis.
References: 1. Wald ER, Applegate KE, Bordley C, et al. Clinical practice guideline for the diagnosis and management of acute bacterial sinusitis in children aged 1 to 18 years. Pediatrics 2013; 132: e262. 2. Chow AW, Benninger MS, Brook I, et al. IDSA clinical practice guideline for acute bacterial rhinosinusitis in children and adults. Clin Infect Dis 2012; 54: e72. 3. Charles PS Hui: Canadian Paediatric Society. Paediatr Child Health 2013;18(2):96-98 4. The Sanford Guide to Antimicrobial therapy 2018. 5. American Academy of Paediatrics. Committee on Infectious Diseases. Red Book: Report of the committee on Infectious Diseases (2018).			

RESPIRATORY INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Community-acquired Pneumonia			
Pneumonia (outpatient) Infant (≥3 months) & children Viral infection is more common (Influenza, RSV, human metapneumovirus (hMPV), Parainfluenza, Adenovirus) Bacteria (<i>S. pneumoniae</i>, Group A <i>Streptococcus</i>, <i>S. aureus</i>, <i>H. influenza</i>)	*High-dose PO Amoxicillin for 5-7 days <u>For influenza:</u> PO Oseltamivir	PO Erythromycin Ethylsuccinate	Antibiotics are not routinely recommended since viral infection is more common. For infants & children admitted to hospital, treat as presumed bacterial unless viral origin is known. Macrolide antibiotics should be used if either mycoplasma or chlamydia pneumonia is suspected. It may be started in school-going children where disease predominates. Duration: minimum 5 days & until afebrile for 2-3 days in empiric therapy with absence of an identified specific etiology & specific therapy with known pneumonia due to pneumococcus, <i>HI</i> & <i>Moraxella catarrhalis</i>. *Experts recommend using high dose amoxicillin to overcome resistance conferred by cell wall changes of the bacteria (pneumococcus). If a child cannot tolerate high doses, the

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			standard amoxicillin dose can be used.
Pneumonia (inpatient, fully immunised)	IV Benzylpenicillin for 5-7 days	Second line/partially treated IV Cefuroxime OR IV Amoxicillin/clavulanate	Macrolide antibiotics should be used if pneumonia due to <i>Mycoplasma</i> or <i>Chlamydia</i> is suspected.
Severe Community-acquired Pneumonia			
Child not fully immunised/life-threatening	IV Cefotaxime OR IV Ceftriaxone PLUS/MINUS IV Azithromycin for 5 days	IV Cefuroxime PLUS/MINUS IV Azithromycin for 5 days	For the management of empyema, refer Surgical Infection section .
References: 1. M. Harris, J. Clark, N. Coote et al. British Thoracic Society guidelines for the management of community acquired pneumonia in children: update 2011. Thorax, vol. 66, no. 2, pp. ii1– ii23. The Management of Community-Acquired Pneumonia in Infants and Children Older Than 3 Months of Age: Clinical Practice Guidelines by the Pediatric Infectious Diseases Society and the Infectious Diseases Society of America Clinical Infectious Diseases 2011; 53(7):e25–e76 2. Daniele Dona et al. Treatment of Community-Acquired Pneumonia: Are All Countries Treating Children in the Same Way? A Literature Review. International Journal of Pediatrics Volume 2017. 3. The Sanford guide to antimicrobial therapy 2018. 4. American Academy of Paediatrics. Committee on Infectious Diseases. Red Book: Report of the committee on Infectious Diseases (2018).			

SKIN AND SOFT TISSUE INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Abscess <i>Staphylococcus aureus</i>	Mild: PO Cloxacillin for 5-7 days Severe: IV Cloxacillin for 5-7 days CA-MRSA: PO Clindamycin for 5-7 days OR PO Trimethoprim/sulfamethoxazole for 5-7 days	PO Cephalexin for 5-7 days	Incision & drainage (I&D) is the MAINSTAY of therapy. Needle aspiration is inadequate, can sent pus obtained during I&D for C&S. Use parenteral route for severe infections. Consider CA- MRSA if poorly resolving, based on local epidemiology (still generally uncommon).
Animal bites <i>Pasteurella multocida</i> , <i>Staphylococcus</i> spp., <i>Streptococcus</i> spp., <i>Capnocytophaga</i> spp., anaerobes	PO Amoxicillin/clavulanate for 5-7 days	IV Amoxicillin/clavulanate	Consider rabies prophylaxis according to local epidemiology.
Cellulitis <i>Staphylococcus aureus</i> , <i>Streptococcus pyogenes</i>	IV Cloxacillin for 5-7 days	PO Amoxicillin for 7 days OR PO Cephalexin for 5-7 days	Administer using parenteral route for extensive lesions. Total treatment until 3 days after acute inflammation disappears.
Hansen's Disease (leprosy) in children	Paucibacillary: PO Rifampicin PLUS PO Dapsone		-Duration of treatment: 6 months -Surveillance: 5 years

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	Multibacillary: PO Rifampicin PLUS PO Dapsone PLUS PO Clofazimine		-Duration of treatment: 1 year for BI<4; 2 years for BI≥4 (BI: Bacteriological index) -Surveillance: 15 years
Impetigo <i>Staphylococcus aureus</i> , <i>Streptococcus pyogenes</i>	Localised: Topical 2% fusidic acid for 7 days (outpatient)		
	Generalised: PO Cloxacillin	PO Cephalexin	Duration: 5-7 days
Necrotizing fasciitis <i>Streptococcus</i> spp.: GroupA <i>Streptococcus</i> (GABHS), other <i>Streptococcus</i> Staphylococcal: <i>Staphylococcal aureus</i> (MSSA & CA-MRSA)	Streptococcal necrotising Fasciitis: IV Benzylpenicillin PLUS IV Clindamycin		50% of patients have associated streptococcal toxic shock syndrome (STSS). Aggressive surgical debridement of the deep-seated infection is the mainstay of the therapy. Combination therapy is needed with clindamycin to block toxin production whether or not patient manifests toxic shock syndrome. Tissues should be sent for Gram staining & C&S. IVIG can be used as an adjunct. Vancomycin is NOT RECOMMENDED for the treatment of serious MSSA infections because outcomes are INFERIOR compare with cases in which anti-staphylococcus beta-lactam
	Staphylococcal Necrotising Fasciitis: IV Cloxacillin PLUS IV Clindamycin	If CA-MRSA is suspected: IV Vancomycin	

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			(cloxacillin) is used AND to minimise the emergence of vancomycin resistance Duration of treatment: at least 2 weeks if no foci is found (no deep-seated involvement plus no involvement of heart, bone, joint etc.)
Scalded skin syndrome (SSSS) <i>Staphylococcus aureus</i>	IV Cloxacillin <u>Step down</u> PO Cloxacillin Total treatment duration: 7-10 days		If no positive blood culture associated with SSSS, then intravenous therapy can be stopped following clinical improvement & switch to oral.

References:

1. Guidelines for the diagnosis and management of skin and soft tissue infections: 2014 Update by the Infectious Diseases Society of America (IDSA): Clin Infect Dis 2014;59:e10.
2. Malaysian clinical practice guideline on management of leprosy 2014.
3. The Sanford guide to Antimicrobial therapy 2018.
4. American Academy of Paediatrics Committee on Infectious Diseases. Red Book: Report of the committee on Infectious Diseases (2018).

SURGICAL INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
A. General Surgery			
Empyema thoracis (Lung empyema): <i>Staphylococcus aureus</i> <i>Streptococcus pneumoniae</i> Empiric treatment is used to cover organisms mentioned above. Other bacteria: <i>Streptococcus pyogenes</i> , <i>Haemophilus influenzae</i> , other Gram-negative organisms in immunocompromised individuals In patients not responding to treatment need to rule out TB	IV Cefuroxime PLUS IV Cloxacillin	<i>S. aureus</i> (methicillin-sensitive): IV Cloxacillin <i>S. pneumoniae</i> (penicillin-sensitive): IV Benzylpenicillin <i>S. pneumoniae</i> (penicillin-resistant, use result of C&S): IV Cefotaxime OR IV Ceftriaxone	Based on C&S of pleural fluid/tissue or blood culture. All children with empyema need to receive high dose antibiotic therapy via intravenous route to ensure pleural penetration. Pneumatocoeles on CXR indicate <i>S. aureus</i> BUT they can also be seen in pneumococcal disease. There is NO need to routinely use a macrolide antibiotic but its use should be considered in children whom <i>Mycoplasma pneumoniae</i> is thought to be the cause (<i>Mycoplasma</i> usually causes effusion, not empyema). There is NO CONSENSUS on how long antibiotics need to be given. Most recommend 4-6 weeks of total antibiotics. For other adjunct therapy, refer MOH consensus guideline 2013.
Enterocolitis	IV Cefotaxime PLUS	IV Ampicillin PLUS	Antibiotics should be adjusted with results of C&S.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks	
	Preferred	Alternative		
<i>Enterobacteriaceae</i> , <i>Enterococci</i> , <i>Bacteroides</i>	IV Metronidazole	IV Metronidazole		
B. Bone & Joints Infections				
Septic Arthritis (SA) & Osteomyelitis (OM): <u>0-2 months:</u> <i>Staph. aureus</i> <i>Streptococcus agalactiae</i> Gram-negative enteric organisms <u>Less than 5 yrs:</u> <i>Staph. aureus</i> <i>Streptococcus pyogenes</i> <i>Streptococcus pneumoniae</i> Non-typeable <i>Haemophilus</i> spp. <i>Kingella Kingae</i> <u>Older than 5 yrs:</u> <i>Staph. aureus</i> <i>Streptococcus pyogens</i>	0-2 months: IV Cloxacillin PLUS IV Cefotaxime	Optimize antimicrobial treatment based on C&S	Empiric antibiotics should be started based on clinical diagnosis of SA or OM. Surgical debridement is often not required in OM. Urgent washout & drainage is needed in SA in hip and other joints to reduce pressure on growth plate.	
	Children less than 5 yrs: IV Cefuroxime (monotherapy)	IV Cefazolin (can be use in children with suspected <i>S. aureus</i> or <i>S. pyogenes</i> ; less hypersensitivity reaction compared to cloxacillin and dosing convenience) <i>*Kingella kingae</i> - uncommon organism causing infection in <5 yrs old; sensitive to β -lactam antibiotics e.g. cefuroxime or amoxicillin/clavulanate		*IV antibiotics can be switch to oral if no concurrent bacteremia when: Child afebrile and pain free for at least 24 hrs and CRP <20mg/L or CRP decreased by $\geq 2/3$ of highest value.
	Children older than 5 yrs: IV Cloxacillin			Duration of antibiotics: -SA: total of 3-4 weeks -OM: 4-6 weeks -In complex disease (multifocal, significant bone destruction, immunocompromised host and resistant/unusual pathogens), need prolonged intravenous antibiotics and duration might exceed 6 weeks.
References: 1. American Academy of Pediatrics: Pickering LK, BakerCJ, Kimberlin DW, Long SS, eds.Red Book 2012 Report of the committee on Infectious Diseases.				

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
2. Paediatric Empyema Thoracis recommendations for management-Position Statement from the Thoracic society of Australia and New Zealand 2010. 3. Manual of childhood infections-Blue Book 3rd edition; Oxford University Press. 4. Guideline for the management of community acquired pneumonia in children; update 2011.Thorax October 2011: vol 66 (supplement 2) 5. Kathleen Gutierrez. Bone and Joint infections in children. Pediatr Clin N Am 52(2005); 779-794. 6. Approach and management of empyema thoracis in children: a consensus guideline from the paediatric empyema working group 2013-MOH			

TROPICAL INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Typhoid fever	Refer to Gastrointestinal infections Section		
Cholera	Refer to Gastrointestinal infections Section		
Scrub Typhus			
<i>Rickettsia tsutsugamushi</i>	PO Doxycycline for 5-7 days	PO Azithromycin for 3 days	Doxycycline can be used in young children (even below 8 years old) since safety data approved its use for rickettsial diseases.
Brucellosis			
<i>B. melitensis</i> , <i>B. abortus</i> , <i>B. suis</i> and <i>B. canis</i>	PO Rifampicin for 6 weeks PLUS For children <8 years old: PO Trimethoprim/sulfamethoxazole (TMP dose) for 6 weeks OR >8 years old: PO Doxycycline for 6 weeks <u>Serious illness</u> PLUS IV Gentamicin for 7-14 days		For non-localized disease: Can use two-drug combination. Drug of choice for Brucellosis for children >8 years old: Doxycycline (plus rifampicin)
Leptospirosis (<i>L. icterohaemorrhagiae</i> , <i>L. canicola</i>)			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Moderate to severe disease	IV Benzylpenicillin	IV Ceftriaxone OR IV Cefotaxime	Duration: 7 days
Mild disease	PO Amoxicillin	For children >8 years old: PO Doxycycline	
Tetanus (<i>Clostridium tetani</i>)			
	IV Metronidazole for 7-10 days	IV Benzylpenicillin for 7-10 days	Primary tetanus infection: Clinical diagnosis to be made as negative culture is often. Steps in care: 1. Early airway protection & treatment of reflex spasm with benzodiazepine (midazolam). 2. Neutralization of toxin: TIG single dose, administered IM (optimal dose not established). 3. Surgical debridement of infected tissues.
	Neutralization of toxin: Human tetanus globulin (TIG) IM as a single dose	<u>If TIG not available:</u> IVIG as a single dose	
Melioidosis (<i>Burkholderia pseudomallei</i>)			
Intensive/Induction therapy:	IV Ceftazidime	IV Imipenem/cilastatin OR IV Meropenem	Duration: 2-8 weeks ● Uncomplicated : 2 weeks ● Complicated pneumonia, deep-seated infection, neurological melioidosis, osteomyelitis & septic arthritis: 4-8 weeks.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Maintenance therapy:	Child > 8yrs old: PO Trimethoprim/sulfamethoxazole (TMP) PLUS PO Doxycycline in 2 divided doses	Child < 8yrs old: PO Amoxicillin/clavulanate (higher relapse rate)	Duration: 20 weeks PO Folic acid to be given for patients on trimethoprim/sulfamethaxazole. Consider combination therapy of two drugs in maintenance phase if high risk of relapse.
Malaria			
<i>Plasmodium falciparum</i> a) Uncomplicated b) Treatment failure c) Complicated (Almost always due to <i>P. falciparum</i> , suspect mixed infections if <i>P. vivax</i> / <i>P. knowlesi</i> malaria appears more severe than usual)	Uncomplicated PO Artemether/lumefantrine (Riamet®) Duration: 3 days	PO Artesunate/mefloquine FDC (ASMQ) (ASMQ is available as FDC tablet 25/55mg & 100/220mg) Duration: 3 days	Artesunate/mefloquine may cause seizure in children with epilepsy. Riamet® should be served with high fat diet e.g milk to enhance absorption.
	Treatment failure An alternative artemisinin-based combination therapy (ACT) regimen to be used. (If Riamet® is used as the first line regimen, use ASMQ & vice versa)	PO Artesunate PLUS PO Clindamycin for 7 days OR PO Quinine PLUS PO Clindamycin for 7 days	Primaquine 0.25mg base/kg to be given on Day 1 as a single dose in addition to artemisinin-based combination therapy (ACT) (G6PD testing is not required prior to administration of this dose).
	Complicated Day 1: IV Artesunate on admission, then repeat again at 12 & 24 hours.	*IV Quinine PLUS	Avoid using ASMQ (Artesunate/mefloquine) if a patient presents initially with impaired consciousness as increased incidence

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	Day 2 -7: IV Artesunate or switch to PO ACT Duration: 7 days [Parenteral artesunate should be given for a minimum of 24 hours (3 doses) or until patient is able to tolerate orally & thereafter to complete treatment with a complete course of oral ACT (3 days of ASMQ or Riamet®)]	PO Doxycycline OR PO Clindamycin Duration: 7 days	of neuropsychiatric complications associated with mefloquine following cerebral malaria have been reported. Do not use IV artesunate as monotherapy. If IV artesunate needs to be continued indefinitely, clindamycin must be added to the regimen to complete 7 days of treatment. IM artesunate (same dose as IV) can be used in patients with difficult intravenous access. Children with severe malaria should be started on broad-spectrum antibiotic treatment immediately at the same time as antimalarial treatment. *Change to PO quinine if able to tolerate orally. Reduce IV quinine dose by one third of total dose if unable to change to PO quinine after 48 hours or in renal failure or liver impairment.
<i>P. vivax, P. malariae, P. knowlesi</i> (<i>P. malariae</i> & <i>P. knowlesi</i> are chloroquine-sensitive) a) New infection	New infection: PO chloroquine for 3 days PLUS PO Primaquine for 14 days		Primaquine is ONLY needed for <i>P. vivax</i> for a total of 14 days. Primaquine (0.5mg/kg) may cause haemolysis in individuals with G6PD

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
b) Chloroquine-resistant (<i>P. vivax</i>) or relapse c) Severe & complicated <i>P. vivax</i> , <i>P. knowlesi</i> or <i>P. malariae</i>	Chloroquine-resistant (<i>P. vivax</i>) or relapse: (*In Sabah, considered chloroquine-resistant) ACT (Riamet or ASMQ) (dosing as per <i>P. falciparum</i> treatment) PLUS PO Primaquine for 14 days	PO Quinine for 7 days PLUS PO Primaquine for 14 days Mefloquine single dose combined with primaquine have been found to be effective (except for <i>P. knowlesi</i>)	deficiency, hence G6PD testing is required before administration of primaquine above 0.25mg/kg. For those found to have mild to moderate G6PD deficiency, an intermittent primaquine regimen of 0.75mg base/kg weekly for 8 weeks can be given under medical supervision. In severe G6PD deficiency, primaquine is contraindicated & should not be used. <i>P. knowlesi</i> (monkey malaria) can cause severe malaria & should be treated as severe malaria secondary to <i>P. Falciparum</i> .
	Severe & complicated <i>P. vivax</i>, <i>P. knowlesi</i> or <i>P. malariae</i> Treat as per severe <i>P. falciparum</i> malaria		
Mixed Infection	Treat as <i>P. falciparum</i>		
References: 1. Keren Saklsky et al. Treatment of Human Brucellosis: systematic review and meta-analysis of randomized controlled trials. BMJ online. 2008. doi:10.1136/bmj.39497.500903.25 2. Silpapojakul K. Paediatric scrub typhus in Thailand: a study of 73 confirmed cases. Trans R Soc Trop Med & Hygiene 2004;98:354-9 3. Phimda K et al. Doxycycline versus azithromycin for treatment of leptospirosis and scrub typhus. Antimicrob Agents Chemother. 2007;51(9):3259 4. Szue-Chia Lee et al. Comparative effectiveness of Azithromycin for treating scrub typhus. A PRISMA- compliant systematic review and meta-analysis. Medicine. 2017. 96:36(e7992) 5. Fang Y, Huang Z, Tu C, Zhang L, Ye D, Zhu B. Meta-analysis of Drug Treatment for Scrub Typhus in Asia. Intern Med. 2012;;51: 233-2320 6. Guidelines for the Diagnosis, Management, Prevention and Control of Leptospirosis in Malaysia, MOH 2011			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
7. Watt G, Padre LP, Tuazon ML, et al. Placebo-controlled trial of intravenous penicillin for severe and late leptospirosis. Lancet 1988; 1:433-5 8. Panaphut T. Ceftriaxone compared with sodium penicillin G for treatment of severe leptospirosis. ClinInfect Dis 2003; 36:1507-13 9. Suputtamongkol Y. An Open, Randomized, Controlled Trial of Penicillin, Doxycycline, and Cefotaxime for Patients with Severe Leptospirosis. Clin Infect Dis 2004; 39:1417-24 10. Currie BJ. Melioidosis: Evolving Concepts in Epidemiology, Pathogenesis, and Treatment. Semin Respir Crit Care Med. 2015;36:111-125 11. McLeod C, Morris PS, Bauert PA, Kilburn CJ, Ward LM, Baird RW, Currie BJ. Clinical Presentation and Medical Management of Melioidosis in children A 24-Years Prospective Study in the Northern Territory of Ausralia and Reivew of the Literature. Clin Inf Dis. 2015; 60(1):21-6 12. Cheng AC, Chierakul W, Chaowagul W, et al. Consensus guidelines for dosing of amoxicillin- clavulanate in melioidosis. Consensus guidelines for dosing of amoxicillin-clavulanate in melioidosis. Am J Trop Med Hyg. 2008; 78(2):208 13. WHO Guidelines for the treatment of malaria 2015.WHO/HTM/MAL/2015 14. The Sanford guide to antimicrobial therapy 2018. 15. American Academy of Paediatrics. Committee on Infectious Diseases. Red Book: Report of the Committee on Infectious Diseases (2018).			

TUBERCULOSIS IN CHILDREN

1. Anti Tuberculosis Drugs

Table 1: Recommended doses of anti-TB drugs in children

Drug	Recommended Daily Dose	
	Dose (range) in mg/kg	Maximum dose in mg
Isoniazid(H)	10 (7- 15) ^a	300
Rifampicin(R)	15(10- 20)	600
Pyrazinamide(Z)	35(30- 40)	2000
Ethambutol(E)	20(15- 25)	1000

^a The higher end of the range for INH dose applies to younger children. As the children grow older the lower end of the dosing range becomes more appropriate.

Source: World Health Organization. Guidance for National Tuberculosis Programmes on The Management of Tuberculosis in Children (2nd Edition). Geneva: WHO;2014

Pyridoxine 5 - 10 mg daily needs to be added if isoniazid is prescribed.

Medication dose requires recalculation **every two to four weeks** as children gain weight rapidly, particularly in neonates and young children.

2. Treatment Regimen

- Treatment has 2 phases: an initial intensive phase and a second continuation phase.
- Daily directly-observed therapy (DOT) is recommended for treatment of active disease.
- Use of steroids:
Corticosteroids should be used in tuberculous meningitis and pericarditis (pericardial effusion).

3. Latent Tuberculosis Infection Treatment (LTBI)

*Active TB needs to be ruled out before starting LTBI treatment.

- Current regimens used to treat LTBI in children include:
 - a. RIF daily for four months (4R)
 - b. INH and rifapentine weekly for three months (12 doses) (3HP)
 - c. INH and RIF daily for three months (3HR)
 - d. INH daily for six or nine months (6H/9H)
- In the treatment of children with latent tuberculosis infection (LTBI), the preferred regimens are: 4R for all children >28 days of age or 3HP* for children aged >2 years. 6H for all newborns aged 28 days and below.
- Alternative regimens of LTBI in children are 3HR, 6H or 9H.

Table 2: Recommended dosage for LTBI treatment in children

Drug	Duration	Interval	Dosage
Isoniazid	6 months	Daily	1. Age 10 years and older: 5mg/kg/day 2. Age < 10 years: 10mg/kg/day (Range: 7 – 15mg/kg) Max dose: 300mg
Rifampicin (4R)	4 months	Daily	1. Age 10 years and older: 10mg/kg/day 2. Age < 10 years: 15mg/kg/day (Range: 10-20 mg/kg)

			Max dose: 600mg	
Isoniazid + Rifampicin (3HR)	3 months	Daily	Dose of INH and RIF same as above	
Rifapentine + Isoniazid (3HP)	3 months*	Weekly	Isoniazid: 10-15 kg : 300 mg 16-23 kg : 500 mg 24-30 kg : 600 mg >31 kg : 700 mg (For children age 2-14 years old)	Rifapentine: 10-15 kg : 300 mg 16-23 kg : 450 mg 24-30 kg : 600 mg >31 kg : 750 mg (For children age 2-14 years old)

*Given in total of 12 doses

Pyridoxine 5 - 10 mg/day should be given to patients on Isoniazid.

Source: World Health Organization. WHO Consolidated Guidelines on Tuberculosis: Tuberculosis Preventive Treatment: Module 1: Prevention. Geneva: World Health Organization; 2020

References:

- 1.Clinical Practice Guideline Management of Tuberculosis. 4 th edition. 2021.
- 2.World Health Organization. Guidance for National Tuberculosis Programmes on The Management of Tuberculosis in Children (2nd Edition). Geneva: WHO;2014
- 3.World Health Organization. WHO Consolidated Guidelines on Tuberculosis: Tuberculosis Preventive Treatment: Module 1: Prevention. Geneva: World Health Organization; 2020.

URINARY TRACT INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Urinary Tract Infection: <i>Escherichia coli</i> <i>Proteus</i> spp <i>Klebsiella</i> spp. <i>Enterobacter</i> spp.	0-2 months old: IV Ampicillin PLUS IV Gentamicin > 2 months old: IV Cefuroxime PLUS/MINUS IV Gentamicin	III Young Infant: IV Cefotaxime	Adjust therapy base on culture results Duration: 7-14 days Switch to oral therapy when improving and being able to tolerate oral therapy.
Prophylaxis for UTI for infants & children with recurrent UTI	PO Trimethoprim at night	PO Nitrofurantoin at night OR PO Cephalexin at night	Antibiotic prophylaxis should not be routinely recommended in children with first-time UTI. Avoid nitrofurantoin for infants less than 3 months old, G6PD deficiency, and estimated glomerular filtration rate < 50 ml/min/1.73m ² . Prophylactic antibiotics should be given for 3 days with MCUG (micturating cystourethrogram) taking place on the second day. Children with Grade 4-5 vesicoureteral reflux (VUR) require prophylaxis antibiotic whereas Grade 3 VUR may require prophylaxis antibiotic if associated with many genitourinary tract anomaly with high risk of reflux

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			nephropathy or frequent symptomatic UTI.
References: 1. NICE Guidelines: Urinary tract infection: diagnosis, treatment and long term management of urinary tract infection in children 2007. Last update 2017 2. UTI Clinical Practise Guideline, Pediarics 2011 3. American Academy of Paediatrics. Commitee on Infectious Diseases. Red Book: Report of the Committee on Infectious Diseases (2018). 4. The Sanford Guide to Antimicrobial Therapy 2018 5. BNF Children 2022			

VASCULAR INFECTIONS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Catheter-related BloodStream Infection (CRBSI)			
*Coagulase-negative staphylococcus (CoNS)			
Methicillin-sensitive (MSCoNS)	IV Cloxacillin	IV Cefazolin	Diagnosis needs: 1. Paired blood cultures drawn from both catheters & peripheral veins. 2. If blood cultures cannot be drawn from peripheral veins, it is recommended that two or more blood cultures should be drawn through different catheter lumen. Long term catheters should be removed in patients with CRBSI with: Severe sepsis, suppurative thrombophlebitis, endocarditis, blood stream infections that continues despite 72 hours of antimicrobial therapy or longer to which the infecting organism is susceptible or infection due to <i>Staphylococcus aureus</i> , <i>Pseudomonas aeruginosa</i> , fungi & <i>Mycobacterium</i> . Attempts at catheter salvage are only recommended in uncomplicated CRBSI or CLABSI caused by bacteria that are neither too virulent nor too difficult to eradicate.
Methicillin-resistant (MRCoNS)	IV Vancomycin		
Coagulase-positive staphylococcus			
Methicillin-sensitive (MSSA)	IV Cloxacillin	IV Cefazolin	
Methicillin-resistant (MRSA)	IV Vancomycin		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			Exact optimal duration of therapy has not been established in children with or without catheter removal. 10-14 days after the first negative blood culture is usually recommended. *For CONS, we need to decide whether isolates from blood culture are colonisers or true pathogens.
Gram-negative bacilli Enterobacterales (<i>Escherichia coli</i> , <i>Klebsiella pneumonia</i> , <i>Enterobacter</i> , <i>Proteus</i> sp etc.)			
ESBL negative	*IV Piperacillin/tazobactam		*Empiric treatment with piperacillin/tazobactam covers most of the Gram-negative organisms (Enterobacterales), Gram-positive organisms & <i>Pseudomonas</i> ; follow through with C&S.
ESBL positive	IV Ertapenem	IV Imipenem OR IV Meropenem	
<i>Pseudomonas aeruginosa</i>	IV Piperacillin/tazobactam	IV Cefepime	Not all <i>Pseudomonas</i> is drug resistant. If ceftazidime remains susceptible. Please use it with antipseudomonal aminoglycoside to treat. De-escalation is important to preserve antibiotics for future use.
<i>Candida albicans</i> or other <i>Candida</i> species			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	IV Fluconazole	IV Caspofungin OR IV Amphotericin B lipid complex	Fungaemia: treatment without catheter removal is associated with low success rate & higher mortality.
Suppurative Thrombophlebitis (<i>Staphylococcus aureus</i>)			
MSSA	IV Cloxacillin	IV Cefazolin	Diagnosis requires positive blood culture plus radiographic demonstration of thrombus. Remove catheter & a minimum antibiotic treatment of 3-4 weeks. Surgical resection of involved vein if failed conservative therapy.
MRSA	IV Vancomycin		
References: 1. IDSA guidelines for intravascular catheter-related infection. CID 2009; 49:1-45. 2. Patricia MF. Diagnosis and management of central-venous catheter-related bloodstream infections in pediatric patients. Pediatr Infect Dis J. 2009; 28(11):1016-1017. 3. Michael JS, Catheter related bloodstream infection in children. Am J Infect Control 2008; 36:S173.e1-S173.e3. 4. Janum S et al.Bench to bedside review: Challenges in diagnosis, care and prevention of central catheter related bloodstream infections in children. Critical Care 2013.17:238. 5. The Sanford guide to antimicrobial therapy 2018.			

CHEMOPROPHYLAXIS

A. SURGICAL CHEMOPROPHYLAXIS

Guidelines for prevention of surgical site infections (SSIs) have been published. General principles:

1. Agent used for antimicrobial prophylaxis should prevent SSIs and related morbidity and mortality
2. Reduce duration and cost of care
3. Produce no adverse effect
4. Minimise adverse consequences to the microbial flora

Timing:

Effective chemoprophylaxis occurs only when the appropriate antimicrobial drug is present in tissues at sufficient local concentration at the time of intra-operative bacterial contamination. Administration of antimicrobial agents is recommended within 60 minutes before surgical incision to ensure adequate tissue concentration at the start of the procedure. Agents that require longer administration time such as vancomycin should begin within 120 minutes before surgery begins. Adequate antimicrobial concentration should be maintained throughout the surgical procedure and in most instances, single dose of antimicrobial agent is sufficient and the duration of prophylaxis after any procedure should not exceed 24 hours. Intraoperative dosing is required if the duration of the procedure is greater than two times the half-life of the antimicrobial agent or if there is excessive blood loss.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Cardiac surgery			
<i>S. epidermidis</i> , <i>S. aureus</i> , <i>Corynebacterium sp.</i> , <i>Enteric Gram-negative bacilli</i>	IV Cefazolin	If known to have MRSA/MRSE colonisation, use IV Vancomycin	β-lactam Allergy: IV Clindamycin
Thoracic surgery			
Non-cardiac including lobectomy, pneumonectomy, lung resection & thoracotomy	IV Cefuroxime	IV Ampicillin/sulbactam	β-lactam Allergy: IV Clindamycin
Abdominal surgery			
Gastroduodenal	IV Cefotaxime PLUS/MINUS IV Metronidazole	IV Ampicillin/sulbactam	β-lactam Allergy: IV Clindamycin PLUS IV Gentamicin

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Biliary tract Kasai procedure/Choledochal cyst excision	IV Cefotaxime PLUS IV Metronidazole	IV Ceftriaxone PLUS IV Metronidazole	β-lactam Allergy: IV Clindamycin PLUS IV Gentamicin
Inguinal surgery Orchidopexy (Open procedure/Laparoscopic procedure)/Hernia repair (Herniotomy)	No antibiotic routinely	No antibiotic routinely	
Head & neck			
Clean (tonsillectomy, adenoidectomy, tracheostomy, lymph node biopsy)	No antibiotic routinely	No antibiotic routinely	
Clean (preauricular sinus, dermoid cyst, branchial anomaly, thyroidectomy, thyroglossal cyst excision, parotidectomy)	IV Augmentin	IV Cefuroxime OR IV Ceftriaxone	β-lactam Allergy: IV Clindamycin
Clean with placement of prosthesis (excludes tympanostomy tubes)	IV Cefazolin	IV Ampicillin/sulbactam OR IV Cefuroxime	β-lactam Allergy: IV Clindamycin
Clean-contaminated procedures with the exception of tonsillectomy & functional endoscopic sinus procedure	IV Cefazolin PLUS IV Metronidazole	IV Ampicillin/sulbactam OR IV Cefuroxime PLUS IV Metronidazole	β-lactam Allergy: IV Clindamycin
Clean-contaminated cancer surgery	IV Cefotaxime PLUS IV Metronidazole	IV Ampicillin/sulbactam OR	β-lactam Allergy: IV Clindamycin

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
		IV Cefuroxime PLUS IV Metronidazole	
Neurosurgery			
Elective craniotomy & cerebrospinal fluid shunting procedures Types of surgery: Craniotomy/crani-facial Transsphenoidal surgery Spine surgery (Intradural/Extradural lesion) Shunt surgery & Neuroendoscopy Neuroendovascular procedure	IV Cefazolin	If known to have MRSA/MRSE colonisation, use IV Vancomycin	β-lactam Allergy: IV Clindamycin
Orthopaedics			
Clean operations involving hand, knee or foot & not involving implantation of foreign materials	None	None	
Spinal procedure with or without instrumentation/hip surgery/implantation of internal fixation devices (e.g. nails, screws, plates, wires)	IV Cefazolin		β-lactam Allergy: IV Clindamycin
Urology			
Low tract instrumentation with risk factors for infections	IV Cefuroxime	IV Cefazolin	β-lactam Allergy: IV Gentamicin
Clean without entry into urinary tract/clean with entry into	IV Cefuroxime	IV Amoxicillin/clavulanate	UTI should be treated before procedure when possible.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
urinary tract (e.g. hypospadias surgery)			Medical literature does not support continuing antimicrobial prophylaxis until the urinary catheter has been removed.
Clean-contaminated (entering gastrointestinal tract)	IV Cefotaxime PLUS IV Metronidazole	IV Amoxicillin/clavulanate	β-lactam Allergy: IV Gentamicin
Plastic surgery			
Elective soft tissue surgery	No prophylaxis unless complex prolonged procedure If complex, IV Cloxacillin		
Elective hand & foot surgery involving bone	IV Cloxacillin		β-lactam Allergy: IV Clindamycin
Cleft lip & palate surgery	IV Amoxicillin/clavulanate		β-lactam Allergy: IV Clindamycin
Excision & grafting surgery	IV Amoxicillin/clavulanate		β-lactam Allergy: IV Clindamycin
Interventional radiology			
Percutaneous endoscopic gastrostomy (PEG) or jejunostomy (PEJ) or nephrostomy tube placement	IV Cefazolin	IV Amoxicillin/clavulanate	
Micturating cystourethrogram (MCUG)	PO Trimethoprim (If patient is already on existing antibiotic UTI prophylaxis, increase antibiotic to therapeutic dose for a single dose prior procedure)		
Tenckhoff peritoneal dialysis catheter insertion	IV Cefuroxime	IV Amoxicillin/clavulanate	
Burns	No prophylaxis required		

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
References: 1. Clinical Practical Guideline for Antimicrobial Prophylaxis for Surgery 2013. American Society of Hospital Pharmacists (ASHP) guideline, IDSA, Surgical Infection Society (SIS) and Society of Healthcare Epidemiology of America (SHEA). Bratzler DW, Dellinger EP, Olsen KM, Perl TM, Auwaerter PG, Bolon MK, Fish DN, Napolitano LM, Sawyer RG, Slain D, Steinberg JP, Weinstein RA. American Journal of Health-system Pharmacy 2013. 70(3):195-283. 2. Antibiotic Prophylaxis for Paediatric Surgery. Royal Hospital for Children Glasgow. 2018.			

B. NON-SURGICAL CHEMOPROPHYLAXIS

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
Rheumatic fever (Secondary prevention)	IM Benzathine Penicillin <u>Duration</u> 1. With carditis & residual heart disease (persistent valvular disease): 10 years since last ARF or until 40 years of age. Whichever is longer. Consider lifelong prophylaxis 2. With carditis but no residual heart disease (no valvular disease): 10 years since last ARF or 21 years of age, whichever is longer 3. Without carditis: 5 years since last ARF or until 21 years of age, whichever is longer	PO Phenoxymethylpenicillin (Penicillin V) Penicillin Allergy: PO Erythromycin Ethylsuccinate	
Infective Endocarditis (IE)	PO Amoxycillin 30-60 minutes before procedure OR IV Ampicillin 30 -60 minutes before procedure	Penicillin Allergy: IV/PO Clindamycin 30-60 minutes before procedure Other alternative: IV Cefazolin (Cephalosporin should not be used in children with anaphylaxis, angioedema or urticaria)	IE prophylaxis recommended for patients with the highest risk cardiac conditions undergoing procedures likely to result in bacteremia with a microorganism that has the potential ability to cause bacterial endocarditis. Prophylaxis always required for: Dental procedures that involve extraction, periodontal procedure including surgery, subgingival scaling, root planning, re-planting of avulsed teeth and other surgical procedure.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			- Incision and drainage of local abscess in brain, skin, subcutaneous tissue, epidural, lung orbital area, per rectal area, liver, tooth and surgical procedure through infected skin. - Percutaneous endoscopic gastrotomy. Maintenance of optimal oral hygiene may reduce the incidence of bacteremia from daily activities and is more important than prophylactic antibiotics for a dental procedure to reduce the risk of IE.
Post-splenectomy At risk for <i>Pneumococcus</i> , <i>Meningococcus</i> , <i>Haemophilus</i>	PO Penicillin V Duration: Minimum 3 years post-splenectomy or until 18 years of age (some expert) OR At least 1 year post splenectomy Asplenia attributed to other causes unknown - most experts recommend throughout childhood and adulthood.	PO Amoxicillin <u>Penicillin Allergy:</u> PO Erythromycin Ethylsuccinate	Risk of sepsis is lifelong but especially high in the first two years of splenectomy. Important adjunct: Immunization against <i>Pneumococcus</i> <i>Haemophilus</i> , <i>Meningococcus</i> at least 14 days prior to splenectomy (if not possible then as soon as possible, 14 days or more after surgery) Yearly influenza vaccine is also recommended. Not all pneumococcal isolates are sensitive to these antibiotics. Limitation stressed to parents so that all febrile illness in this group of

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			children are taken seriously since initial signs & symptoms of fulminant septicemia can be subtle.
<i>Haemophilus influenza b</i> exposure	PO Rifampicin for 4 days		Chemoprophylaxis indicated for: 1. <u>All household</u> contacts in the following circumstances (Household contact is defined as a person who resides with the index patient or who spent ≥ 4 hours with index patient for at least five of seven days before the day of hospital admission of index case): a) Household with at least one contact <4 years old who is unimmunised or incompletely immunised. b) Household with contact who is an immunocompromised child, regardless of the child's Hib immunisation status. c) Household with a child younger than 12 months who has not complete the primary Hib series. 2. <u>Nursery Contact</u>: For ALL attendees in childcare & preschool (regardless of age or vaccine status) when unimmunized or incompletely immunized children attend the facility and two or more cases of Hib invasive

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			disease have occurred among attendees within 60 days. 3. <u>Index case</u>: Prior to discharge if did not receive at least ONE dose of cefotaxime/ceftriaxone & infants younger than 2 years old. For contacts < 2 years old who are not immunised, recommended to complete immunisation.
Meningococcal exposure	PO Rifampicin for 2 days	IM Ceftriaxone stat	Chemoprophylaxis is provided to close contact at HIGH RISK which include: - All household especially children younger than 2 years old - Childcare or preschool contact at anytime during 7days before onset of illness. - Direct exposure to index patient's secretion through kissing or through sharing toothbrushes or eating utensils at any time during 7 days before onset of illness. - Frequently slept in the same place as the index during 7 days before onset of illness. <u>Healthcare staff</u> Routine prophylaxis is not recommended unless there is intimate exposure to respiratory secretion, mouth to mouth resuscitation. Unprotected contact

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			during intubation at any time 7 days before onset of illness or within 24hours of initiation of effective antimicrobial therapy. Give chemoprophylaxis to index case prior to discharge if treated with regimes other than Ceftriaxone or Cefotaxime. Chemoprophylaxis is ideally initiated within 24hours after index patient is identified; prophylaxis is not indicated more than 2 week after exposure.
Neonatal Group B <i>Streptococcus</i> Infection	Intrapartum maternal prophylaxis till delivery: IV Benzylpenicillin	IV Ampicillin <u>Penicillin allergy:</u> IV Clindamycin (according to susceptibility)	Treat during labour if previously delivered infants with invasive GBS, GBS bacteriuria or antenatally screening swab positive OR if GBS status is not known AND any of the following: - Preterm <37 weeks - PPROM > 18 hours - Intrapartum temperature > 38°C
Malaria prophylaxis	PO Chloroquine weekly (begin 1-2 weeks before travelling and take weekly throughout & 4 weeks after leaving the area)	For prophylaxis (chloroquine-resistant): *Atovaquone-proguanil OR **PO Mefloquine (Start 2-3 weeks before, continue weekly during exposure and 4 weeks thereafter) OR Age > 8 years old:	*Atovaquone/proguanil is another drug used in malaria prophylaxis in children (chloroquine resistance) BUT not yet registered in Blue book (available commercially in Malaysia). **Mefloquine: Not available in HRPB Formulary

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
		PO Doxycycline daily 1-2 days before, during & 4 weeks after travelling.	**Mefloquine: Not recommended if there are cardiac conduction abnormalities, seizures or psychiatric disorders eg: depression, psychosis (Black box warning: Neuropsychiatric reaction may persist even after discontinuation). To carefully assess risk and benefit of starting antimalarial prophylaxis to any children to prevent development of drug resistance.
Pertussis (Post-Exposure prophylaxis, PEP)	<u>Less than 1 month - 5 months old:</u> Azithromycin for 5 days <u>6 months or older:</u> Erythromycin Ethylsuccinate for 14 days OR Azithromycin 5 days	Erythromycin is not preferred in young infants. *Use only if azithromycin is not available. <u>2 months and older:</u> Trimethoprim/sulphamethoxazole for 14 days	Drug of choice for PEP & treatment is macrolide. Azithromycin is the preferred macrolide. *Association between orally administered azithromycin & erythromycin with infantile hypertrophic pyloric stenosis (especially in infant < 6 weeks) has been reported but azithromycin remains the drug of choice in very young infants because risk of developing severe disease outweighs the potential risk. Antimicrobial prophylaxis is recommended for: ALL household contacts of the index cases & other close contacts including children in childcare regardless of immunization status.

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
			When considering borderline degree of exposure for non household contact, PEP administered if contact is personally at high risk** or lives in a household with high risk of severe disease (e.g. young infant, pregnant women person who has contact with infant). Close contact who are unimmunized or underimmunized should have pertussis immunization initiated or continue with age appropriate products according to the recommended schedule as soon as possible. (includes Tdap in children 7-9 years old who did not complete DTaP series). **High risk: infant, women at 3rd trimester of pregnancy & people with pre-existing health conditions. (not limited to immunocompromised and those with moderate to severe asthma)
Chicken pox (Post-exposure prophylaxis) Potential interventions for people without immunity exposed to varicella following significant exposure* 1. Vaccine	Varicella vaccine:		*Exposure is significant if: 1. Household: Residing in same household 2. Playmate: Face to face indoor play >1 hour 3. Hospital: In same 2 to 4 bedroom or adjacent beds in a large

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
	Within 3-5 days of exposure for the susceptible healthy adult/child 12 months or older (followed by second dose at age-appropriate interval)		ward, face to face contact with an infectious staff member or patient or visit by a person who is contagious. 4. Newborn infants
2. When indicated & available, Varicella zoster immune globulin (VZIG)	For patient who are at high risk for severe infection & complication* & significant exposure (and have contraindication to vaccine): VZIG as soon as possible after exposure up to 10 days after	Patients receiving monthly high dose IVIG ($\geq 400\text{mg/kg}$) are likely to be protected & probably do not require VZIG if the most recent dose IVIG was administered ≤ 3 weeks before exposure.	*Susceptible host include: 1. Immunocompromised children 2. Pregnant women, newborn of mothers with varicella shortly before or after delivery. (5 days before or within 2 days after delivery) 3. Premature born infants born ≥ 28 weeks of gestation who are exposed during hospitalization & whose mother does not have any evidence of immunity 4. Premature born infants born < 28 weeks of gestation or birth weight $\leq 1000\text{g}$ regardless of mothers' immunity.
3. When VZIG not available	For patient who are at high risk for severe infection & complication* & significant exposure (and have contraindication to vaccine): IVIG stat OR PO Acyclovir beginning 7 to 10 days after exposure & continued for 7 days		
References: - Davies JM, Lewis MP, and Wimperis J et al. Review of guidelines for the prevention and treatment of infection in patient with an absent or dysfunctional spleen: prepared on behalf of the British Committee for Standards in Haematology by a working party of the Haemato- Oncology Task Force. Br Haematol. 2011; 155:308-17. - Rubin LG, Schaffner W. Care of the Asplenic Patient. N Engl J Med 2014; 317:349-56 - Gardner P. Clinical practice. Prevention of meningococcal disease. N Engl J Med 2006;355:1466 - Guidance for the public health management of meningococcal disease in UK. Updated February 2018. Public Health England. - Verani JR, McGee L, Schrag SJ, Division of Bacterial Disease, National Center for Immunization and Respiratory Diseases, Centers for Disease Control and Prevention (CDC). Prevention of perinatal group B streptococcus disease-- revised guidelines from CDC, 2010. MMWR Recomm Resp 2010; 59:1 - Updated recommendation for use of VariZIG-- United States, 2013. Centres for Disease Control and Prevention (CDC). MMWR Mortal Wkly Rep. 2013;62(28):574			

Infection/ Condition & Likely Organism	Suggested Treatment		Remarks
	Preferred	Alternative	
7.	WHO Malaria Treatment Guideline 2015		
8.	The Sanford Guide to Antimicrobial therapy 2018		
9.	American Academy of Paediatrics. Committee on Infectious Diseases. Red Book: Report on the committee on Infectious Disease (2018).		

(APPENDICES)

APPENDIX 1: ANTIBIOTIC RESISTANCE DATA (2019 TO 2021)

A. Common organisms isolated

1. By isolates




Figure 1: Common organisms isolated from all samples by isolates

2. By patients




Figure 2: Common organisms isolated from all samples by patients

B. Percentage of antibiotic resistance among bacteria

Gram-positive bacteria

1. *Staphylococcus aureus*

MSSA




Figure 3: Antibiotic resistance trend for MSSA isolated from all samples against selective antibiotics

MRSA




Figure 4: Antibiotic resistance trend for MRSA isolated from all samples against selective antibiotics

2. *Streptococcus pneumoniae*




Figure 5: Antibiotic resistance trend for *Streptococcus pneumoniae* isolated from all samples against selective antibiotics

3. *Enterococcus faecium*




Figure 6: Antibiotic resistance trend for *Enterococcus faecium* isolated from all samples against selective antibiotics

4. *Enterococcus faecalis*




Figure 7: Antibiotic resistance trend for *Enterococcus faecalis* isolated from all samples against selective antibiotics

Gram-negative bacteria

1. *Acinetobacter baumannii*




Figure 8: Antibiotic resistance trend for *Acinetobacter baumannii* isolated from all samples against selective antibiotics

2. *Escherichia coli*




Figure 9: Antibiotic resistance trend for *Escherichia coli* isolated from all samples against selective antibiotics

3. *Klebsiella pneumoniae*




Figure 10: Antibiotic resistance trend for *Klebsiella pneumoniae* isolated from all samples against selective antibiotics

4. *Pseudomonas aeruginosa*




Figure 11: Antibiotic resistance trend for *Pseudomonas aeruginosa* isolated from all samples against selective antibiotics

APPENDIX 2: ANTIBIOTIC UTILISATION DATA (2018 TO 2021)

1. Penicillins




Figure 12: Utilisation of injection of the penicillin group

2. Third- and fourth-generation cephalosporins




Figure 13: Utilisation of cephalosporin injection

3. Carbapenems




Figure 14: Utilisation of carbapenem injection vs rate of ESBL infection

4. Polymyxins




Figure 15: Utilisation of polymyxin injection vs rate of CRE infection

5. Fluoroquinolones




Figure 16: Utilisation of fluoroquinolone injection

6. Vancomycin




Figure 15: Utilisation of vancomycin injection vs rate of MRSA infection

APPENDIX 3: ANTIMICROBIAL DOSAGES IN PATIENTS WITH IMPAIRED RENAL FUNCTION (ADULTS)

The Cockcroft-Gault equation estimates creatinine clearance (CrCl) in mL/minute and has been successfully used in the calculation of adults with renal impairment. Weight, age, gender and serum creatinine are required. The formula below should not be used to estimate CrCl in severe renal insufficiency, or with rapidly changing renal function.

(i) Modified Cockcroft-Gault equation:

$$CrCl (mL/min) = \frac{(140 - age) \times IBW (kg)}{Serum creatinine (\mu mol/L)} \times (1.04 \text{ for females}) \text{ or } (1.23 \text{ for males})$$

*If actual body weight (ABW) is < ideal body weight (IBW), use actual body weight (ABW)

(ii) Ideal body weight calculation:

Males: IBW (kg) = 50kg + 2.3kg for each inch over 5 feet

Females: IBW(kg) = 45.5kg + 2.3kg for each inch over 5 feet

*Use IBW if overweight but non-obese

(iii) Dosing weight calculation:

$$\text{Dosing weight (DW)} = IBW + [0.4 \times (ABW - IBW)]$$

*Use DW if obesity (ABW ≥ 125% of IBW)

ANTIMICROBIAL DOSAGES IN PATIENTS WITH IMPAIRED RENAL FUNCTION (ADULTS)

**** Unless stated otherwise, adjusted doses are % of dose for normal renal function.**

**** For critical care patients, kindly refer to Guide to Antimicrobial Therapy in the Adult ICU 2017 page 123-131 for dosage adjustment for renal impairment.**

ANTIMICROBIAL	DOSE FOR NORMAL RENAL FUNCTION	ADJUSTMENT FOR RENAL FAILURE Estimated Creatinine Clearance (CrCl), mL/min			DIALYSIS DOSE	COMMENTS									
		> 50	10-50	<10											
ANTIBACTERIAL															
(A) Aminoglycosides															
Amikacin	7.5mg/kg q12h	100% of normal dose q12h or 24hr by levels	100% of normal dose q24-72h by levels	100% of normal dose q48h-72h by levels	HD: 7.5mg/kg q48h post HD	Dosage adjustment should be based on TDM level where possible.									
Gentamicin	LD: 2mg/kg; MD: 1.7-2mg/kg q8h	100% of normal dose q8h by levels	100% of normal dose q12-48h by levels	100% of normal dose q48h by levels	HD: 1.7-2mg/kg q48h post HD	Dosage adjustment should be based on TDM level where possible.									
Streptomycin (IM)	15mg/kg (max. 1gm) q24h OR 25-30mg/kg 2x/week OR 25-30mg/kg 3x/week	100% of normal dose q24h	100% of normal dose q24-72h	100% of normal dose q72-96h	HD: 15mg/kg q72-96h post HD										
(B) Carbapenems															
Imipenem	<table><tr><td rowspan="2">Normal dose</td><td colspan="4">CrCl (mL/min/1.73 m²)</td></tr><tr><td>≥ 60</td><td>≥ 30 to < 60</td><td>≥ 15 to < 30</td><td>< 15</td></tr></table>				Normal dose	CrCl (mL/min/1.73 m²)				≥ 60	≥ 30 to < 60	≥ 15 to < 30	< 15	HD: 250-500mg q12h	
Normal dose	CrCl (mL/min/1.73 m²)														
	≥ 60	≥ 30 to < 60	≥ 15 to < 30	< 15											

ANTIMICROBIAL	DOSE FOR NORMAL RENAL FUNCTION	ADJUSTMENT FOR RENAL FAILURE Estimated Creatinine Clearance (CrCl), mL/min				DIALYSIS DOSE	COMMENTS									
		> 50	10-50		<10											
	<table><tr><td>500mg q6h</td><td>100% of normal dose q6h</td><td>250mg q6h OR 500mg q8h</td><td>250mg q8h OR 500mg q12h</td><td rowspan="3">500mg BD *as long as HD will be started within 48hrs</td></tr><tr><td>1000mg q8h</td><td>100% of normal dose q8h</td><td>500mg q8h</td><td>250mg q8h OR 500mg q12h</td></tr><tr><td>1000mg q6h</td><td>100% of normal dose q6h</td><td>500mg q6h</td><td>250mg q6h</td></tr></table>	500mg q6h	100% of normal dose q6h	250mg q6h OR 500mg q8h	250mg q8h OR 500mg q12h	500mg BD *as long as HD will be started within 48hrs	1000mg q8h	100% of normal dose q8h	500mg q8h	250mg q8h OR 500mg q12h	1000mg q6h	100% of normal dose q6h	500mg q6h	250mg q6h		
500mg q6h	100% of normal dose q6h	250mg q6h OR 500mg q8h	250mg q8h OR 500mg q12h	500mg BD *as long as HD will be started within 48hrs												
1000mg q8h	100% of normal dose q8h	500mg q8h	250mg q8h OR 500mg q12h													
1000mg q6h	100% of normal dose q6h	500mg q6h	250mg q6h													
Meropenem	1-2g q8h	100% of normal dose q8h	26-50mL/min: 100% of normal dose q12h 10-25mL/min: 50% of normal dose q12h	50% of normal dose q24h	CAPD, HD: 500mg 24h											
Ertapenem	1g q24h	100% of normal dose q24h	<30 mL/min: 50% of normal dose q24h	50% of normal dose q24h	CAPD, HD: 500mg q24h	If dosed <6 hours prior to HD, to give 150mg supplemental dose after dialysis.										
(C) Cephalosporins																
Cefazolin (IV/IM)	1-2g q8h (max 12g/day)	100% of normal dose q8h	100% of normal dose q12h	50% of normal dose q24-48h	HD: 1-2g q24h											
Cefepime	1-2g q8-12h	100% of normal dose q12h	50-100% of normal dose q24h	25-50% of normal dose q24h	HD: 1g q24h (+ extra 1g post HD)	For continuous infusion, start with LD of 15mg/kg IV over 30 minutes and then immediately begin the continuous infusion:										

ANTIMICROBIAL	DOSE FOR NORMAL RENAL FUNCTION	ADJUSTMENT FOR RENAL FAILURE Estimated Creatinine Clearance (CrCl), mL/min			DIALYSIS DOSE	COMMENTS								
		> 50	10-50	<10										
						<table><tr><td>CrCl</td><td>Dose</td></tr><tr><td>>60</td><td>6g q24h</td></tr><tr><td>30-60</td><td>4g q24h</td></tr><tr><td>11-29</td><td>2g q24h</td></tr></table>	CrCl	Dose	>60	6g q24h	30-60	4g q24h	11-29	2g q24h
CrCl	Dose													
>60	6g q24h													
30-60	4g q24h													
11-29	2g q24h													
Cefotaxime	2g q8h	100% of normal dose q8-12h	100% of normal dose q12-24h	100% of normal dose q24h	HD: 2g q24h									
Cefoperazone	1-2g q12h	100% of normal dose q12h	100% of normal dose q12h	100% of normal dose q12h		Max dose 4g/day for patients who CrCl < 18mL/min or serum creatinine > 309µmol/L								
Ceftazidime	2g q8h	100% of normal dose q8-12h	100% of normal dose q12-24h	100% of normal dose q24-48h	HD: 2g q24-48h	For continuous infusion, start with LD of 15mg/kg IV over 30 minutes and then immediately begin the continuous infusion: <table><tr><td>CrCl</td><td>Dose</td></tr><tr><td>>60</td><td>6g q24h</td></tr><tr><td>30-60</td><td>4g q24h</td></tr><tr><td>11-29</td><td>2g q24h</td></tr></table>	CrCl	Dose	>60	6g q24h	30-60	4g q24h	11-29	2g q24h
CrCl	Dose													
>60	6g q24h													
30-60	4g q24h													
11-29	2g q24h													
Cefuroxime sodium (IV)	750-1500mg q8h	100% of normal dose q8h	10-30mL/min: 100% of normal dose q12h	100% of normal dose q24h	CAPD, HD: 750-1500mg q24h	In more serious infection, can increase to IV 1500mg q6h								

ANTIMICROBIAL	DOSE FOR NORMAL RENAL FUNCTION	ADJUSTMENT FOR RENAL FAILURE Estimated Creatinine Clearance (CrCl), mL/min			DIALYSIS DOSE	COMMENTS
		> 50	10-50	<10		
Cefuroxime axetil (PO)	250mg-500mg q12h	100% of normal dose q12h	10-30mL/min: 250mg q12h OR 250-500mg q24h	250mg q24h OR 250-500mg q48h	CAPD, HD: 250mg q24h OR 250-500mg q48h	
(D) Fluoroquinolones						
Ciprofloxacin (IV)	400mg q12h	100% of normal dose q12h	100% of normal dose q24h	100% of normal dose q24h OR 50% of normal dose q12h	HD: 400mg IV q24h	
Ciprofloxacin (PO)	500-750mg q12h	100% of normal dose q12h	30-50mL/min: 250-500mg q12hr 10-30mL/min: 500mg q24h	500mg q24h	CAPD, HD: 250-500mg q24h	
Levofloxacin (IV/PO)	750mg q24h	100% of normal dose q24h	20-49mL/min: 100% of normal dose q48h	< 20 mL/min: 750mg q24h then 500mg q48h	CAPD, HD: 750mg q24h then 500mg q48h	
Ofloxacin (PO)	200-400mg q12h	100% of normal dose q12h	100% of normal dose q24h	200mg q24h	HD: 200mg q24h	
(E) Macrolides						
Clarithromycin (PO)	500mg q12h	100% of normal dose q12h	50-100% of normal dose q12h	50-75% of normal dose q12h	CAPD, HD: 500mg q24h	

ANTIMICROBIAL	DOSE FOR NORMAL RENAL FUNCTION	ADJUSTMENT FOR RENAL FAILURE Estimated Creatinine Clearance (CrCl), mL/min			DIALYSIS DOSE	COMMENTS
		> 50	10-50	<10		
(F) Miscellaneous Antibacterials						
Colistin	LD (regardless of renal function): IBW > 75kg: 9 MU IBW 61-74kg: 8 MU IBW 51-60kg: 7 MU IBW < 50kg: 6 MU MD: Based on renal function	4.5 MU q12h	30-50mL/min: 3 MU q12h 10-30mL/min: 2.5 MU q12h	2 MU q12h	HD: 2 MU q12h (+ extra 1.5 MU as supplemental dose post HD)	
Linezolid (IV/PO)	600mg q12h	100% of normal dose q12h	100% of normal dose q12h	100% of normal dose q12h	CAPD: No dosage adjustment HD: No dose adjustment, but administer one of the dose post HD	
Metronidazole	250-500mg q8-12h	100% of normal dose q8-12h	100% of normal dose q8-12h	100% of normal dose q8-12h	CAPD, HD: 7.5mg/kg q12h	
Nitrofurantoin (PO)	UTI treatment: 50-100mg q6h UTI Prophylaxis: 50-100mg at night	Avoid use when CrCl < 60mL/min	Avoid use	Avoid use	Avoid use	

ANTIMICROBIAL	DOSE FOR NORMAL RENAL FUNCTION	ADJUSTMENT FOR RENAL FAILURE Estimated Creatinine Clearance (CrCl), mL/min			DIALYSIS DOSE	COMMENTS																								
		> 50	10-50	<10																										
Trimethoprim (TMP)/sulfamethoxazole (SMZ) (IV/PO) TMP 80mg: SMZ 4000mg	5-20mg/kg/day (TMP) (divided q6-12h)	100% of normal dose (divided q6-12h)	10-29mL/min: 5-10mg/kg/day (divided q12h)	5-10mg/kg q24h																										
Vancomycin	<table><tr><th>CrCl (mL/min)</th><th>Loading Dose (LD)</th><th>Initial Maintenance Dose (MD)</th><th>Dosing Interval</th></tr><tr><td>> 90</td><td>25-30mg/kg*</td><td>15-20mg/kg</td><td>q8-12h</td></tr><tr><td>50 - 90</td><td>20-25mg/kg</td><td>15-20mg/kg</td><td>q12h</td></tr><tr><td>20 - 49</td><td>20-25mg/kg</td><td>10-15mg/kg</td><td>q24h</td></tr><tr><td>< 20</td><td>20-25mg/kg</td><td colspan="2">Dosage adjustment according to TDM level</td></tr><tr><td>HD</td><td>25mg/kg</td><td colspan="2">Dosage adjustment according to TDM level</td></tr></table>					CrCl (mL/min)	Loading Dose (LD)	Initial Maintenance Dose (MD)	Dosing Interval	> 90	25-30mg/kg*	15-20mg/kg	q8-12h	50 - 90	20-25mg/kg	15-20mg/kg	q12h	20 - 49	20-25mg/kg	10-15mg/kg	q24h	< 20	20-25mg/kg	Dosage adjustment according to TDM level		HD	25mg/kg	Dosage adjustment according to TDM level		Maximum loading dose 1. Normal body weight: 2g/dose 2. Obese (BMI ≥ 30kg/m²): 3g/dose *Suggestion for 30mg/kg in seriously ill patients
CrCl (mL/min)	Loading Dose (LD)	Initial Maintenance Dose (MD)	Dosing Interval																											
> 90	25-30mg/kg*	15-20mg/kg	q8-12h																											
50 - 90	20-25mg/kg	15-20mg/kg	q12h																											
20 - 49	20-25mg/kg	10-15mg/kg	q24h																											
< 20	20-25mg/kg	Dosage adjustment according to TDM level																												
HD	25mg/kg	Dosage adjustment according to TDM level																												
(G) Penicillins																														
Amoxicillin (PO)	250-1000mg q8h	100% of normal dose q8h	100% of normal dose q8-12h	100% of normal dose q24h	HD: 250-500mg post HD																									
Ampicillin sodium	1-2g q4-6h	100% of normal dose q4-6h	30-50mL/min: 100% of normal dose q6-8h 10-30mL/min: 100% of normal dose q8-12h	100% of normal dose q12h	HD: 1-2g q12h																									

ANTIMICROBIAL	DOSE FOR NORMAL RENAL FUNCTION	ADJUSTMENT FOR RENAL FAILURE Estimated Creatinine Clearance (CrCl), mL/min			DIALYSIS DOSE	COMMENTS
		> 50	10-50	<10		
Amoxicillin/clavulanate (IV)	1.2g q8h In serious infection can increase frequency up to q6h	100% of normal dose q8h	10-30mL/min: 1.2g first dose then 600mg q12h	1.2g first dose then 600mg q12h	CAPD: No data HD: 1.2g first dose then 600mg q12h	
Amoxicillin/clavulanate (PO)	500/125mg q8h	100% of normal dose q8h	>30mL/min: 100% of normal dose q8h 10-30mL/min: 100% of normal dose q12h	100% of normal dose q24h	HD: 625mg q24h post HD	
Ampicillin/sulbactam (IV)	3g q6h	100% of normal dose q6h	15-29mL/min: 100% of normal dose q12h	< 15mL/min: 100% of normal dose q24h	CAPD: 3g q12h HD: 3g q24h post HD	
Benzylpenicillin	0.5-4MU q4-6h	100% of normal dose q4-6h	75% of normal dose q4-6h	20-50% of normal dose q4-6h	HD: Normal dose first dose then 25-50% of normal dose q4-6h OR 50-100% of normal dose q8-12h	
Piperacillin/tazobactam	3.375-4.5g q6h	100% of normal dose q6h	20-40mL/min: 2.25g-3.375g q6h <20mL/min: 2.25g-3.375g q8h	2.25g q6-8h	HD: 2.25g q8-12h	

ANTIMICROBIAL	DOSE FOR NORMAL RENAL FUNCTION	ADJUSTMENT FOR RENAL FAILURE Estimated Creatinine Clearance (CrCl), mL/min			DIALYSIS DOSE	COMMENTS
		> 50	10-50	<10		
ANTIFUNGAL						
Amphotericin	Amphotericin B Test dose 1mg: starting 0.1mg/kg/day over 6 hours; increase to target dose 0.25-1mg/kg q24h (over 2-6h)	100% of normal dose. No changes in frequency.	100% of normal dose. No changes in frequency.	100% of normal dose. No changes in frequency.	100% of normal dose. No changes in frequency	
	Amphotericin, colloidal dispersion 3-6mg/kg q24h	100% of normal dose. No changes in frequency.	100% of normal dose. No changes in frequency.	100% of normal dose. No changes in frequency.	100% of normal dose. No changes in frequency	
	Amphotericin, lipid complex 5mg/kg/day as single infusion	100% of normal dose. No changes in frequency.	100% of normal dose. No changes in frequency	100% of normal dose. No changes in frequency	100% of normal dose. No changes in frequency	
	Amphotericin, liposomal 1-5mg/kg/day as single infusion	100% of normal dose. No changes in frequency	100% of normal dose. No changes in frequency	100% of normal dose. No changes in frequency	100% of normal dose. No changes in frequency	
Fluconazole (IV/PO)	100-400mg q24h	100% of normal dose q24h	50% of normal dose q24h	50% of normal dose q24h	CAPD: 200mg then 50-100mg q24h HD: 100-400mg q24h	

ANTIMICROBIAL	DOSE FOR NORMAL RENAL FUNCTION	ADJUSTMENT FOR RENAL FAILURE Estimated Creatinine Clearance (CrCl), mL/min			DIALYSIS DOSE	COMMENTS
		> 50	10-50	<10		
Flucytosine	25mg/kg q6h	> 40mL/min: 100% of normal dose q6h	21-40mL/min: 100% of normal dose q12h 10-20mL/min: 100% of normal dose q24h	100 % of normal dose q48h	HD: 25-50mg/kg q48-72h post HD	
Itraconazole (IV)	200mg q12h for 4 doses, then 200mg q24h	100% of normal dose. No changes in frequency.	<30mL/min: Avoid	Avoid	Avoid	IV formulation is contraindicated for CrCl <30mL/min due to accumulation of cyclodextrin vehicle)
Itraconazole (PO)	100-200mg q12h	100% of normal dose q12h	100% of normal dose q12h	50% of normal dose q12h	CAPD, HD: 100mg q12-24h	
Voriconazole (IV)	6mg/kg q12h for 2 doses, then 4mg/kg q12h	100% of normal dose. No changes in frequency.	Switch to PO or suspension (IV formulation is contraindicated for CrCl <30mL/min due to accumulation of cyclodextrin vehicle)			
Voriconazole (PO)	>40kg: 400mg PO q12h x2 doses, then 200mg PO q12h <40kg: 200mg PO q12h x2 doses, then 100mg PO q12h	100% of normal dose. No changes in frequency.	100% of normal dose. No changes in frequency.	100% of normal dose. No changes in frequency.	CAPD, HD: No data, but prefer oral route to avoid accumulation of IV cyclodextrin vehicle	

ANTIMICROBIAL	DOSE FOR NORMAL RENAL FUNCTION	ADJUSTMENT FOR RENAL FAILURE Estimated Creatinine Clearance (CrCl), mL/min			DIALYSIS DOSE	COMMENTS
		> 50	10-50	<10		
ANTIPARASITIC						
Pentamidine (IV)	IV/IM 4mg/kg q24h	100% of normal dose q24h	100% of normal dose q24h	100% of normal dose q24-36h	CAPD: 4mg/kg q24-36h HD: 4mg/kg q48h	
ANTIMYCOBACTERIAL						
Ethambutol	15-25mg/kg q24h (max 1600mg/day)	100% of normal dose q24h	30-50mL/min: 100% normal dose q24-36h 15-25mL/min: 100% of normal dose q36-48h	15mg/kg q48h	CAPD, HD: 15mg/kg q48h	
Isoniazid (PO)	5mg/kg q24h (max 300mg/day)	100% of normal dose q24h	100% of normal dose q24h	100% of normal dose q24h	HD: 5mg/kg q24h	
Pyrazinamide (PO)	25mg/kg q24h (max 2500mg/day)	100% of normal dose q24h	10-20mL/min: 100% of normal dose q48h	100% of normal dose q48h	CAPD: 25mg/kg q24h HD: 25mg/kg q48h	
Rifampin (PO)	10mg/kg q24h (max 600mg/day)	100% of normal dose q24h	50-100% of normal dose q24h	50-100% of normal dose q24h	CAPD, HD: 300-600mg q24h	
Ethionamide (PO)	15-20mg/kg/day in 1-3 divided doses	100% of normal dose. No changes in frequency.	100% of normal dose. No changes in frequency.	50% of normal dose. No changes in frequency.	CAPD, HD: 50% of normal dose. No	

ANTIMICROBIAL	DOSE FOR NORMAL RENAL FUNCTION	ADJUSTMENT FOR RENAL FAILURE Estimated Creatinine Clearance (CrCl), mL/min			DIALYSIS DOSE	COMMENTS
		> 50	10-50	<10		
	(max 1000mg/day)				changes in frequency.	
ANTIVIRAL						
Acyclovir	5-12.5mg/kg q8h	100% of normal dose q8h	100% of normal dose q12-24h	5-6.25mg/kg q24h	CAPD, HD: 2.5-6.25mg/kg q24h	
Ganciclovir (IV)	Induction: 5mg/kg q12h Maintenance: 5mg/kg q24h OR 6mg/kg 5 times weekly	50-69mL/min: 50% of normal dose q12h	25-49mL/min: 50% of normal dose q24h 10-24mL/min: 25% of normal dose q24h	25% of normal dose 3x/week	CAPD, HD: 1.25mg/kg 3x/week	
Valganciclovir (PO)	Induction: 900mg q12h	≥60mL/min: 900mg q12h 40-59mL/min: 450mg q12h 25-39mL/min: 450mg q24h 10-24mL/min: 450mg q48h		Avoid use		
	Maintenance (prophylaxis): 900mg q24h	≥60mL/min: 900mg q24h 40-59mL/min: 450mg q24h 25-39mL/min: 450mg q48h 10-24mL/min: 450mg twice weekly		Avoid use		
Lamivudine (HIV) (PO)	300mg q24h	100% of normal dose q24h	300mg first dose; then 50-150mg q24h	50mg first dose; then 25-50mg q24h	CAPD, HD: 25-50mg q24h	

ANTIMICROBIAL	DOSE FOR NORMAL RENAL FUNCTION	ADJUSTMENT FOR RENAL FAILURE Estimated Creatinine Clearance (CrCl), mL/min			DIALYSIS DOSE	COMMENTS
		> 50	10-50	<10		
Zidovudine	300mg q12h	100% of normal dose q12h	100% of normal dose q12h	100% of normal dose q24h or 100mg q8h	HD: 100mg q8h or 300mg q24h CAPD: 100mg q8h	

HD = Hemodialysis; CAPD = Continuous Peritoneal Dialysis; LD = Loading Dose; MD: Maintenance Dose; IV = Intravenous; IM = Intramuscular; PO = Oral

References:

1. The Sanford Guide to Antimicrobial Therapy 2022.
2. The Renal Drug Handbook 5th Edition.
3. ISPD peritonitis guideline recommendations: 2022 update on prevention and treatment.
4. Guide to Antimicrobial Therapy in the Adult ICU 2017.
5. Micromedex (online).
6. Lexicomp (online).

APPENDIX 4: CLINICAL PHARMACOKINETICS GUIDELINES (AMINOGLYCOSIDES AND VANCOMYCIN)

I. AMINOGLYCOSIDE DOSING STRATEGIES

A. EXTENDED-INTERVAL THERAPY/SINGLE DAILY DOSING (EID/SDD)

EID/SDD is an approach of giving high-dose aminoglycoside over 30 minutes at an extended interval (e.g 24 hourly, 36 hourly or more).

The theoretical benefits of EID/SDD:

- Aminoglycosides display concentration-dependent bactericidal action - that is, higher dose and serum concentrations result in more rapid bacterial killing.¹
- Optimize concentration-dependent bacterial killing by achieving a high peak (>10x MIC).²
- Minimize nephrotoxicity by administering larger, less frequent doses and potentially decreasing renal cortical aminoglycoside concentrations.
- Utilize the post-antibiotic effect (PAE) (2-8 hours), defined as a recovery period before organisms can resume growth after drug removal.¹
- Minimize the development of adaptive resistance by allowing a recovery period during the dosing interval.

Exclusion criteria:

EID/SDD is reasonable in most patients, with the following exceptions:³

- Pregnancy
- Ascites
- Burns (>20%)
- Endocarditis
- Creatinine clearance <30ml/min
- Dialysis
- Neutropenic patients
- Patients with gram positive infections (synergistic effect).
- Hemodynamically unstable.
- History of hearing loss/vestibular dysfunction.
- Mycobacterium infection.

SDD Dosing Strategy Based on Creatinine Clearance:⁴

Creatinine Clearance (mL/min)	Dose in 24 hours	
	Gentamicin	Amikacin
> 80	5mg/kg	15mg/kg
60 - 79	4mg/kg	12mg/kg
40 – 59	3.5mg/kg	7.5mg/kg

30 – 39	2.5mg/kg	4mg/kg
< 30	Conventional dosing	Conventional dosing

EID Dosing Strategy Based on Serum Concentration:⁵

Gentamicin	Amikacin
7mg/kg per dose	15mg/kg per dose


1. Initial level monitoring*

Single level drawn 8-12 hours after the first dose (Only applicable for 7mg/kg – plotting doses lower or higher than 7mg/kg may under or overestimate clearance)

Concentration Gentamicin (7mg/kg/dose): Plot level on graph

Concentration Amikacin (15mg/kg/dose): Divide level in half, then, plot on graph

****Please consult pharmacists for dosage adjustment.***



2. Follow up trough level

Trough monitoring (30-60 minutes prior to dose) should be considered in patients demonstrating acute changes in renal function or suspicion of extended interval failure.

Maintenance trough levels should be monitored at least once weekly.

Sample Parameters	Gentamicin	Amikacin
-------------------	------------	----------

Time to sample ¹⁰	At the 2 nd dose			
Sampling time ¹⁰	Take two samples at minimum 4 hours interval (e.g. post-2H and post-6H)			
Target levels (mcg/ml) ^{5,6}	TROUGH	PEAK*	TROUGH	PEAK*
	Adult: < 1 mcg/ml Children: < 1 mcg/ml IP: < 3 mcg/ml	Adult: 10-20 mcg/ml Children: 5-12 mcg/ml	Adult: < 5 mcg/ml Children: < 5 mcg/ml IP: < 15 mcg/ml	Adult: 20-60 mcg/ml Children: 20-30 mcg/ml

*The target reference range may be individualized based on institutional MIC value.

B. CONVENTIONAL/TRADITIONAL DOSING

Traditional dosing includes reduced doses and frequent administration of aminoglycosides using pharmacokinetic parameters to determine dose and frequency. However, in practice, dose below 7mg/kg for Gentamicin at 24 hours interval is frequently used for patients with normal renal function.

Creatinine Clearance (ml/min)	Gentamicin	Amikacin
> 60 ⁷	1.5 – 2mg/kg every 8 hourly	5 – 7.5mg/kg every 8 hourly
40 - 60 ⁷	1.5 – 2mg/kg every 12 hourly	5 – 7.5mg/kg every 12 hourly
20 - 40 ⁷	1.5 – 2mg/kg every 24 hourly	5 – 7.5mg/kg every 24 hourly
< 20 ⁷	1.5 – 2mg/kg every 48 – 72 hourly	5 – 7.5mg/kg every 48 – 72 hourly
CVVH / CVVHD / CVVHDF ⁸	Loading dose 3mg/kg followed by 2mg/kg every 24 – 48 hourly	Loading dose 10mg/kg followed by 7.5mg/kg every 24 – 48 hourly
CAPD ⁹	Intermittent: 0.6mg/kg in night dwell Continuous: Loading dose 8mg/L followed by 4mg/L	Intermittent: 2mg/kg in night dwell Continuous: Loading dose 25mg/L followed by 12mg/L

Sample Parameters	Gentamicin	Amikacin
Time to sample ¹⁰	After the 3 rd dose	
Sampling time ¹⁰	PRE: obtained just prior to the next dose OR within 30 minutes before the next dose POST: 30 minutes after completion of 30 minutes infusion OR	

	Bolus: 1 hour after dose is given			
Sampling time for ESRF¹¹	PRE dialysis			
Target levels (mcg/ml)^{6,10}	TROUGH	PEAK*	TROUGH	PEAK*
	< 2	5 - 10	< 10	20 - 30

**The target reference range may be individualized based on institutional MIC value.*

References:

1. Bennett WM, Plamp CE, Gilbert DN, Parker RA, Porter GA. The influence of dosage regimen on experimental gentamicin nephrotoxicity: dissociation of peak serum levels from renal failure. J Infect Dis 1979; 140:576-580
2. Freeman C.D. et al. Once daily dosing of aminoglycosides: review and recommendations for clinical practice. Journal of antimicrobial chemotherapy (1997) 39, 677-686
3. Once-Daily Dosing of Aminoglycosides, A consensus Document. 1997. Nasr Anaizi. Obtained from www.rxkinetics.com/oda.html on 23 May 2011
4. The Sanford Guide to Antimicrobial Therapy 2014 (Forty-fourth edition)
5. Hartford Hospital Once Daily Aminoglycoside nomogram.
6. Bauer LA 2006.Clinical Pharmacokinetics Handbook. Chapter 4: Aminoglycoside Antibiotics. New York. McGraw Hill.
7. Guide to Antimicrobial Therapy in the Adult ICU 2012.
8. Robin T et.al. Antibiotic Dosing in Critically Ill Adult Patients Receiving Continuous Renal Replacement Therapy. Clinical Infectious Disease 2005;41:1159-1166
9. ISPD Guidelines/Recommendation-Peritoneal Dialysis-Related Infections Recommendations: 2010 Update. Peritoneal Dialysis International, Vol 30,402.
10. Winter ME.2010.Basic Clinical Pharmacokinetics 5th Edition. Philadelphia. Lippincott Williams and Wilkins.
11. Clinical Pharmacokinetics: Pharmacy Handbook 1st Edition.

II. THERAPEUTIC DRUG MONITORING PROTOCOL FOR EXTENDED-INTERVAL GENTAMICIN IN NEONATES

Gentamicin initial empiric dosage regimen is 5mg/kg^{1,2} Q36H

Sampling time: post 22 hours², or 30 minutes prior second dose^{1,3}

Nomogram for sampling post 22 hours²

Suggested Gentamicin Dosing Interval Based on 22-Hour Concentration ²	
Concentration at 22 hours, mg/L	Suggested Dosing Regime,
<1.2	5mg/kg, Q24H
1.3-2.6	5mg/kg, Q36H
2.7-3.5	5mg/kg, Q48H
>3.6	Hold dose, repeat concentration in 24 hours

Nomogram for sampling 30 minutes prior next dose^{1,3}

1. Therapeutic range:

Cmin: <2mg/L (pre-dose of second dose)

Cmin: <1mg/L (if more than 3 doses of gentamicin)

2. If Cmin ABOVE therapeutic range, repeat sampling by **ADDITIONAL 12 HOURS INTERVAL** from the last dose.

Empirical interval	Repeat sampling
Q36H	48 hours from last dose

3. Recommendation of Dosing interval must be based on the time to withhold until Cmin in therapeutic range.

4. If intended Cmin is not available (eg. sampling error), **DO NOT WITHHOLD** the next dose unless there is evidence of renal dysfunction (for example: an elevated serum urea or creatinine concentration or anuria).

5. Repeat sampling EVERY THIRD DOSE (eg. 2nd dose, 5th dose, 8th dose etc).

Cmax¹ (1 hour after dose administered)

- Aim Cmax>8mg/L if neonate has Gram-negative or staphylococcal infection

Suggest Cmax sampling if:

1. Neonate with edema
2. Macrosomia (birthweight more than 4.5kg)
3. Unsatisfactory response to treatment
4. Proven Gram-negative infection.

Note:

1. Aim Cmax : MIC³ = 8:1 to 10:1, an AUC 80-100mg⁻¹L⁻¹
2. PAC= 4 hours after undetectable level of Cmin (<0.3mg/L) before redosing⁵.
3. Potential advantage of Extended interval **similar** safety to traditional dosing.⁴
4. Serum creatinine **elevations by ≥ 50% from baseline** should investigate gentamicin induced nephrotoxicity.³
5. Demographic distribution for 5mg/kg Q36h

	Group I (≤ 34 6/7 weeks of GA)			Group II (> 35 0/7 weeks of GA)		
	Control arm	Study arm	P	Control arm	Study arm	P
N	25	25		23	23	
Male, n (%)	16 (64%)	12 (48%)	0.4	9 (39%)	12 (52%)	0.6
GA, weeks (range)	30.6 ± 3.1 ^a (24–34)	29 ± 3.7 (24–34)	0.3	38.1 ± 2 (35–41)	38 ± 1.7 (35–41)	0.7
PNA, days (range)	3.4 ± 6.3 (1–26)	6 ± 8.8 ^a (1–32)	0.09	1 ± 0.2 (1–2)	1.4 ± 0.6 (1–3)	0.01
PCA, weeks (range)	31.1 ± 2.7 (26–34.14)	30.1 ± 3.3 (24.14–34.14)	0.4	38.3 ± 2 (35.14–41.14)	38.2 ± 1.8 (35.14–41.14)	0.95
Birth weight, kg (range)	1.6 ± 0.6 (0.53–2.8)	1.4 ± 0.7 (0.59–2.98)	0.6	3.2 ± 0.4 (2.31–3.86)	3.2 ± 0.5 (2.37–4.057)	0.79

Abbreviations: GA, gestational age; PCA, postconceptional age; PNA, postnatal age. ^aMean ± s.d.

6. Population pharmacokinetic profile

PK data	Group I (≤ 34 6/7 weeks of GA)			Group II (> 35 0/7 weeks of GA)		
	Control arm, n = 25	Study arm, n = 25	P	Control arm, n = 23	Study arm, n = 23	P
K _e	0.074 ± 0.02 ^a	0.076 ± 0.02	0.83	0.093 ± 0.02	0.093 ± 0.02	0.629
t _{1/2}	9.9 ± 2.4	9.8 ± 2.6	0.83	7.7 ± 1.4	7.7 ± 1.6	0.829
V _d (l kg ⁻¹)	0.5 ± 0.1	0.6 ± 0.1	0.69	0.5 ± 0.1	0.48 ± 0.1	0.688
Cl (l h ⁻¹ kg ⁻¹)	0.06 ± 0.02	0.06 ± 0.03	0.72	0.15 ± 0.04	0.14 ± 0.04	0.718

Abbreviations: Cl, clearance; GA, gestational age; K_e, elimination rate constant; t_{1/2}, half-life; V_d, volume of distribution. ^aMean ± s.d.

References:

1. NICE Guidance, 2020. Neonatal Infection (early onset): antibiotics Treatment and Monitoring for Suspected or Confirmed Early-Onset Neonatal Infection.
2. Deonne, D, Akierman, A., Alshaikh, B. & Yusuf, K. 2012. Validation of a Dosage Individualization Table for Extended-Interval Gentamicin in Neonates. Neonatology (46) 935-942
3. El-Chaar, G.M., Supaswud-Franks, T., Venugopalan, L., Kohn, N. & Castro-Alcaraz, S. 2016. Extended-interval gentamicin administration in neonates: a simplified approach. Journal of Perinatology. 1-6.
4. Rao, S.C., Srinivasjois, R. Hagan, R. & Ahmed, M. 2011. One dose per day compare to multiple doses per day of gentamicin for treatment of suspected or proven sepsis in neonate. Cochrane Database syst Rev. 11: CD005091.
5. Zhanel, G.G., Hoban, D.J., Harding, G.K.M. 1991. The postantibiotic effect: a review of in vitro and in vivo data. DICP. 25(2): 153-163.

III. THERAPEUTIC DRUG MONITORING PROTOCOL FOR VANCOMYCIN FOR PATIENTS WITH COMPLICATED INVASIVE MRSA INFECTION ONLY*

*Defined as “including but not limited to bacteremia, sepsis, infective endocarditis, pneumonia, osteomyelitis and meningitis”¹

Intermittent Infusion ^{2,3}:

- Loading dose (using actual body weight, irrespective renal function)
 - Normal body weight : 20-35mg/kg over 1-2 hours (max: 3g/dose)
 - Obesity (BMI≥30kg/m²) : 20-25mg/kg (max: 3g/dose)
- Maintenance dose (max: 4.5g/day):
 - 15-20mg/kg/dose frequency according to table below ³:

CrCL (mL/min)	Dosage Adjustment
>50	bd
20-49	od
<20	Dose per Vancomycin's level

SPECIAL POPULATION EMPIRIC DOSING REGIMEN ^{2,3}:

Critically ill:

- Continuous Infusion:
 - Loading dose : 15-20mg/kg over 1-2 hours
 - Maintenance dose : 30-40mg/kg/day (max: 4.5g/day)

Unstable Pharmacokinetic Parameter (AKI, Burn, Augmented renal function)

- Loading dose (using actual body weight)
 - Normal body weight: 20-35mg/kg over 1-2 hours (max: 3g/dose)
- Maintenance dose:
 - 15-20mg/kg/dose frequency according to table below ³:

CrCL (mL/min)	Dosage Adjustment
>50	bd
20-49	od
<20	Dose per Vancomycin's level

Hemodialysis:

- CRRT (effluent rate: 20-25ml/kg/h):
 - Loading dose: 20-25mg/kg
 - Maintenance dose: 7.5mg-10mg/kg/dose bd

- Hybrid (SLED)
 - Loading dose: 20-25mg/kg
 - 15mg/kg/ dose (dose per level)
- Intermittent Hemodialysis (thrice weekly dose administration)
 - Post HD (after dialysis end)
 - Low Flux:
 - Loading dose: 25mg/kg
 - Maintenance: 7.5mg/kg
 - High Flux:
 - Loading dose: 25mg/kg
 - Maintenance: 10mg/kg
 - Intradialytic (Last hour of HD)
 - Low Flux
 - Loading dose: 30mg/kg
 - Maintenance: 7.5-10mg/kg
 - High Flux
 - Loading dose: 35mg/kg
 - Maintenance: 10-15mg/kg

EFFICACY PARAMETER ^{2,3}

AUC₂₄/MIC: 400-600 mg.h/L

- Assume MIC=1 if MIC is not available
- AUC calculated based on past 24hours accumulation dose
- Must be at steady state

Cmin: 15-20mg/L

- For Intermittent Hemodialysis only
- Noninvasive OR uncomplicated MRSA OR non-MRSA organism

TOXICODYNAMIC PARAMETERS ²

- AUC₂₄ >600mg.h/L
- Increase SCr in 23mcmol/L from baseline (according to AKIN, KDIGO criteria).

DILUTION AND ADMINISTRATION ^{3,4}

- Strength/unit: 500mg
- Reconstitution: 10ml Water for Injection
- Dilution for infusion: Normal saline / D5
- Rate of administration: 10-15mg/min
- Max concentration:
 - Non fluid restriction: ≤ 5mg/ml
 - Fluid restriction: ≤10mg/ml
- Stability:
 - Reconstituted: 14 days
 - Diluted:
 - Under refrigeration: 14 days

- Room temperature: 7 days

References:

1. CADTH Technology Review: Vancomycin Therapeutic Drug Monitoring for Serious Methicillin-Resistant *Staphylococcus aureus* infections.2020. Cuello,C. & Loshak, H.Canadian Agency of Drugs and Technologies in Health (CADTH).
2. Therapeutic monitoring of vancomycin for serious methicillin-resistant *Staphylococcus aureus* infections: A revised consensus guideline and review by the American Society of Health-System Pharmacists, the Infectious Diseases Society of America, the Pediatric Infectious Diseases Society and the Society of Infectious Diseases Pharmacist. 2020. ASHP Report.
3. Clinical Pharmacokinetic Pharmacy Handbook, 2019. Clinical Pharmacy Working Committee (Clinical Pharmacokinetics Subspecialty) Pharmacy Practice & Development division. Ministry of Health Malaysia.
4. Injectable Drugs Dilution, 2010. Bahagian Perkhidmatan Farmasi, Jabatan Kesihatan Negeri Perak.

